

Strong invariance and noise-comparison principles for some parabolic stochastic PDEs*

Mathew Joseph
University of Sheffield

Davar Khoshnevisan
University of Utah

Carl Mueller
University of Rochester

January 27, 2015

Abstract

We consider a system of interacting diffusions on the integer lattice. By letting the mesh size go to zero and by using a suitable scaling, we show that the system converges (in a strong sense) to a solution of the stochastic heat equation on the real line. As a consequence, we obtain comparison inequalities for product moments of the stochastic heat equation with different nonlinearities.

Keywords: Stochastic PDEs, comparison theorems, white noise.

AMS 2010 subject classification: Primary 60H15; Secondary: 35K57.

1 Introduction

We propose to study the solution $u := \{u_t(x)\}_{t \geq 0, x \in \mathbf{R}}$ to the following partial differential equation:¹

$$\frac{\partial}{\partial t} u_t(x) = (\mathcal{L}u_t)(x) + \sigma(u_t(x))\xi(t, x) \quad (t > 0, x \in \mathbf{R}); \quad (\text{SHE})$$

subject to $u_0(x) := 1$ for all $x \in \mathbf{R}$. The function $\sigma : \mathbf{R} \rightarrow \mathbf{R}$ is assumed to be non-random and globally Lipschitz continuous. The forcing term ξ in (SHE) is *space-time white noise*. That is, ξ is assumed to be a generalized Gaussian random field on $\mathbf{R}_+ \times \mathbf{R}$ with mean function zero and covariance measure

$$\text{Cov}(\xi(t, x), \xi(s, y)) = \delta_0(t - s)\delta_0(x - y). \quad (1.1)$$

*Research supported in part by the United States National Science Foundation grants DMS-0747758 (M.J.), DMS-1307470 (M.J. and D.K.), and DMS-1102646. (C.M.).

¹Note that $u_t(x)$ denotes the evaluation of the solution at (t, x) ; this is consistent with the standard nomenclature of stochastic process theory. To be consistent, we will never write u_t in place of the time derivative $\partial_t u$ of u .

Finally, the operator \mathcal{L} acts on the variable x only, and is the fractional Laplacian, on \mathbf{R} , of order α . That is,

$$\mathcal{L} = -\nu(-\Delta)^{\alpha/2}, \quad (1.2)$$

where $\nu > 0$ is a parameter that is held fixed throughout this paper [ν is the so-called ‘‘viscosity coefficient’’].

We are interested only in the cases where the semigroup $t \mapsto \exp(-t\mathcal{L})$ is positive and conservative; that is precisely when the fundamental solution to the heat operator $(\partial/\partial t) - \mathcal{L}$ is a probability density at all times. It is well known that this property holds if and only if $\alpha \in (0, 2]$. Moreover, in that case, \mathcal{L} is a negative-definite pseudo-differential operator with Fourier symbol $\hat{\mathcal{L}}(z) = -\nu|z|^\alpha$, where our Fourier transform is normalized so that

$$\hat{f}(z) = \int_{-\infty}^{\infty} e^{izx} f(x) dx \quad \text{for all } z \in \mathbf{R} \text{ and } f \in L^1(\mathbf{R}). \quad (1.3)$$

In probabilistic terms, the preceding means that we can identify \mathcal{L} as the generator of a symmetric α -stable Lévy process $\{S_t\}_{t \geq 0}$ whose characteristic function at time $t \geq 0$ is normalized as

$$\mathbf{E} \exp(izS_t) = \exp(-\nu t|z|^\alpha) \quad (t \geq 0, z \in \mathbf{R}). \quad (1.4)$$

Even though $\alpha \in (0, 2]$ always leads us to a nice pseudo-differential operator \mathcal{L} , we will further assume that

$$1 < \alpha \leq 2. \quad (1.5)$$

It is well known that under the condition (1.5), (SHE) has an almost surely unique Hölder-continuous solution u that is predictable with respect to the filtration of the white noise ξ and satisfies

$$\sup_{t \in [0, T]} \sup_{x \in \mathbf{R}} \mathbf{E} (|u_t(x)|^k) < \infty \quad \text{for all } T > 0 \text{ and } k \geq 2. \quad (1.6)$$

See, for example, Dalang [6] and Peszat and Zabczyk [22], and also Foondun and Khoshnevisan [9] for proofs of these assertions. Moreover, if $\alpha \in (0, 1]$ and $\sigma(u) = 1$, then (SHE) does not have a function solution [6, 22].

We may think of $u_t(x)$ as the conditional expected density of a branching particle system at space-time point (t, x) given the white noise ξ , where the particles move in \mathbf{R} as an α -stable Lévy process. Moreover, these particles interact with one another only through the random environment ξ and the function σ ; the latter describes the underlying [non-linear] branching mechanism.

Let $\bar{\sigma} : \mathbf{R} \rightarrow \mathbf{R}$ be a non random Lipschitz continuous function and suppose \bar{u} solves (SHE) but with σ replaced by $\bar{\sigma}$; that is,

$$\frac{\partial}{\partial t} \bar{u}_t(x) = (\mathcal{L}\bar{u}_t)(x) + \bar{\sigma}(\bar{u}_t(x))\xi(t, x) \quad (t > 0, x \in \mathbf{R}), \quad (1.7)$$

subject to the same initial condition as u was; namely, $\bar{u}_0(x) := 1$ for all $x \in \mathbf{R}$. One of the goals of this project is to prove that if $\sigma \leq \bar{\sigma}$ pointwise, then frequently,

$$\mathbb{E} \left[\prod_{i=1}^m u_t(x_i) \right] \leq \mathbb{E} \left[\prod_{i=1}^m \bar{u}_t(x_i) \right], \quad (1.8)$$

for all $t > 0$ and $x_1, \dots, x_m \in \mathbf{R}$. We will conclude this fact—see Theorem 2.6—from a general approximation result that we describe next.

Let \mathcal{L} denote the generator of a rate 1, continuous-time random walk on \mathbf{Z} . That is, there is a probability measure q on \mathbf{Z} such that

$$(\mathcal{L}f)(m) := \sum_n q(n-m)[f(n) - f(m)], \quad (1.9)$$

for appropriate functions $f : \mathbf{Z} \rightarrow \mathbf{R}$. Consider the interacting particle system

$$dU_t(x) = (\mathcal{L}U_t)(x) dt + \sigma(U_t(x)) dB_t(x) \quad (t > 0, x \in \mathbf{Z}), \quad (\text{SHE}_1)$$

subject to $U_0(x) \equiv 1$ for all $x \in \mathbf{Z}$. Here, the jump rate of the underlying walk is assumed to be one, and $\tilde{\xi}(t, x) := dB_t(x)/dt$ denotes space-time white noise on $\mathbf{R}_+ \times \mathbf{Z}$; equivalently, $\{B_\bullet(x)\}_{x \in \mathbf{Z}}$ is a countable family of independent standard Brownian motions on \mathbf{R} , and (SHE_1) is understood as an infinite system of Itô stochastic differential equations in the sense of Shiga and Shimizu [23] [see §2.3 below for more details].

One might expect that *under appropriate conditions on the generator \mathcal{L}* , if we first rescale space in (SHE_1) so that $x \in \varepsilon\mathbf{Z}$ instead of $x \in \mathbf{Z}$, and then speed up the rate of the jumps of the underlying walks suitably, then the resulting rescaled version of (SHE_1) converges to the continuum equation (SHE) as $\varepsilon \downarrow 0$. This is an example of [an infinite dimensional] invariance principle.

This program has been carried out successfully by Funaki [10] in the special case that \mathcal{L} denotes the generator of a simple random walk, where $\nu = 1$ and \mathbf{R} is replaced by $[0, 1]$ in (SHE).

Our main contribution includes an extension of Funaki's program in two directions: Firstly, we study the full equation on \mathbf{R} , in place of $[0, 1]$, where compactness problems arise in noteworthy ways. Secondly, we establish an invariance principle of a sort that we will explain next.

In order to describe our invariance result, let π_ε denote the rescaling that maps the parameter set $\varepsilon\mathbf{Z}$ to the rescaled parameter set \mathbf{Z} . That is,

$$(\pi_\varepsilon f)(x) := f(\varepsilon x) \quad (x \in \mathbf{Z}, \varepsilon \in (0, 1]), \quad (1.10)$$

for all functions $f : \varepsilon\mathbf{Z} \rightarrow \mathbf{R}$. Now we consider the semi-discrete stochastic heat equation (SHE_1) , rescaled so that $x \in \varepsilon\mathbf{Z}$. That is, we study the solution $U^{(\varepsilon)}$ to the following infinite system of stochastic differential equations: For $t > 0$ and $x \in \varepsilon\mathbf{Z}$,

$$dU_t^{(\varepsilon)}(x) = \varrho_\varepsilon \times \left(\mathcal{L} \left[\pi_\varepsilon U_t^{(\varepsilon)} \right] \right) (x/\varepsilon) dt + \frac{1}{\sqrt{\varepsilon}} \sigma \left(U_t^{(\varepsilon)}(x) \right) dB_t(x/\varepsilon), \quad (\text{SHE}_\varepsilon)$$

subject to $U_0^{(\varepsilon)}(x) \equiv 1$ for all $x \in \varepsilon\mathbf{Z}$, where $\varrho_\varepsilon > 0$ is a suitable normalization.

As it turns out, the following is a good choice for our normalization factor:

$$\varrho_\varepsilon := \varepsilon^{-\alpha} \quad (0 < \varepsilon \leq 1). \quad (1.11)$$

[In fact, this factor is, in a sense, uniquely optimal.] Our choice for ϱ_ε is justified by the following corollary to our main theorem Theorem 2.4, stated in Section 2.4.

Corollary 1.1 (A weak invariance principle). *Suppose \mathcal{L} is the generator of a symmetric random walk $\{X_t\}_{t \geq 0}$ on \mathbf{Z} whose characteristic function satisfies Assumption 2.1 below. Then, the finite-dimensional distributions of $U^{(\varepsilon)}$ —normalized via (1.11)—converge to those of u as $\varepsilon \downarrow 0$.*

In fact, there is a very strong sense in which “functional weak convergence” takes place; see the strong approximation assertion—in the spirit of Csörgő and Révész [5]—of Theorem 2.5 below.

In due time, we will show also that Corollary 1.1 has a number of consequences in the analysis of the solution to the [continuous] stochastic partial differential equation (SHE); see, for example: (i) Theorem 2.5, which is a strong approximation theorem; (ii) Theorem 2.6, which is a comparison principle for the diffusion coefficient σ of (SHE); and (iii) Theorem 2.7, which establishes nearly-optimal bounds on the moment Lyapunov exponents of the solution to (SHE).

Our methods are connected loosely to earlier ideas from the literature on the numerical analysis of SPDEs. See, in particular, Gyöngy and Krylov [14, 15, 16].

2 Preliminaries, and main results

In this section we review some basic facts about SPDEs, mainly in order to state the notation, and then list the main contributions of the paper.

2.1 White noise and integration

It is well-known, and can be verified directly, that our space-time white noise can be realized as the distributional mixed derivative,

$$\xi(t, x) = \frac{\partial^2 W(t, x)}{\partial t \partial x}, \quad (2.1)$$

of a *two-sided Brownian sheet* $W := \{W(t, x)\}_{t \geq 0, x \in \mathbf{R}}$. The latter object is a mean-zero Gaussian process whose covariance function is

$$\text{Cov}(W(t, x), W(s, y)) = \min(s, t) \times \min(|x|, |y|) \times \mathbf{1}_{(0, \infty)}(xy), \quad (2.2)$$

for all $s, t \geq 0$ and $x, y \in \mathbf{R}$.

Standard results [25, Chapter 1] imply that W has Hölder-continuous sample functions. Since $W_t := W(t, \bullet)$ is an infinite-dimensional Brownian motion, [with values say in $C(\mathbf{R}, \mathbf{R})$], we may then define the Walsh integral process

$$M_t := \int_{(0,t) \times \mathbf{R}} Z_s(y) W(ds dy) \quad (t \geq 0), \quad (2.3)$$

in the spirit of an Itô integral, as a continuous martingale with quadratic variation

$$\langle M \rangle_t = \int_0^t ds \int_{-\infty}^{\infty} dy [Z_s(y)]^2 \quad (t \geq 0), \quad (2.4)$$

provided that $Z := \{Z_t(\bullet)\}_{t \geq 0}$ is predictable with respect to the Brownian filtration generated by $\{W_t\}_{t \geq 0}$, and the preceding displayed integral is finite for all $t \geq 0$. See Walsh [25, Chapter 2]. We mention the following variation of the Burkholder–Davis–Gundy inequality (Theorem 1.1 in [8], also see Remark 2.2 in [9]): For every $k \geq 2$ and $t \geq 0$,

$$\|M_t\|_k^2 \leq (4k)^{1/k} \int_0^t ds \int_{-\infty}^{\infty} dy \|Z_s(y)\|_k^2, \quad (\text{BDG})$$

where, here and throughout, we adopt the standard notation,

$$\|X\|_\nu := \{\mathbf{E}(|X|^\nu)\}^{1/\nu}, \quad (2.5)$$

valid for every $\nu \in [1, \infty)$ and $X \in L^\nu(\Omega)$.

2.2 The mild form of (SHE)

Define $p_t(x)$ to be the transition density of a symmetric α -stable Lévy process. In light of (1.4), this means that

$$\int_{-\infty}^{\infty} e^{ixz} p_t(x) dx = \exp(-\nu t |z|^\alpha) \quad (t > 0, z \in \mathbf{R}). \quad (2.6)$$

Because $(s, t, x, y) \mapsto p_{t-s}(y-x)$ is the fundamental solution to the heat operator $(\partial/\partial t) - \mathcal{L}$, the theory of Walsh [25, Chapter 3] tells us that we may write (SHE) in mild form as follows:

$$u_t(x) = 1 + \int_{(0,t) \times \mathbf{R}} p_{t-s}(y-x) \sigma(u_s(y)) W(ds dy), \quad (2.7)$$

for all $t > 0$ and $x \in \mathbf{R}$. The integral is understood to be a stochastic Itô-type integral in the sense of Walsh [25, Chapter 2].

2.3 The mild form of (SHE $_\varepsilon$)

Before we write (SHE $_\varepsilon$) in mild form, it might help to recall a few facts about continuous-time random walks. We use this opportunity to also set forth some notation.

Recall that $X := \{X_t\}_{t \geq 0}$ denotes a rate-one symmetric continuous-time random walk on \mathbf{Z} , whose generator is \mathcal{L} and satisfies $X_0 = 0$. [We further assume that the random walk satisfies the technical Assumption 2.1 below. But that assumption is not germane to the present discussion, and so will be introduced later on.]

Let us define, for every fixed $\varepsilon > 0$, a random walk $X^{(\varepsilon)} := \{X_t^{(\varepsilon)}\}_{t \geq 0}$ on the rescaled lattice $\varepsilon\mathbf{Z}$ by setting

$$X_t^{(\varepsilon)} := \varepsilon X_{t/\varepsilon^\alpha} \quad (t \geq 0). \quad (2.8)$$

We emphasize that $X_0^{(\varepsilon)} = 0$ and $\mathbb{E}X_t^{(\varepsilon)} = 0$ for all $t \geq 0$. Thus, the random walk X is rescaled to $X^{(\varepsilon)}$, so that $X^{(\varepsilon)}$ might possibly converge weakly to an α -stable Lévy process S —see (1.4)—as $\varepsilon \downarrow 0$.

The transition probabilities of each random walk $X^{(\varepsilon)}$ are

$$P_t^{(\varepsilon)}(x) := \mathbb{P}_0\{X_t^{(\varepsilon)} = x\} \quad (t \geq 0, x \in \varepsilon\mathbf{Z}). \quad (2.9)$$

Let $\mathcal{L}^{(\varepsilon)}$ denote the generator of $X^{(\varepsilon)}$, and choose and fix a bounded measurable function $\phi : \varepsilon\mathbf{Z} \rightarrow \mathbf{R}$. Because

$$(\mathcal{L}^{(\varepsilon)}\phi)(x) = \varepsilon^{-\alpha} (\mathcal{L}(\pi_\varepsilon\phi))(x/\varepsilon) \quad \text{for all } x \in \varepsilon\mathbf{Z}, \quad (2.10)$$

we may rewrite (SHE $_\varepsilon$) as the following system of Itô SDEs: For all $x \in \varepsilon\mathbf{Z}$,

$$dU_t^{(\varepsilon)}(x) = \varepsilon^\alpha \varrho_\varepsilon \cdot \left(\mathcal{L}^{(\varepsilon)} U_t^{(\varepsilon)} \right) (x) dt + \varepsilon^{-1/2} \sigma \left(U_t^{(\varepsilon)}(x) \right) dB_t(x/\varepsilon), \quad (2.11)$$

subject to $U_0^{(\varepsilon)} \equiv 1$. From now on, we adopt the normalization (1.11) without further mention. In that case, we have

$$dU_t^{(\varepsilon)}(x) = \left(\mathcal{L}^{(\varepsilon)} U_t^{(\varepsilon)} \right) (x) dt + \varepsilon^{-1/2} \sigma \left(U_t^{(\varepsilon)}(x) \right) dB_t(x/\varepsilon), \quad (2.12)$$

for $t > 0$ and $x \in \varepsilon\mathbf{Z}$, subject to $U_0^{(\varepsilon)} \equiv 1$. It follows from elementary Markov chain theory that $(s, t, x, y) \mapsto P_{t-s}^{(\varepsilon)}(y - x)$ is the fundamental solution to the Kolmogorov–Fokker–Planck equation $(\partial/\partial t)p - \mathcal{L}^{(\varepsilon)}p = 0$. Therefore, we may rewrite (2.12)—equivalently (SHE $_\varepsilon$)—in mild form as follows:

$$U_t^{(\varepsilon)}(x) = 1 + \varepsilon^{-1/2} \sum_{y \in \varepsilon\mathbf{Z}} \int_0^t P_{t-s}^{(\varepsilon)}(y - x) \sigma \left(U_s^{(\varepsilon)}(y) \right) dB_s(y/\varepsilon), \quad (2.13)$$

valid for all $t > 0$ and $x \in \varepsilon\mathbf{Z}$. According to Shiga and Shimizu [23], the system (2.13) of interacting SDEs has a unique strong solution $U^{(\varepsilon)}$ that is continuous in t almost surely. We appeal to this fact tacitly from now on.

2.4 The main result, and some consequences

We will first state the assumption needed for our main result. Let μ denote the *dislocation distribution* of the random walk X on \mathbf{Z} ; that is, for all $A \subseteq \mathbf{R}$,

$$\mu(A) := \mathbb{P}\{X_\gamma \in A\}, \quad \text{where } \gamma := \inf\{s > 0 : X_s \neq 0\}. \quad (2.14)$$

The Fourier transform $\hat{\mu}$ of μ is computed easily as

$$\hat{\mu}(z) := \sum_{x \in \mathbf{Z}} e^{izx} \mu(\{x\}) = \mathbb{E} \exp(izX_\gamma) \quad (-\pi \leq z \leq \pi). \quad (2.15)$$

We will refer to $\hat{\mu}$ as the *characteristic function* of the random walk X . It is possible to show that, if

$$1 - \hat{\mu}(z) = \nu|z|^\alpha + o(|z|^\alpha) \quad \text{as } |z| \rightarrow 0, \quad (2.16)$$

then the finite-dimensional distributions of the normalized random walk $X^{(\varepsilon)}$ —see (2.8)—converge to those of a symmetric α -stable Lévy process with transition function $p_t(x)$, as $\varepsilon \downarrow 0$. We will need to strengthen the preceding property for our results. The following assumption restricts a little further the behavior of the characteristic function of μ near 0.

Assumption 2.1. We assume that the dislocation measure μ satisfies the following: (i) $\{z \in [-\pi, \pi] : \hat{\mu}(z) = 1\} = \{0\}$; and (ii) There exists $a > 0$ such that

$$1 - \hat{\mu}(z) = \nu|z|^\alpha + O(|z|^{\alpha+a}) \quad \text{as } |z| \rightarrow 0. \quad (2.17)$$

Example 2.2 ($\alpha = 2$). Recall [24] that the random walk X is called *strongly aperiodic* when $|\hat{\mu}(z)| = 1$ if and only if z is an integer multiple of 2π . Thus, it follows that Assumption 2.1(i) holds for all strongly aperiodic random walks. There are also random walks that satisfy Assumption 2.1(i) but are not strongly aperiodic. The simple symmetric random walk [on \mathbf{Z}] is a good example. In fact, the simple walk satisfies all of Assumption 2.1, with $\nu = \frac{1}{2}$. More generally, we now show that if $Y_1 = X_\gamma$ satisfies $\mathbb{E}(Y_1^2) = 2\nu$ and $Y_1 \in L^{2+\eta}(\Omega)$ for some $\eta > 0$, then Assumption 2.1(ii) holds for the same value $\nu = \frac{1}{2}\mathbb{E}(Y_1^2)$. [It might help to recall that the stopping time γ was defined earlier in (2.14).] Without loss of generality, we may—and will—assume that $\eta \in (0, 1)$. Equivalently, we are assuming that

$$\mathbb{E}(|Y_1|^{3-\delta}) < \infty \quad \text{for some } \delta \in (0, 1). \quad (2.18)$$

Because $|1 - \cos x - (x^2/2)| \leq C \min(x^2, |x|^3)$ for all $x \in \mathbf{R}$, and since $\mathbb{E}(Y_1^2) = 2\nu$, we plug in $x := zY_1$ in the above inequality and integrate [dP] in order to see that

$$|1 - \hat{\mu}(z) - \nu z^2| \leq Cz^2 \mathbb{E}[\min(|z||Y_1|^3, Y_1^2)], \quad (2.19)$$

for all $z \in \mathbf{R}$. In particular, we can see that Assumption 2.1 holds as long as there exists $a > 0$ such that

$$\mathbb{E}[\min(|z||Y_1|^3, Y_1^2)] = O(|z|^a) \quad \text{as } |z| \rightarrow 0. \quad (2.20)$$

Next, let us write, for any value of $\delta \in (0, 1)$,

$$\begin{aligned} \mathbf{E} [\min(|z||Y_1|^3, Y_1^2)] &= |z| \mathbf{E} [|Y_1|^3; |Y_1| \leq |z|^{-1}] + \mathbf{E} [Y_1^2; |Y_1| > |z|^{-1}] \\ &\leq |z|^{1-\delta} \mathbf{E} (|Y_1|^{3-\delta}) + \mathbf{E} [Y_1^2; |Y_1| > |z|^{-1}]. \end{aligned} \quad (2.21)$$

We first apply Hölder's inequality and then Chebyshev's inequality in order to see that

$$\begin{aligned} \mathbf{E} [Y_1^2; |Y_1| > |z|^{-1}] &\leq [\mathbf{E} (|Y_1|^{3-\delta})]^{2/(3-\delta)} [\mathbf{P}\{|Y_1| > |z|^{-1}\}]^{(1-\delta)/(3-\delta)} \\ &\leq |z|^{1-\delta} \mathbf{E} (|Y_1|^{3-\delta}). \end{aligned} \quad (2.22)$$

Thus we deduce (2.20) from (2.18). \square

Example 2.3 ($\alpha < 2$). There are also random walks on \mathbf{Z} that satisfy Assumption 2.1 when $1 < \alpha < 2$. Let us choose and fix $\alpha \in (1, 2)$ and define

$$\mu(\{j\}) := \frac{1}{2\zeta(\alpha+1)|j|^{\alpha+1}} \quad \text{for } j = \pm 1, \pm 2, \dots, \quad (2.23)$$

where ζ denotes the zeta function. Evidently, μ defines a symmetric probability measure on \mathbf{Z} , and

$$1 - \hat{\mu}(z) = \frac{1}{\zeta(\alpha+1)} \sum_{j=1}^{\infty} \frac{1 - \cos(jz)}{j^{\alpha+1}} \quad \text{for every } z \geq 0. \quad (2.24)$$

Assumption 2.1(i) follows readily from this representation. Also, a few lines of direct computation show that there exists a finite constant C such that

$$\sum_{j=1}^{\infty} \left| \int_j^{j+1} \frac{1 - \cos(xz)}{x^{\alpha+1}} dx - \frac{1 - \cos(jz)}{j^{\alpha+1}} \right| \leq C \max(z^2, z^{1+\alpha}), \quad (2.25)$$

for all $z \geq 0$. This yields immediately that $1 - \hat{\mu}(z) = \nu z^\alpha + O(z^2)$ as $z \downarrow 0$, where

$$\nu := \frac{1}{\zeta(\alpha+1)} \int_0^\infty \frac{1 - \cos x}{x^{\alpha+1}} dx = -\frac{\pi}{2\zeta(\alpha+1)\Gamma(\alpha+1)\sin(\alpha\pi/2)}; \quad (2.26)$$

see [18, §4, p. 13] for the details of this last computation. Thus, Assumption 2.1(ii) also holds for the preceding value of ν [with $a := 2 - \alpha > 0$]. \square

The following is our main invariance principle.

Theorem 2.4. *Suppose \mathcal{L} is the generator of a symmetric random walk $\{X_t\}_{t \geq 0}$ on \mathbf{Z} whose characteristic function satisfies Assumption 2.1. Then there exists a coupling of $\{U^{(\varepsilon)}\}_{\varepsilon \in (0,1)}$ and u —all on a common underlying probability space—such that*

$$\sup_{t \in [0, T]} \sup_{x \in \varepsilon \mathbf{Z}} \mathbf{E} \left(\left| U_t^{(\varepsilon)}(x) - u_t(x) \right|^k \right) = o(\varepsilon^{\rho k/2}) \quad (\varepsilon \downarrow 0), \quad (2.27)$$

for every fixed $T > 0$, $k \geq 2$, and $\rho \in (0, a \wedge [\alpha - 1])$.

We will see also that Theorem 2.4 implies the following, which in turns implies Corollary 1.1 readily [i.e., without need for proof].

Theorem 2.5 (Strong approximations). *Under the assumptions of Theorem 2.4, the preceding coupling satisfies the following: For all $M, T > 0$ and $\rho \in (0, a \wedge \alpha - 1)$,*

$$\varepsilon^{-\rho/2} \sup_{t \in (0, T)} \sup_{\substack{x \in \varepsilon \mathbf{Z}: \\ |x| \leq \varepsilon^{-M}}} \left| U_t^{(\varepsilon)}(x) - u_t(x) \right| \xrightarrow{\mathbf{P}} 0 \quad \text{as } \varepsilon \downarrow 0. \quad (2.28)$$

Using a comparison of moments argument for interacting diffusions, one can also obtain the following consequence of Theorem 2.4; this was alluded to earlier.

Theorem 2.6 (Comparison of moments). *Let the assumptions of Theorem 2.4 hold. Suppose \bar{u} is the solution to (SHE) subject to $\bar{u}_0 \equiv 1$, where σ is replaced everywhere by $\bar{\sigma}$ for a function $\bar{\sigma} : \mathbf{R} \rightarrow \mathbf{R}$ that is Lipschitz continuous, and $0 \leq \sigma(z) \leq \bar{\sigma}(z)$ for all $z \in \mathbf{R}$. Let us assume further that $\sigma(0) = \bar{\sigma}(0) = 0$. Then,*

$$\mathbf{E} \left[\prod_{j=1}^m u_t(x_j) \right] \leq \mathbf{E} \left[\prod_{j=1}^m \bar{u}_t(x_j) \right], \quad (2.29)$$

for all $t \geq 0$ and $x_1, \dots, x_m \in \mathbf{R}$.

The following theorem gives upper and lower bounds on the exponential rate of growth of the moments (Lyapunov-exponents) of the stochastic heat equation when the nonlinearity is “roughly” linear.

Theorem 2.7 (Lyapunov-exponent bounds). *Suppose u solves (SHE) with $\mathcal{L}f = \nu f''$ subject to initial profile $u_0 \equiv 1$. If there exists nonnegative constants $0 \leq L_\sigma \leq \text{Lip}_\sigma < \infty$ such that $L_\sigma |z| \leq \sigma(z) \leq \text{Lip}_\sigma |z|$ for all z , then for all integers $k \geq 2$ and reals $x \in \mathbf{R}$,*

$$\begin{aligned} \frac{L_\sigma^4 k(k^2 - 1)}{48\nu} &\leq \liminf_{t \rightarrow \infty} \frac{1}{t} \log \mathbf{E} (|u_t(x)|^k) \\ &\leq \limsup_{t \rightarrow \infty} \frac{1}{t} \log \mathbf{E} (|u_t(x)|^k) \leq \frac{\text{Lip}_\sigma^4 k^3}{\nu}. \end{aligned} \quad (2.30)$$

Ivan Corwin and Roger Tribe have independently asked us whether one can transfer the knowledge of Lyapunov exponents for linear SPDEs to nonlinear ones [private communications]. The preceding answers their question in the affirmative.

Let us conclude this discussion by stating two open problems.

Open Problem. Consider the parabolic Anderson model; that is, (SHE) with $\sigma(x) = \lambda x$, where $\lambda \in \mathbf{R}$ is a fixed constant. According to a predication of Kardar [17], one would expect that

$$\lim_{t \rightarrow \infty} \frac{1}{t} \log \mathbf{E} (|u_t(x)|^k) = \frac{\lambda^4 k(k^2 - 1)}{48\nu} \quad \text{for every } x \in \mathbf{R}. \quad (2.31)$$

[The left-hand side is independent of x , actually; see the proof of Lemma 18 in Dalang [6].] Bertini and Cancrini [1] have invented a rigorous Feynman–Kac moment formula, which suggests the Kardar formula (2.31) though their claim to proof of (2.31) is false because of an incorrect appeal to the Skorohod lemma. As far as we know a rigorous proof of (2.31) is still not available. Our Theorem 2.6 shows that when $L_\sigma|z| \leq \sigma(z) \leq \text{Lip}_\sigma|z|$ for all z ,

$$\frac{(\text{Lip}_\sigma^4 + o(1))k(k^2 - 1)}{48\nu} \leq \frac{1}{t} \log \mathbb{E} (|u_t(x)|^k) \leq \frac{(\text{Lip}_\sigma^4 + o(1))k^3}{\nu}, \quad (2.32)$$

as $t \rightarrow \infty$. In the case that the initial profile is a point mass [instead of $u_0(x) \equiv 1$, as is the case here], Borodin and Corwin [3, §4.2] have succeeded in proving (2.31). Can their methods be used to establish (2.31) when $u_0(x) \equiv 1$? If so, then comparison methods of this paper will improve (2.32) to the following, which is the best possible bound of its type: As $t \rightarrow \infty$.

$$\frac{(\text{Lip}_\sigma^4 + o(1))k(k^2 - 1)}{48\nu} \leq \frac{1}{t} \log \mathbb{E} (|u_t(x)|^k) \leq \frac{(\text{Lip}_\sigma^4 + o(1))k(k^2 - 1)}{48\nu}. \quad (2.33)$$

Open Problem. For a more challenging problem consider (SHE) in the case that $\lim_{z \rightarrow \infty} \sigma(z)/z = \lambda > 0$; this is an “asymptotic Anderson model.” Is it true that the $\lim_{t \rightarrow \infty} t^{-1} \sup_{x \in \mathbf{R}} \log \mathbb{E} (|u_t(x)|^k)$ exists for some integer $k \geq 2$? Theorem 2.10 of Foondun and Khoshnevisan [9] suggests that this might be the case provided that the initial profile is a bounded and measurable non-random function that satisfies $\inf_{x \in \mathbf{R}} u_0(x) \geq \eta$ for a sufficiently large $\eta > 0$. However, a proof, or disproof, is completely missing.

3 Proof of Theorem 2.4

Let us say that $X^{(\varepsilon)} \approx Y^{(\varepsilon)}$ whenever $X^{(\varepsilon)} := \{X_t^{(\varepsilon)}(x)\}$ and $Y^{(\varepsilon)} := \{Y_t^{(\varepsilon)}(x)\}$ are space-time random fields that are indexed as $t > 0$ and $x \in A_\varepsilon \subset \mathbf{R}$, and satisfy the following:

$$\sup_{t \in [0, T]} \sup_{x \in A_\varepsilon} \left\| X_t^{(\varepsilon)}(x) - Y_t^{(\varepsilon)}(x) \right\|_k = o(\varepsilon^{\rho/2}) \quad (\varepsilon \downarrow 0), \quad (3.1)$$

for all $T > 0$, $k \geq 2$ and $0 < \rho < a \wedge (\alpha - 1)$.

Our surrogates from $X^{(\varepsilon)}$ and $Y^{(\varepsilon)}$ will be selected from a list of 5 random fields. The first 4 are:

$$u_t^{(\varepsilon)}(x) := 1 + \int_{(0, t - \varepsilon^\alpha) \times \mathbf{R}} p_{t-s}(y-x) \sigma(u_s(\varepsilon[y/\varepsilon])) W(ds dy); \quad (3.2)$$

$$v_t^{(\varepsilon)}(x) := 1 + \int_{(0, t - \varepsilon^\alpha) \times \mathbf{R}} p_{t-s}(\varepsilon[y/\varepsilon] - \varepsilon[x/\varepsilon]) \sigma(u_s(\varepsilon[y/\varepsilon])) W(ds dy);$$

$$V_t^{(\varepsilon)}(x) := 1 + \varepsilon^{-1} \int_{(0, t - \varepsilon^\alpha) \times \mathbf{R}} P_{t-s}^{(\varepsilon)}(\varepsilon[y/\varepsilon] - \varepsilon[x/\varepsilon]) \sigma(u_s(\varepsilon[y/\varepsilon])) W(ds dy);$$

$$W_t^{(\varepsilon)}(x) := 1 + \varepsilon^{-1} \int_{(0, t) \times \mathbf{R}} P_{t-s}^{(\varepsilon)}(\varepsilon[y/\varepsilon] - \varepsilon[x/\varepsilon]) \sigma(u_s(\varepsilon[y/\varepsilon])) W(ds dy),$$

where $[w] :=$ the greatest integer $\leq w$ when $w \geq 0$, and $[w] :=$ the smallest integer $\geq w$ when $w \leq 0$.

There is a fifth random field $\bar{U}^{(\varepsilon)} := \{\bar{U}_t^{(\varepsilon)}(x)\}_{t \geq 0, x \in \varepsilon \mathbf{Z}}$ of interest to us; that is the solution to an infinite system of Itô stochastic differential equations that is defined by

$$\bar{U}_t^{(\varepsilon)}(x) = 1 + \varepsilon^{-1/2} \sum_{j=-\infty}^{\infty} \int_0^t P_{t-s}^{(\varepsilon)}(j\varepsilon - x) \sigma\left(\bar{U}_s^{(\varepsilon)}(j\varepsilon)\right) dB_s^{(j,\varepsilon)}, \quad (3.3)$$

where

$$B_t^{(j,\varepsilon)} := \varepsilon^{-1/2} \int_{(0,t) \times (j\varepsilon, (j+1)\varepsilon)} W(ds dy). \quad (3.4)$$

Elementary properties of the Wiener integral imply that $\{B_t^{(j,\varepsilon)}\}_{t \geq 0, j \in \mathbf{Z}}$ is a [relabelled] version of the Brownian motions $\{B_t(x)\}_{t \geq 0, x \in \mathbf{Z}}$ of the Introduction. As such, the theory of Shiga and Shimizu [23] assures us that the infinite system (3.3) of Itô stochastic ODEs has a unique strong solution. Moreover, the strong existence and uniqueness of the solutions to (2.12) and (3.3) together imply that $\{\bar{U}_t^{(\varepsilon)}(x)\}_{t \geq 0, x \in \varepsilon \mathbf{Z}}$ is a modification of $\{U_t^{(\varepsilon)}(x)\}_{t \geq 0, x \in \varepsilon \mathbf{Z}}$. Thus, Theorem 2.4 is a quantitative way to say that $u \approx \bar{U}^{(\varepsilon)}$, and follows once we verify the following chain of successive approximations (with $A_\varepsilon = \varepsilon \mathbf{Z}$):

- (1). $u \approx u^{(\varepsilon)}$;
- (2). $u^{(\varepsilon)} \approx v^{(\varepsilon)}$;
- (3). $v^{(\varepsilon)} \approx V^{(\varepsilon)}$;
- (4). $V^{(\varepsilon)} \approx W^{(\varepsilon)}$; and
- (5). $W^{(\varepsilon)} \approx \bar{U}^{(\varepsilon)}$.

We now proceed to prove the theorem using the outlined five steps. Henceforth, all unimportant finite constants are denoted by C . These constants might vary from line to line, and might depend only on the parameters (k, T, ν, α) .

3.1 Verification of Assertion (1)

We bound the mean square difference between $u_t(x)$ and $u_t^{(\varepsilon)}(x)$, using (BDG), as follows:

$$\begin{aligned} & \left\| u_t(x) - u_t^{(\varepsilon)}(x) \right\|_k^2 \\ & \leq 2 \cdot (4k)^{1/k} \int_{t-\varepsilon^\alpha}^t ds \int_{-\infty}^{\infty} dy |p_{t-s}(y-x)|^2 \|\sigma(u_s(y))\|_k^2 \\ & \quad + 2 \cdot (4k)^{1/k} \int_0^{t-\varepsilon^\alpha} ds \int_{-\infty}^{\infty} dy |p_{t-s}(y-x)|^2 \|\sigma(u_s(y)) - \sigma(u_s(\varepsilon[y/\varepsilon]))\|_k^2 \\ & \leq C \int_0^{\varepsilon^\alpha} ds \int_{-\infty}^{\infty} dy |p_s(y)|^2 + C \int_0^t ds \int_{-\infty}^{\infty} dy |p_{t-s}(y)|^2 \sup_{|z-z'| \leq \varepsilon} \|u_s(z) - u_s(z')\|_k^2. \end{aligned} \quad (3.5)$$

In the final step we have used the facts that: (i) σ is a Lipschitz continuous function; and (ii) $\sup_{0 \leq s \leq T} \mathbb{E}(|u_s(y)|^k) < \infty$. By the semigroup property and the inversion theorem of Fourier transforms,

$$\int_{-\infty}^{\infty} |p_s(y)|^2 dy = (p_s * p_s)(0) = p_{2s}(0) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{-2\nu|z|^\alpha s} dz = \frac{C}{s^{1/\alpha}}. \quad (3.6)$$

Therefore,

$$\int_0^{\varepsilon^\alpha} ds \int_{-\infty}^{\infty} dy |p_s(y)|^2 \leq C\varepsilon^{\alpha-1}. \quad (3.7)$$

This estimates the first term on the right-most side of (3.5).

Now we bound the second term. Thanks to the Itô isometry (2.4) and the uniform bound of the k th moment of u , we conclude that whenever $|z - z'| \leq \varepsilon$,

$$\|u_s(z) - u_s(z')\|_k^2 \leq C \int_0^s dr \int_{-\infty}^{\infty} dw [p_{s-r}(w - z) - p_{s-r}(w - z')]^2. \quad (3.8)$$

Therefore, Plancherel's identity yields the bound

$$\begin{aligned} \|u_s(z) - u_s(z')\|_k^2 &\leq C \int_0^s dr \int_{-\infty}^{\infty} d\zeta e^{-2\nu(s-r)|\zeta|^\alpha} |e^{i(z-z')\zeta} - 1|^2 \\ &\leq C e^{2\nu s} \int_0^s dr \int_0^{\infty} d\zeta e^{-2\nu r(1+\zeta^\alpha)} (1 \wedge |z - z'| \zeta)^2 \\ &\leq C \int_0^{\infty} \frac{(1 \wedge |z - z'| \zeta)^2}{1 + \zeta^\alpha} d\zeta. \end{aligned} \quad (3.9)$$

We split the final integral according to whether or not $\zeta < |z - z'|^{-1}$ and compute directly to find that the preceding integral is at most $C|z - z'|^{\alpha-1}$. Therefore, (3.6) yields the inequality,

$$\int_0^t ds \int_{-\infty}^{\infty} dy |p_{t-s}(y)|^2 \sup_{0 \leq z, z' \leq \varepsilon} \|u_s(z) - u_s(z')\|_k^2 \leq C\varepsilon^{\alpha-1}. \quad (3.10)$$

Because $\alpha - 1 > \rho$, Assertion (1) follows from (3.5), (3.7), and (3.10). \square

3.2 Verification of Assertion (2)

For every integer $k \geq 2$, the k th moment of $\sigma(u_s(y))$ is finite uniformly for all $y \in \mathbf{R}$ and $s \in [0, T]$. Therefore, Itô's isometry yields

$$\left\| u_t^{(\varepsilon)}(x) - v_t^{(\varepsilon)}(x) \right\|_k^2 \leq C \int_{\varepsilon^\alpha}^t ds \int_{-\infty}^{\infty} dy [p_s(y) - p_s(\varepsilon[y/\varepsilon])]^2 \quad (3.11)$$

It is well known that the stable probability density p_1 is C^∞ and satisfies

$$|p_1'(x)| \leq \frac{C}{1 + |x|^{\alpha+1}} \quad \text{for all } x \in \mathbf{R}. \quad (3.12)$$

When $1 < \alpha < 2$, this inequality is classical (see Głowacki [13, Corollary (5.30)] for some of the more recent developments for stable densities over homogeneous spaces). When $\alpha = 2$, one has $|p'_1(x)| \leq C(1 + |x|^3)^{-1}$ by explicit computation. Consequently, scaling reveals that for all $\alpha \in (1, 2]$, $s > 0$, and $x \in \mathbf{R}$,

$$|p'_s(x)| = \left| s^{-2/\alpha} p'_1\left(\frac{x}{s^{1/\alpha}}\right) \right| \leq \frac{C s^{(\alpha-1)/\alpha}}{s^{(\alpha+1)/\alpha} + |x|^{\alpha+1}}. \quad (3.13)$$

With the above bound in hand, we obtain

$$\begin{aligned} & \left\| u_t^{(\varepsilon)}(x) - v_t^{(\varepsilon)}(x) \right\|_k^2 \\ & \leq C \int_{\varepsilon^\alpha}^t ds \int_0^\infty dy \left| \int_{(y-\varepsilon)_+}^y \frac{s^{(\alpha-1)/\alpha}}{s^{(\alpha+1)/\alpha} + x^{\alpha+1}} dx \right|^2 \\ & \leq C \varepsilon^2 \int_{\varepsilon^\alpha}^t ds \int_0^\infty dy \left| \frac{s^{(\alpha-1)/\alpha}}{s^{(\alpha+1)/\alpha} + (y-\varepsilon)_+^{\alpha+1}} \right|^2 \\ & = C \varepsilon^3 \int_{\varepsilon^\alpha}^t \left| \frac{s^{(\alpha-1)/\alpha}}{s^{(\alpha+1)/\alpha}} \right|^2 ds + C \varepsilon^2 \int_{\varepsilon^\alpha}^t ds \int_0^\infty dw \left| \frac{s^{(\alpha-1)/\alpha}}{s^{(\alpha+1)/\alpha} + w^{\alpha+1}} \right|^2. \end{aligned} \quad (3.14)$$

By scaling, the latter dw -integral is equal to $C s^{-3/\alpha}$, whence

$$\begin{aligned} \left\| u_t^{(\varepsilon)}(x) - v_t^{(\varepsilon)}(x) \right\|_k^2 & \leq C \varepsilon^3 \int_{\varepsilon^\alpha}^\infty \frac{ds}{s^{4/\alpha}} + C \varepsilon^2 \int_{\varepsilon^\alpha}^\infty \frac{ds}{s^{3/\alpha}} \\ & \leq C \varepsilon^{\alpha-1}. \end{aligned} \quad (3.15)$$

Because $\alpha - 1 > \rho$, this completes our demonstration of Assertion (2). \square

3.3 Verification of Assertion (3)

The proof of Assertion (3) requires a certain form of a local limit theorem for continuous-time random walks that are in the domain of attraction of a symmetric α -stable Lévy processes. As a first step in this direction we develop a suitable quantitative form of a local central limit theorem for continuous-time random walks that lie in the domain of attraction of a symmetric α -stable Lévy process and satisfy the slightly stronger Assumption 2.1.

Proposition 3.1 (A local CLT). *For all $T > 0$ there exist positive and finite constants K, C, ε_0 such that*

$$\begin{aligned} & \sup_{x \in \varepsilon \mathbf{Z}} \left| \frac{P_t^{(\varepsilon)}(x)}{\varepsilon} - p_t(x) \right| \\ & \leq C \times \begin{cases} \frac{\varepsilon^\alpha |\ln \varepsilon|^{(a+\alpha)/\alpha}}{t^{(a+1)/\alpha}} & \text{if } t \geq K \varepsilon^\alpha |\ln \varepsilon|^{(a+\alpha)/\alpha}, \\ t^{-1/\alpha} + \varepsilon^{-1} & \text{otherwise,} \end{cases} \end{aligned} \quad (3.16)$$

uniformly for all $t \in (0, T]$ and $\varepsilon \in (0, \varepsilon_0)$.

When $\alpha = 2$, the preceding is a sharp form of the usual local central limit theorem for random walks. The standard method to obtain such a result in continuous time is to appeal to the classical local limit theorem for discrete-time random walks, and then “Poissonize.” See Theorem 2.5.6 of Lawler and Limic [20]. Proposition 3.1 is a much stronger version of the resulting estimate. As was mentioned earlier, Proposition 3.1 tells us that $\varepsilon^{-1}P_t^{(\varepsilon)}(x) \approx p_t(x)$ when ε is small. The more significant part of our estimate is that it states that this approximation is good uniformly for all times $t \geq \varepsilon^{\alpha+o(1)}$; it is not hard to see from the forthcoming proof that this time bound is basically sharp. The proof is similar to—though not identical to—the standard proof of the classical local limit theorem for discrete-time random walks, but we work directly in continuous time, rather than “Poissonize.”

Proof. By the Fourier inversion theorem,

$$p_t(x) = \frac{1}{\pi} \int_0^\infty \cos(zx) e^{-\nu t z^\alpha} dz \leq p_t(0) = \frac{C}{t^{1/\alpha}}. \quad (3.17)$$

Since $P_t^{(\varepsilon)}(x) = \mathbb{P}\{X_{t/\varepsilon^\alpha} = \varepsilon^{-1}x\} \leq 1$, it follows readily from the nearly-tautological inequality $|\varepsilon^{-1}P_t^{(\varepsilon)}(x) - p_t(x)| \leq \varepsilon^{-1}P_t^{(\varepsilon)}(x) + p_t(x)$ that

$$\sup_{x \in \varepsilon \mathbf{Z}} \left| \frac{P_t^{(\varepsilon)}(x)}{\varepsilon} - p_t(x) \right| \leq C \left(\varepsilon^{-1} + t^{-1/\alpha} \right). \quad (3.18)$$

The preceding is valid for all t, ε , etc., but is useful to us in the case that $t < K\varepsilon^\alpha |\ln \varepsilon|^{(a+\alpha)/a}$. From now on, we will concentrate on the more subtle case that $t \geq K\varepsilon^\alpha |\ln \varepsilon|^{(a+\alpha)/a}$, where the value of K will be determined during the course of the proof.

Recall that we can write $X_t = \sum_{j=0}^{\Pi(t)} Y_j$, where Y_1, Y_2, \dots are i.i.d. random variables, each distributed as μ , and Π is an independent rate-one Poisson process. This representation, and elementary facts about Poisson random variables, together yield the following special case of the Lévy–Khintchine formula:

$$\mathbb{E} \exp(izX_{t/\varepsilon^\alpha}) = \exp(-t\varepsilon^{-\alpha}[1 - \hat{\mu}(z)]); \quad (3.19)$$

valid for all $\varepsilon > 0$, $t \geq 0$, and $z \in \mathbf{R}$. In accord with the inversion formula,

$$\frac{2\pi}{\varepsilon} P_t^{(\varepsilon)}(x) = \int_{-\pi/\varepsilon}^{\pi/\varepsilon} \exp\left(-ixz - \frac{t[1 - \hat{\mu}(\varepsilon z)]}{\varepsilon^\alpha}\right) dz. \quad (3.20)$$

Let us now choose and fix $\varepsilon \in (0, \frac{1}{2})$. Assumption 2.1 and the uniform boundedness of $\hat{\mu}$ together guarantee that

$$|1 - \hat{\mu}(w) - \nu|w|^\alpha| \leq C|w|^{a+\alpha} \quad \text{for all } w \in \mathbf{R}. \quad (3.21)$$

This bound has the following ready consequence: There exists $r_0 \in (0, \pi)$ such that

$$1 - \hat{\mu}(w) \geq \frac{\nu|w|^\alpha}{2} \quad \text{whenever } |w| \leq r_0. \quad (3.22)$$

[It might help to recall that $\hat{\mu}$ is real-valued by symmetry.]

Let us define

$$\lambda := \left(\frac{(10+2a)|\ln \varepsilon|}{\nu t} \right)^{1/\alpha}, \quad (3.23)$$

for notational simplicity. If $t > K\varepsilon^\alpha |\ln \varepsilon|^{(a+\alpha)/a}$, and $K > 20 + 4a$, then

$$\lambda \leq \left(\frac{|\ln \varepsilon|}{2\varepsilon^\alpha |\ln \varepsilon|^{(a+\alpha)/a}} \right)^{1/\alpha} < \frac{r_0}{\varepsilon}, \quad (3.24)$$

for all $\varepsilon < \varepsilon_0$, as long as $\varepsilon_0 > 0$ is sufficiently small. In this case, we may split up the Fourier integral of (3.20) into three regions as follows:

$$\frac{2\pi}{\varepsilon} P_t^{(\varepsilon)}(x) := \mathbb{I}_1 + \mathbb{I}_2 + \mathbb{I}_3; \quad (3.25)$$

where

$$\mathbb{I}_1 := \int_{|z| \leq \lambda} ; \quad \mathbb{I}_2 := \int_{\lambda < |z| \leq r_0/\varepsilon} ; \quad \mathbb{I}_3 := \int_{r_0/\varepsilon < |z| \leq \pi/\varepsilon}. \quad (3.26)$$

According to the inversion theorem,

$$2\pi p_t(x) = \int_{-\infty}^{\infty} \exp(-ixz - \nu t|z|^\alpha) dz. \quad (3.27)$$

Recall that our goal is to show that $\varepsilon^{-1} P_t^{(\varepsilon)}(x) \simeq p_t(x)$; with this aim in mind, we use a well-known strategy. Namely, we will demonstrate that:

- (i) $\mathbb{I}_1 \simeq 2\pi p_t(x)$; whereas
- (ii) $\mathbb{I}_2 \simeq \mathbb{I}_3 \simeq 0$.

Let us begin by estimating \mathbb{I}_1 , whose analysis turns out to require the most care. Clearly,

$$\begin{aligned} & \sup_{x \in \varepsilon \mathbf{Z}} |\mathbb{I}_1 - 2\pi p_t(x)| \\ & \leq \int_{-\lambda}^{\lambda} \left| \exp\left(-\frac{t[1 - \hat{\mu}(\varepsilon z)]}{\varepsilon^\alpha}\right) - e^{-\nu t|z|^\alpha} \right| dz + 2 \int_{\lambda}^{\infty} e^{-\nu t z^\alpha} dz \\ & = 2 \int_0^{\lambda} e^{-\nu t z^\alpha} \left| 1 - \exp\left(-\frac{t[1 - \hat{\mu}(\varepsilon z) - \nu(\varepsilon z)^\alpha]}{\varepsilon^\alpha}\right) \right| dz + \frac{2}{t^{1/\alpha}} \int_{\lambda t^{1/\alpha}}^{\infty} e^{-\nu y^\alpha} dy. \end{aligned} \quad (3.28)$$

Since $0 < \lambda < r_0 \varepsilon^{-1} < \pi \varepsilon^{-1}$, Assumption 2.1 implies that uniformly for all $z \in (0, \lambda)$ and $t \in (0, T]$,

$$\begin{aligned} t \left| \frac{1 - \hat{\mu}(\varepsilon z) - \nu(\varepsilon z)^\alpha}{\varepsilon^\alpha} \right| & \leq C t z^{a+\alpha} \varepsilon^a \\ & < C t \lambda^{a+\alpha} \varepsilon^{-a} \\ & = \frac{C \cdot (10+2a)^{(a+\alpha)/\alpha}}{t^{a/\alpha}} \varepsilon^a |\ln \varepsilon|^{(a+\alpha)/\alpha}. \end{aligned} \quad (3.29)$$

Since $t > K\varepsilon^\alpha |\ln \varepsilon|^{(a+\alpha)/a}$ for some $K > 20 + 4a$, the preceding implies that uniformly for all $z \in (0, \lambda)$ and $t \in (0, T]$,

$$t \left| \frac{1 - \hat{\mu}(\varepsilon z) - \nu(\varepsilon z)^\alpha}{\varepsilon^\alpha} \right| \leq \frac{C \cdot (10 + 2a)^{(a+\alpha)/\alpha}}{K^{a/\alpha}} < 1, \quad (3.30)$$

provided that we select K sufficiently large. Henceforth, we choose and fix $K > 20 + 4a$ large enough to ensure the preceding condition. Since $|1 - \exp(-w)| \leq C|w|$ for all $w \in \mathbf{C}$ with $|w| < 1$, it follows from Assumption 2.1 that whenever $t > K\varepsilon^\alpha |\ln \varepsilon|^{(a+\alpha)/a}$ for our choice of K ,

$$\begin{aligned} & \sup_{x \in \varepsilon \mathbf{Z}} |\mathbb{I}_1 - 2\pi p_t(x)| \\ & \leq \frac{C \cdot (10 + 2a)^{(a+\alpha)/\alpha}}{t^{a/\alpha}} \varepsilon^a |\ln \varepsilon|^{(a+\alpha)/\alpha} \int_0^\lambda e^{-\nu t z^\alpha} dz + \frac{2}{t^{1/\alpha}} \int_{\lambda t^{1/\alpha}}^\infty e^{-\nu y^\alpha} dy \\ & \leq \frac{C \cdot (10 + 2a)^{(a+\alpha)/\alpha}}{t^{(a+1)/\alpha}} \varepsilon^a |\ln \varepsilon|^{(a+\alpha)/\alpha} + \frac{2}{t^{1/\alpha}} \int_{\lambda t^{1/\alpha}}^\infty e^{-\nu y^\alpha} dy. \end{aligned} \quad (3.31)$$

Because $\alpha > 1$, l'Hôpital's rule implies that $\int_q^\infty \exp(-\nu y^\alpha) dy \leq C \exp(-\nu q^\alpha/2)$ for all $q > 0$. Hence, for every $t \in (K\varepsilon^\alpha |\ln \varepsilon|^{(a+\alpha)/a}, T]$,

$$\begin{aligned} \sup_{x \in \varepsilon \mathbf{Z}} |\mathbb{I}_1 - 2\pi p_t(x)| & \leq \frac{C\varepsilon^a |\ln \varepsilon|^{(a+\alpha)/\alpha}}{t^{(a+1)/\alpha}} + \frac{C e^{-\nu t \lambda^\alpha/2}}{t^{1/\alpha}} \\ & = \frac{C\varepsilon^a |\ln \varepsilon|^{(a+\alpha)/\alpha}}{t^{(a+1)/\alpha}} + \frac{C\varepsilon^{5+a}}{t^{1/\alpha}} \\ & \leq \frac{C\varepsilon^a |\ln \varepsilon|^{(a+\alpha)/\alpha}}{t^{(a+1)/\alpha}}, \end{aligned} \quad (3.32)$$

thanks to the definition of λ .

Next we bound \mathbb{I}_2 , using (3.22), as follows:

$$\begin{aligned} \sup_{x \in \varepsilon \mathbf{Z}} |\mathbb{I}_2| & \leq \int_{\lambda < |z| \leq r_0/\varepsilon} \exp\left(\frac{-t[1 - \hat{\mu}(\varepsilon z)]}{\varepsilon^\alpha}\right) dz \\ & \leq C \int_\lambda^\infty e^{-t\nu z^\alpha/2} dz \\ & \leq \frac{C e^{-\nu t \lambda^\alpha/2}}{t^{1/\alpha}} \\ & = \frac{C\varepsilon^{5+a}}{t^{1/\alpha}}. \end{aligned} \quad (3.33)$$

Finally we bound the quantity \mathbb{I}_3 as follows: Since $\hat{\mu}$ is continuous and real valued, thanks to symmetry, part (i) of Assumption 2.1 ensures that $\theta :=$

$\sup_{r_0 \leq |z| \leq \pi} \hat{\mu}(z) < 1$, and hence

$$\begin{aligned}
\sup_{x \in \varepsilon \mathbf{Z}} |\mathbb{I}_3| &\leq \frac{C}{\varepsilon} \exp \left[-\frac{t(1-\theta)}{\varepsilon^\alpha} \right] \\
&\leq \frac{C}{\varepsilon} \exp \left(-\frac{t}{C\varepsilon^\alpha} \right) \\
&\leq \frac{C}{t^{1/\alpha}} \exp \left(-\frac{t}{C\varepsilon^\alpha} \right) \\
&\leq \frac{C\varepsilon^5}{t^{1/\alpha}},
\end{aligned} \tag{3.34}$$

uniformly for $t > K\varepsilon^\alpha |\ln \varepsilon|^{(a+\alpha)/a}$. Therefore, (3.32), (3.33), and (3.34) together imply the proposition. \square

Proposition 3.1 has the following consequence for us.

Corollary 3.2. *Choose and fix $T > 0$. Then,*

$$\int_{\varepsilon^\alpha}^T dt \int_{-\infty}^{\infty} dy \left| \frac{P_t^{(\varepsilon)}(\varepsilon[y/\varepsilon])}{\varepsilon} - p_t(\varepsilon[y/\varepsilon]) \right|^2 \leq \varepsilon^{a\wedge(\alpha-1)+o(1)} \quad \text{as } \varepsilon \downarrow 0. \tag{3.35}$$

Proof. First of all, let us note that for all $\varepsilon \in (0, \frac{1}{2})$ and $t \in (0, T]$,

$$\int_{-\infty}^{\infty} \frac{P_t^{(\varepsilon)}(\varepsilon[y/\varepsilon])}{\varepsilon} dy = \sum_{j=-\infty}^{\infty} P_t^{(\varepsilon)}(j\varepsilon) = 1. \tag{3.36}$$

The preceding has an analogue for the stable density p_t as well. In order to prove that, recall that we can realize our symmetric stable process as $\{B_{T(t)}\}_{t \geq 0}$, where B denotes Brownian motion, and T an independent stable- 2α subordinator; see Bochner[2, §4.4]. In particular, it follows that we can write the stable heat kernel $p_t(x)$, via subordination, as follows:

$$p_t(x) = \int_0^\infty q_s(x) \tau_t(s) ds, \quad \text{where } q_s(x) := (2\pi s)^{-1/2} \exp\left(-\frac{x^2}{2s}\right) \tag{3.37}$$

denotes the Brownian motion transition probability, and τ_t is a probability density on $(0, \infty)$ for every $t > 0$. Since $x \mapsto q_s(x)$ is nonincreasing on $(0, \infty)$, it follows that p_t is, as well. We now use this conclusion as follows:

$$\begin{aligned}
\int_{-\infty}^{\infty} p_t(\varepsilon[y/\varepsilon]) dy &= 2\varepsilon \sum_{j=0}^{\infty} p_t(j\varepsilon) \\
&= 2\varepsilon p_t(0) + 2 \sum_{j=1}^{\infty} \int_{(j-1)\varepsilon}^{j\varepsilon} p_t(j\varepsilon) dy \\
&\leq 2\varepsilon p_t(0) + 2 \int_0^\infty p_t(y) dy \\
&= \frac{C\varepsilon}{t^{1/\alpha}} + 1.
\end{aligned} \tag{3.38}$$

We may combine (3.36) and (3.38) as follows:

$$\begin{aligned}
& \int_{-\infty}^{\infty} \left| \frac{P_t^{(\varepsilon)}(\varepsilon[y/\varepsilon])}{\varepsilon} - p_t(\varepsilon[y/\varepsilon]) \right|^2 dy \\
& \leq \mathcal{M}_t(\varepsilon) \cdot \int_{-\infty}^{\infty} \left[\frac{P_t^{(\varepsilon)}(\varepsilon[y/\varepsilon])}{\varepsilon} + p_t(\varepsilon[y/\varepsilon]) \right] dy \\
& \leq \mathcal{M}_t(\varepsilon) \cdot \left(\frac{C\varepsilon}{t^{1/\alpha}} + 2 \right),
\end{aligned} \tag{3.39}$$

where

$$\mathcal{M}_t(\varepsilon) := \sup_{y \in \varepsilon \mathbf{Z}} \left| \frac{P^{(\varepsilon)}(y)}{\varepsilon} - p_t(y) \right|. \tag{3.40}$$

In accord with Proposition 3.1,

$$\begin{aligned}
& \int_{\varepsilon^\alpha}^{K\varepsilon^\alpha |\ln \varepsilon|^{(a+\alpha)/a}} dt \int_{-\infty}^{\infty} dy \left| \frac{P_t^{(\varepsilon)}(\varepsilon[y/\varepsilon])}{\varepsilon} - p_t(\varepsilon[y/\varepsilon]) \right|^2 \\
& \leq C \int_{\varepsilon^\alpha}^{K\varepsilon^\alpha |\ln \varepsilon|^{(a+\alpha)/a}} \left(\frac{1}{t^{1/\alpha}} + \frac{1}{\varepsilon} \right) \left(\frac{\varepsilon}{t^{1/\alpha}} + 2 \right) dt \\
& \leq C\varepsilon \int_{\varepsilon^\alpha}^{K\varepsilon^\alpha |\ln \varepsilon|^{(a+\alpha)/a}} \frac{dt}{t^{2/\alpha}} + C \int_{\varepsilon^\alpha}^{K\varepsilon^\alpha |\ln \varepsilon|^{(a+\alpha)/a}} \frac{dt}{t^{1/\alpha}} + C\varepsilon^{\alpha-1} |\ln \varepsilon|^{(a+\alpha)/a} \\
& \leq C\varepsilon^{\alpha-1} |\ln \varepsilon|^{(a+\alpha)/a} \\
& = \varepsilon^{\alpha-1+o(1)}.
\end{aligned}$$

Similarly,

$$\begin{aligned}
& \int_{K\varepsilon^\alpha |\ln \varepsilon|^{(a+\alpha)/a}}^T dt \int_{-\infty}^{\infty} dy \left| \frac{P_t^{(\varepsilon)}(\varepsilon[y/\varepsilon])}{\varepsilon} - p_t(\varepsilon[y/\varepsilon]) \right|^2 \\
& \leq C \int_{K\varepsilon^\alpha |\ln \varepsilon|^{(a+\alpha)/a}}^T \frac{\varepsilon^a |\ln \varepsilon|^{(a+\alpha)/\alpha}}{t^{(a+1)/\alpha}} \left(\frac{\varepsilon}{t^{1/\alpha}} + 2 \right) dt \\
& \leq C\varepsilon^{a+1} |\ln \varepsilon|^{(a+\alpha)/\alpha} \int_{K\varepsilon^\alpha |\ln \varepsilon|^{(a+\alpha)/a}}^T \frac{dt}{t^{(a+2)/\alpha}} \\
& \quad + C\varepsilon^a |\ln \varepsilon|^{(a+\alpha)/\alpha} \int_{K\varepsilon^\alpha |\ln \varepsilon|^{(a+\alpha)/a}}^T \frac{dt}{t^{(a+1)/\alpha}} \\
& \leq \varepsilon^{a \wedge (\alpha-1) + o(1)},
\end{aligned} \tag{3.41}$$

as $\varepsilon \downarrow 0$, because $\alpha \in (1, 2]$. The corollary follows from these computations. \square

We are now ready to close this subsection by verifying Assertion (3). The proof of Corollary 3.2 can be readily adjusted in order to establish the following

variation of Corollary 3.2: Uniformly for all $x \in \varepsilon\mathbf{Z}$ and $t \in (\varepsilon^\alpha, T]$,

$$\int_{\varepsilon^\alpha}^t ds \int_{-\infty}^{\infty} dy \left| \frac{P_s^{(\varepsilon)}(\varepsilon[y/\varepsilon] - \varepsilon[x/\varepsilon])}{\varepsilon} - p_s(\varepsilon[y/\varepsilon] - \varepsilon[x/\varepsilon]) \right|^2 \quad (3.42)$$

$$\leq \varepsilon^{a \wedge (\alpha-1) + o(1)},$$

as $\varepsilon \downarrow 0$. We omit the minor remaining details. Since $\mathbb{E}(|\sigma(u_t(z))|^k)$ is bounded uniformly in $z \in \mathbf{R}$ and $t \in (0, T]$ by a finite constant $K_{T,k}$, an application of the Burkholder–Davis–Gundy inequality (BDG) yields

$$\begin{aligned} & \left\| v_t^{(\varepsilon)}(x) - V_t^{(\varepsilon)}(x) \right\|_k^2 \\ & \leq CK_{T,k} \int_{\varepsilon^\alpha}^t ds \int_{-\infty}^{\infty} dy \left| \frac{P_s^{(\varepsilon)}(\varepsilon[y/\varepsilon] - \varepsilon[x/\varepsilon])}{\varepsilon} - p_s(\varepsilon[y/\varepsilon] - \varepsilon[x/\varepsilon]) \right|^2 \\ & \leq \varepsilon^{a \wedge (\alpha-1) + o(1)}. \end{aligned} \quad (3.43)$$

Because $a \wedge (\alpha - 1) > \rho$, this verifies Assertion (3). \square

3.4 Verification of Assertion (4)

Since $\mathbb{E}(|\sigma(u_s(y))|^k)$ is bounded uniformly in $y \in \mathbf{R}$ and $s \in (0, T]$ for every integer $k \geq 2$, the Burkholder–Davis–Gundy inequality (BDG) implies that, uniformly for all $\varepsilon \in (0, \frac{1}{2})$, $x \in \varepsilon\mathbf{Z}$, and $t \in (0, T]$,

$$\begin{aligned} \left\| V_t^{(\varepsilon)}(x) - W_t^{(\varepsilon)}(x) \right\|_k^2 & \leq \frac{C}{\varepsilon^2} \int_0^{\varepsilon^\alpha} ds \int_{-\infty}^{\infty} dy \left| P_s^{(\varepsilon)}(\varepsilon[y/\varepsilon]) \right|^2 \\ & = \frac{C}{\varepsilon} \int_0^{\varepsilon^\alpha} ds \sum_{j=-\infty}^{\infty} \left| P_s^{(\varepsilon)}(j\varepsilon) \right|^2. \end{aligned} \quad (3.44)$$

Thanks to the Cauchy–Schwarz inequality,

$$\sum_{j=-\infty}^{\infty} \left| P_s^{(\varepsilon)}(j\varepsilon) \right|^2 \leq \left| \sum_{j=-\infty}^{\infty} P_s^{(\varepsilon)}(j\varepsilon) \right|^2 = 1. \quad (3.45)$$

Because $\alpha - 1 > \rho$, this proves Assertion (4). \square

3.5 Verification of Assertion (5)

We begin with a technical lemma.

Lemma 3.3. *The following holds*

$$\sup_{0 < \varepsilon < 1} \sup_{0 \leq t \leq T} \int_0^t ds \int_{-\infty}^{\infty} dy \left[\frac{P_{t-s}^{(\varepsilon)}(\varepsilon[y/\varepsilon] - x)}{\varepsilon} \right]^2 < \infty. \quad (3.46)$$

Proof. We first simplify the expression

$$\begin{aligned}
\int_0^t ds \int_{-\infty}^{\infty} dy \left[\frac{P_{t-s}^{(\varepsilon)}(\varepsilon[y/\varepsilon] - x)}{\varepsilon} \right]^2 &= \frac{1}{\varepsilon} \int_0^t ds \sum_{j=-\infty}^{\infty} [P_s^{(\varepsilon)}(j\varepsilon)]^2 \\
&= \varepsilon^{\alpha-1} \int_0^{t/\varepsilon^\alpha} dr \sum_{j=-\infty}^{\infty} |\mathbb{P}\{X_r = j\}|^2 \\
&= \varepsilon^{\alpha-1} \int_0^{t/\varepsilon^\alpha} dr \mathbb{P}\{Z_r = 0\}, \tag{3.47}
\end{aligned}$$

where $Z_r := X_r - \tilde{X}_r$ for an independent copy \tilde{X} of X . Of course, Z is a random walk with the same jump distribution as X but twice the jump rate. Thus,

$$\begin{aligned}
\int_0^t ds \int_{-\infty}^{\infty} dy \left[\frac{P_{t-s}^{(\varepsilon)}(\varepsilon[y/\varepsilon] - x)}{\varepsilon} \right]^2 &= \frac{\varepsilon^{\alpha-1}}{2} \int_0^{2t/\varepsilon^\alpha} \mathbb{P}\{X_r = 0\} dr \\
&= \frac{\varepsilon^{\alpha-1}}{4\pi} \int_{-\pi}^{\pi} \left\{ 1 - \exp\left(\frac{-2t[1 - \hat{\mu}(\xi)]}{\varepsilon^\alpha}\right) \right\} \frac{d\xi}{1 - \hat{\mu}(\xi)}, \tag{3.48}
\end{aligned}$$

where we used the Fourier inversion formula for the last line. Due to condition (i) of Assumption 2.1, we can find constants $0 < C_1 < C_2$ such that $C_1|\xi|^\alpha < 1 - \hat{\mu}(\xi) < C_2|\xi|^\alpha$ for $|\xi| \leq \pi$. With this observation in mind, we may split the final integral of (3.48) into two integrals: One is computed over the region $\{|\xi| \leq \varepsilon\}$, the other over $\{|\xi| > \varepsilon\}$. For the first integral, we use the bound $1 - e^{-x} \leq x$ valid for $x \geq 0$ and for the second integral, we use the bound $1 - e^{-x} \leq 1$. In this way we find that

$$\begin{aligned}
\int_0^t ds \int_{-\infty}^{\infty} dy \left[\frac{P_{t-s}^{(\varepsilon)}(\varepsilon[y/\varepsilon] - x)}{\varepsilon} \right]^2 &= \frac{\varepsilon^{\alpha-1}}{4\pi} \int_{|\xi| \leq \varepsilon} \frac{2t}{\varepsilon^\alpha} d\xi + \frac{\varepsilon^{\alpha-1}}{4\pi} \int_{\varepsilon < |\xi| \leq \pi} \frac{1}{C_1|\xi|^\alpha} d\xi. \tag{3.49}
\end{aligned}$$

This is a uniformly bounded quantity. \square

We begin our derivation of Assertion (5) with the observation that for $x \in \varepsilon\mathbf{Z}$,

$$\begin{aligned}
\bar{U}_t^{(\varepsilon)}(x) - W_t^{(\varepsilon)}(x) &= \frac{1}{\varepsilon} \int_{(0,t) \times \mathbf{R}} P_{t-s}^{(\varepsilon)}(\varepsilon[y/\varepsilon] - x) [\sigma(\bar{U}_s(\varepsilon[y/\varepsilon])) - \sigma(u_s(\varepsilon[y/\varepsilon]))] W(ds dy). \tag{3.50}
\end{aligned}$$

Consequently, we may apply the Burkholder–Davis–Gundy inequality (BDG), together with the triangle inequality, in order to find that the nonrandom function

$$\mathcal{D}(t) := \sup_{x \in \varepsilon\mathbf{Z}} \left\| \bar{U}_t^{(\varepsilon)}(x) - W_t^{(\varepsilon)}(x) \right\|_k^2 \quad (t \geq 0) \tag{3.51}$$

satisfies the recursive inequality

$$\mathcal{D}(t) \leq \frac{C}{\varepsilon^2} \int_0^t (\mathcal{D}(s) + \mathcal{E}(s)) ds \int_{-\infty}^{\infty} dy \left[P_{t-s}^{(\varepsilon)}(\varepsilon[y/\varepsilon] - x) \right]^2, \quad (3.52)$$

where

$$\mathcal{E}(s) := \sup_{y \in \varepsilon \mathbf{Z}} \left\| u_s(\varepsilon[y/\varepsilon]) - W_s^{(\varepsilon)}(\varepsilon[y/\varepsilon]) \right\|_k^2 \quad (s \geq 0). \quad (3.53)$$

The already-proven Assertions (1)–(4) together show that $\sup_{s \in (0, T]} \mathcal{E}(s) \leq C\varepsilon^\rho$ for all sufficiently small ε and $0 < \rho < a \wedge [\alpha - 1]$. Consequently, we arrive at the following recursive inequality, valid uniformly for every $t \in (0, T]$ and $\varepsilon > 0$ sufficiently small:

$$\begin{aligned} \mathcal{D}(t) &\leq C\varepsilon^\rho \int_0^t ds \int_{-\infty}^{\infty} dy \left| \frac{P_{t-s}^{(\varepsilon)}(\varepsilon[y/\varepsilon] - x)}{\varepsilon} \right|^2 \\ &\quad + C \int_0^t ds \int_{-\infty}^{\infty} dy \left| \frac{P_{t-s}^{(\varepsilon)}(\varepsilon[y/\varepsilon] - x)}{\varepsilon} \right|^2 \mathcal{D}(s) \\ &\leq C\varepsilon^\rho + C \int_0^t ds \mathcal{D}(s) \int_{-\infty}^{\infty} dy \left| \frac{P_{t-s}^{(\varepsilon)}(\varepsilon[y/\varepsilon] - x)}{\varepsilon} \right|^2. \end{aligned} \quad (3.54)$$

Gronwall's inequality reveals that $\mathcal{D}(t) \leq C\varepsilon^\rho$. In order to use that inequality, we need to know that $\sup_{0 \leq t \leq T} \mathcal{D}(t) < \infty$; this follows from the known bounds $\sup_{0 \leq t \leq T} \mathbb{E}(|u_t(x)|^k) < \infty$ and $\sup_{0 \leq t \leq T} \mathbb{E}(|\bar{U}_t^{(\varepsilon)}(x)|^k) < \infty$; see [12]. \square

4 Proof of the remaining results

For the proof of Theorem 2.5, we will need to estimate the Hölder continuity of the random function $t \mapsto \bar{U}_t^{(\varepsilon)}$. The following address this Hölder continuity matter.

Lemma 4.1. *For every real $T > 0$ and integer $k \geq 2$,*

$$\sup_{x \in \varepsilon \mathbf{Z}} \mathbb{E} \left(\left| \bar{U}_t^{(\varepsilon)}(x) - \bar{U}_s^{(\varepsilon)}(x) \right|^k \right) \leq C \left(\frac{t-s}{\varepsilon} \right)^{k/2}, \quad (4.1)$$

uniformly for all $0 < s < t \leq T$.

Proof. Since $\sup_{0 < \varepsilon < 1} \sup_{x \in \varepsilon \mathbf{Z}} \sup_{t \leq T} \mathbb{E}(|\bar{U}_t^{(\varepsilon)}(x)|^k) < \infty$, (BDG) implies that

$$\begin{aligned} &\left\| \bar{U}_t^{(\varepsilon)}(x) - \bar{U}_s^{(\varepsilon)}(x) \right\|_k^2 \\ &\leq \frac{C}{\varepsilon} \int_s^t \sum_{j=-\infty}^{\infty} \left| P_{t-r}^{(\varepsilon)}(j\varepsilon - x) \right|^2 dr + \frac{C}{\varepsilon} \int_0^s \sum_{j=-\infty}^{\infty} \left| P_{t-r}^{(\varepsilon)}(j\varepsilon - x) - P_{s-r}^{(\varepsilon)}(j\varepsilon - x) \right|^2 dr. \end{aligned} \quad (4.2)$$

The first term is bounded by $C(t-s)/\varepsilon$ thanks to (3.45). We appeal to Plancherel's identity—using also the fact that $-1 \leq \hat{\mu}(z) \leq 1$ for all z —in order to obtain the following equivalent representation of the corresponding second term:

$$\begin{aligned}
& \frac{C}{\varepsilon} \int_0^s dr \int_{-\pi}^{\pi} dz \left| \exp\left(-\frac{(t-r)[1-\hat{\mu}(z)]}{\varepsilon^\alpha}\right) - \exp\left(-\frac{(s-r)[1-\hat{\mu}(z)]}{\varepsilon^\alpha}\right) \right|^2 \\
&= \frac{C}{\varepsilon} \int_0^s dr \int_{-\pi}^{\pi} dz \exp\left(-\frac{2(s-r)[1-\hat{\mu}(z)]}{\varepsilon^\alpha}\right) \left| 1 - \exp\left(-\frac{(t-s)[1-\hat{\mu}(z)]}{\varepsilon^\alpha}\right) \right|^2 \\
&\leq \frac{C}{\varepsilon} \int_{-\pi}^{\pi} \frac{\varepsilon^\alpha}{1-\hat{\mu}(z)} \cdot \left| 1 - \exp\left(-\frac{(t-s)[1-\hat{\mu}(z)]}{\varepsilon^\alpha}\right) \right|^2 dz \\
&\leq \frac{C(t-s)}{\varepsilon},
\end{aligned} \tag{4.3}$$

where we used the inequality $1 - e^{-x} \leq \sqrt{x}$, valid for all $x \geq 0$. \square

Proof of Theorem 2.5. The proof is a ready consequence of Theorem 2.4 and a standard application of the Kolmogorov continuity criterion [7, Theorem 4.3, page 10], whose details follow.

Choose and fix some integer $k \in [2, \infty)$. According to [9, p. 566],

$$\mathbb{E}(|u_t(x) - u_s(x)|^k) \leq C|t-s|^{\eta k}, \tag{4.4}$$

uniformly for all $x \in \mathbf{R}$ and $0 \leq s, t \leq T$, where

$$\eta := \frac{\alpha - 1}{2\alpha}. \tag{4.5}$$

Therefore, we appeal to the Kolmogorov continuity theorem and deduce the following bound: For every integer $k \geq 4/(3\eta)$,

$$\sup_{x \in \mathbf{R}} \mathbb{E} \left(\sup_{\substack{0 \leq s, t \leq T \\ |t-s| < \varepsilon^{3T}}} |u_t(x) - u_s(x)|^k \right) \leq C\varepsilon^{3\eta k - 4}, \tag{4.6}$$

simultaneously for every $\varepsilon \in (0, 1)$. Lemma 4.1 and a second appeal to Kolmogorov's continuity theorem, together yield the following:

$$\sup_{x \in \varepsilon \mathbf{Z}} \mathbb{E} \left(\sup_{\substack{0 \leq s, t \leq T \\ |t-s| < \varepsilon^{3T}}} \left| \bar{U}_t^{(\varepsilon)}(x) - \bar{U}_s^{(\varepsilon)}(x) \right|^k \right) \leq C\varepsilon^{k-4} \leq C\varepsilon^{3\eta k - 4}, \tag{4.7}$$

simultaneously for every $\varepsilon \in (0, 1)$. The preceding two observations can be combined as follows. Define

$$D_t^{(\varepsilon)}(x) := u_t(x) - \bar{U}_t^{(\varepsilon)}(x). \tag{4.8}$$

Then

$$\sup_{x \in \varepsilon \mathbf{Z}} \mathbf{E} \left(\sup_{\substack{0 \leq s, t \leq T \\ |t-s| < \varepsilon^3 T}} \left| D_t^{(\varepsilon)}(x) - D_s^{(\varepsilon)}(x) \right|^k \right) \leq C \varepsilon^{3\eta k - 4}, \quad (4.9)$$

simultaneously for every $\varepsilon \in (0, 1)$.

Next, we notice that

$$(0, T] \subseteq \bigcup_{1 \leq j \leq \varepsilon^{-3}} (j\varepsilon^3 T, (j+1)\varepsilon^3 T], \quad (4.10)$$

whence, for every fixed $\nu \in (0, a \wedge [\alpha - 1])$,

$$\begin{aligned} & \mathbf{P} \left\{ \left| u_t(x) - \bar{U}_t^{(\varepsilon)}(x) \right| > \varepsilon^{\nu/2} \text{ for some } x \in \varepsilon \mathbf{Z}, |x| \leq \varepsilon^{-M}, \text{ and } t \in (0, T] \right\} \\ & \leq 2\varepsilon^{-M-1} \sup_{x \in \varepsilon \mathbf{Z}} \mathbf{P} \left\{ \sup_{t \in (0, T]} \left| u_t(x) - \bar{U}_t^{(\varepsilon)}(x) \right| > \varepsilon^{\nu/2} \right\} \\ & \leq J_1 + J_2, \end{aligned} \quad (4.11)$$

where

$$\begin{aligned} J_1 & := 2\varepsilon^{-M-4} \sup_{x \in \varepsilon \mathbf{Z}} \sup_{t \in (0, T]} \mathbf{P} \left\{ \left| u_t(x) - \bar{U}_t^{(\varepsilon)}(x) \right| > \frac{1}{2} \varepsilon^{\nu/2} \right\}, \\ J_2 & := 2\varepsilon^{-M-1} \sup_{x \in \varepsilon \mathbf{Z}} \mathbf{P} \left\{ \sup_{\substack{0 < s, t \leq T \\ |t-s| < \varepsilon^3 T}} \left| D_t^{(\varepsilon)}(x) - D_s^{(\varepsilon)}(x) \right| > \frac{1}{2} \varepsilon^{\nu/2} \right\}. \end{aligned} \quad (4.12)$$

Choose and fix $\rho \in (\nu, a \wedge [\alpha - 1])$ and appeal to Theorem 2.4, and the Chebyshev inequality, in order to see that

$$J_1 \leq C \varepsilon^{-M-4+k(\rho-\nu)/2} \quad \text{as } \varepsilon \downarrow 0, \quad (4.13)$$

where $k \geq 2$ is an integer that we can choose [and fix] as large as we wish. And (4.9) ensures that

$$J_2 \leq C \varepsilon^{-M-5+k(6\eta-\nu)/2} \quad \text{as } \varepsilon \downarrow 0, \quad (4.14)$$

for [say] the same integer k . Since $6\eta > \alpha - 1 > \nu$ and k can be chosen to be as large as needed, we have proved the following: For every $\nu \in (0, a \wedge [\alpha - 1])$ and $Q > 0$,

$$\begin{aligned} & \mathbf{P} \left\{ \left| u_t(x) - \bar{U}_t^{(\varepsilon)}(x) \right| > \varepsilon^{\nu/2} \text{ for some } x \in \varepsilon \mathbf{Z}, |x| \leq \varepsilon^{-M}, \text{ and } t \in (0, T] \right\} \\ & \qquad \qquad \qquad = o(\varepsilon^Q) \quad \text{as } \varepsilon \downarrow 0. \end{aligned}$$

This is more than enough to imply the result. \square

Proof of Theorem 2.6. Choose and fix an arbitrary positive constant β . Let u be the solution to (SHE), and define

$$v_t(x) := u_{\beta t}(x) \quad (t \geq 0, x \in \mathbf{R}). \quad (4.15)$$

Then, v solves the stochastic heat equation,

$$\frac{\partial}{\partial t} v_t(x) = \beta(\mathcal{L}v_t)(x) + \beta^{1/2}\sigma(v_t(x))\tilde{\xi}(t, x), \quad (4.16)$$

subject to $v_0(x) = u_0(x)$ for all x , where $\tilde{\xi} = \tilde{\xi}^{(\beta)}$ is a space-time white noise. This observation shows us that Theorem 2.6 holds for a *fixed* value of $\nu > 0$ if and only if Theorem 2.6 is valid for *every* possible choice of $\nu > 0$. We have seen (Examples 2.2 and 2.3) that there exists $\nu > 0$, together with a random walk model that satisfies Assumption 2.1 for that particular value of ν . [In fact, every $\nu > 0$ works when $\alpha = 2$; see Example 2.2.] We choose and fix that ν and random walk from now on.

The remainder of the proof of the Theorem follows from applying a result of Cox, Fleischmann, and Greven [4, Theorem 1] along with our Theorem 2.4. Theorem 1 of [4] is a comparison of moments for interacting diffusions. For every fixed $\varepsilon, t > 0$ and $x_1, x_2, \dots, x_m \in \mathbf{R}$, that comparison theorem implies that

$$\mathbb{E} \left(\prod_{j=1}^m U_t^{(\varepsilon)}(\varepsilon[x_j/\varepsilon]) \right) \leq \mathbb{E} \left(\prod_{j=1}^m U_t^{1,(\varepsilon)}(\varepsilon[x_j/\varepsilon]) \right) \quad (4.17)$$

where $U_t^{1,(\varepsilon)}$ solves (2.13) with σ replaced by $\bar{\sigma}$. Our assumption that $\sigma(0) = \bar{\sigma}(0) = 0$ guarantees that both $U_t^{(\varepsilon)}$ and $U_t^{1,(\varepsilon)}$ remain nonnegative; this follows from the well-known comparison principle for interacting diffusions; see for example, Geiß and Manthey [11, Theorem 1.1], Mueller [21, Theorem 3.1], or Georgiou et al [12]. The support of $U_t^{(\varepsilon)}(x)$ and $U_t^{1,(\varepsilon)}(x)$ is thus \mathbf{R}_+ and so the conditions of Cox et al [4, Theorem 1] are verified. Now let $\varepsilon \downarrow 0$ in (4.17) and appeal to our Theorem 2.4 to conclude the proof. \square

Proof of Theorem 2.7. The upper bound of $\text{Lip}_\sigma^4 k^3 / \nu$ follows immediately from Example 2.9 of Foondun and Khoshnevisan [9]. We will establish the lower bound next. Let $\theta \in \mathbf{R}$ be fixed, and consider the solution $X_t(x)$ to the *parabolic Anderson model*,

$$\frac{\partial}{\partial t} X_t(x) = \nu \frac{\partial^2}{\partial x^2} X_t(x) + L_\sigma X_t(x) \xi(t, x), \quad (4.18)$$

subject to $X_0(x) \equiv 1$ for all x . That is, $(t, x) \mapsto X_t(x)$ solves (SHE) with $\sigma(x) = L_\sigma x$. According to Theorem 2.6,

$$\mathbb{E} (|u_t(x)|^k) \geq \mathbb{E} (|X_t(x)|^k) \geq \exp \left(\frac{L_\sigma^4 k(k^2 - 1)t}{48\nu} \right), \quad (4.19)$$

pointwise. See Theorem of 6.4 of Khoshnevisan [19, Theorem 6.4] for the last bound. This proves the theorem. \square

Acknowledgements. We thank Nicos Georgiou for many enjoyable conversations on this topic and Gennady Samorodnitsky for his helpful comments on the topic of Example 2.3. Many thanks are also due to Ivan Corwin and Roger Tribe for their question about Lyapunov estimates; that question is answered here in Theorem 2.7.

References

- [1] Bertini, Lorenzo and Nicoletta Cancrini, The stochastic heat equation: Feynman-Kac formula and intermittence, *J. Statist. Phys.* **78**(5-6) (1995) 1377–1401.
- [2] Bochner, Salomon, *Harmonic Analysis and the Theory of Probability*, Dover Publication, Inc., Mineola, New York, 2005 (Reprint of the original, published by the University of California Press, 1955).
- [3] Borodin, Alexei and Ivan Corwin, On the moments of the parabolic Anderson model, to appear in *Ann. Applied Probab.*
- [4] Cox, J. Theodore, Klaus Fleischmann and Andreas Greven, Comparison of interacting diffusions and an application to their ergodic theory, *Probab. Theory Related Fields* **105**(4) (1996) 513–528.
- [5] Csörgő, Miklos and Pál Révész, *Strong Approximations in Probability and Statistics*, Academic Press, New York, 1981.
- [6] Dalang, Robert C., Extending the martingale measure stochastic integral with applications to spatially homogeneous s.p.d.e.’s, *Electron. J. Probab.* **4**, Paper no. 6 (1999) 29 pp. (electronic). Available electronically at <http://www.math.washington.edu/~ejpecp>.
- [7] Dalang, Robert, Davar Khoshnevisan, Carl Mueller, David Nualart, and Yimin Xiao, A minicourse on stochastic partial differential equations (D. Khoshnevisan and F. Rassoul-Agha, Ed’s) *Lecture Notes in Mathematics* **1962**, 2009.
- [8] Davis, Burgess, On the L^p norms of stochastic integrals and other martingales, *Duke Math. J.* **43**(4) (1976) 697704.
- [9] Foondun, Mohammud and Davar Khoshnevisan, Intermittence and nonlinear parabolic stochastic partial differential equations, *Electr. J. Probab.* **14**, Paper no. 12 (2009) 548–568 (electronic). Available electronically at <http://www.math.washington.edu/~ejpecp>.
- [10] Funaki, Tadahisa, Random motion of strings and related stochastic evolution equations, *Nagoya Math. J.* **89** (1983) 129–193.
- [11] Geiß, Christel and Ralf Manthey, Comparison theorems for stochastic differential equations in finite and infinite dimensions, *Stoch. Processes and Their Applications* **53** (1994) 23–35.
- [12] Georgiou, Nicos, Mathew Joseph, Davar Khoshnevisan, and Shang-Yuan Shiu, Semi-discrete semi-linear parabolic SPDEs, to appear in *Ann. Applied Probab.*
- [13] Głowacki, Paweł, Stable semigroups of measures as commutative approximate identities on nongraded homogeneous groups, *Invent. Math.* **83** (1986) 557–582.
- [14] Gyöngy, I., Krylov, N.V., Accelerated numerical schemes for PDEs and SPDEs, *Stochastic analysis* (2010) 131–168.

- [15] Gyöngy, I., Krylov, N.V., On Splitting up method and stochastic partial differential equations, *Ann. Probab.* **31** (2003) 564–591.
- [16] Gyöngy, I., Krylov, N.V., On the rate of convergence of splitting-up approximations for SPDEs, *Prog. Probab.* **56** (2003) 301–321.
- [17] Kardar, Mehran, Replica-Bethe Ansatz studies of two-dimensional interfaces with quenched random impurities, *Nucl. Phys. B* **290** (2007) 582–602.
- [18] Khoshnevisan, Davar, Lévy classes and self-normalization, *Electr. J. Probab.* **1**(1) (1996) 1–18.
- [19] Khoshnevisan, Davar, *Analysis of Stochastic Partial Differential Equations*, Conference Board of the Mathematical Sciences, American Mathematical Society, Providence RI (to appear).
- [20] Lawler, Gregory F. and Vlada Limic, *Random Walk: A Modern Introduction*, Cambridge University Press, Cambridge, UK, 2010.
- [21] Mueller, Carl, On the support of solutions to the heat equation with noise, *Stochastics Stochastics Rep.* **37** no. 4, (1991) 225–245.
- [22] Peszat, Szymon, and Jerzy Zabczyk, Nonlinear stochastic wave and heat equations, *Probab. Theory Rel. Fields* **116** 421–443.
- [23] Shiga, Tokuzo and Akinobu Shimizu, Infinite-dimensional stochastic differential equations and their applications, *J. Math. Kyoto U.* **20**(3) (1980) 395–416.
- [24] Spitzer, Frank, *Principles of Random Walk* (second edition) Springer, New York, 1976.
- [25] Walsh, John B., *An Introduction to Stochastic Partial Differential Equations*, in: *École d’été de probabilités de Saint-Flour, XIV—1984*, 265–439, Lecture Notes in Math., vol. 1180, Springer, Berlin, 1986.

Mathew Joseph (m.joseph@sheffield.ac.uk).

Department of Probability and Statistics, University of Sheffield, Sheffield, England, S3 7RH, UK

Davar Khoshnevisan (davar@math.utah.edu).

Department of Mathematics, University of Utah, Salt Lake City, UT 84112-0090, USA

Carl Mueller (carl.2013@outlook.com).

Department of Mathematics, University of Rochester, Rochester, NY 14627, USA