

# Math 162: Calculus IIA

## Final Exam ANSWERS

December 14, 2021

### HANDY DANDY FORMULAS

Integration by parts formula:

$$\int u dv = uv - \int v du$$

Trigonometric identities:

$$\cos^2 \theta + \sin^2 \theta = 1$$

$$\sec^2 \theta - \tan^2 \theta = 1$$

$$\sin(\alpha + \beta) = \sin \alpha \cos \beta + \cos \alpha \sin \beta$$

$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta$$

$$\cos^2 \theta = \frac{1 + \cos 2\theta}{2}$$

$$\sin^2 \theta = \frac{1 - \cos 2\theta}{2}$$

Derivatives of trig functions.

$$\frac{d \sin x}{dx} = \cos x$$

$$\frac{d \tan x}{dx} = \sec^2 x$$

$$\frac{d \sec x}{dx} = \sec x \tan x$$

$$\frac{d \cos x}{dx} = -\sin x$$

$$\frac{d \cot x}{dx} = -\csc^2 x$$

$$\frac{d \csc x}{dx} = -\csc x \cot x$$

Trigonometric substitution for integrals of the form

$$\int \tan^m x \sec^n x dx \quad \text{with } n > 0,$$

known in Doug's section as *the rabbit trick*.

$$u = \sec x + \tan x$$

$$\sec x dx = \frac{du}{u}$$

$$\sec x = \frac{u^2 + 1}{2u}$$

$$\tan x = \frac{u^2 - 1}{2u}$$

Area of surface of revolution in rectangular coordinates,  $y = f(x)$  with  $a \leq x \leq b$

• about the  $x$ -axis: 
$$S = 2\pi \int_a^b f(x) \sqrt{1 + f'(x)^2} dx$$

• about the  $y$ -axis: 
$$S = 2\pi \int_a^b x \sqrt{1 + f'(x)^2} dx$$

## MORE FORMULAS FOR YOUR ENJOYMENT

## Polar coordinates

$$\begin{aligned}
 r &= \sqrt{x^2 + y^2} & \theta &= \arctan(y/x) & \text{for } x > 0 \\
 \pi + \arctan(y/x) & \text{for } x < 0 \\
 \pi/2 & \text{for } x = 0 \text{ and } y > 0 \\
 3\pi/2 & \text{for } x = 0 \text{ and } y < 0 \\
 \text{undefined} & \text{for } (x, y) = (0, 0) \\
 x &= r \cos \theta & y &= r \sin \theta
 \end{aligned}$$

Changing  $\theta$  by any multiple of  $2\pi$  does not change the location of the point. Changing the sign of  $r$  is equivalent to adding  $\pi$  to  $\theta$ , which is the same as moving the point to one in the opposite direction and the same distance from the origin.

Area in polar coordinates for  $r = f(\theta)$  with  $\alpha \leq \theta \leq \beta$ :

$$A = \int_{\alpha}^{\beta} \frac{r^2}{2} d\theta$$

## Arc length formulas

- Rectangular coordinates,  $y = f(x)$  with  $a \leq x \leq b$ :

$$S = \int_a^b \sqrt{1 + f'(x)^2} dx$$

- Polar coordinates,  $r = f(\theta)$  with  $\alpha \leq \theta \leq \beta$ :

$$S = \int_{\alpha}^{\beta} \sqrt{r^2 + f'(\theta)^2} d\theta$$

- Parametric equations,  $x = x(t)$  and  $y = y(t)$  with  $a \leq t \leq b$ :

$$S = \int_a^b \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

## INFINITE SERIES FORMULAS

The Maclaurin series for  $f(x)$  is

$$\sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{n!} x^n.$$

The Taylor series for  $f(x)$  at  $a$  is

$$\sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x - a)^n.$$

The  $m$ th Taylor polynomial is

$$T_m(x) = \sum_{n=0}^m \frac{f^{(n)}(a)}{n!} (x - a)^n,$$

and the  $m$ th Taylor remainder is

$$R_m(x) = f(x) - T_m(x)$$

Taylor's inequality says that if  $|f^{(n+1)}(x)| \leq M$  for suitable  $x$ , then

$$|R_m(x)| \leq \frac{(x - a)^{n+1} M}{(n + 1)!}.$$

**Part A****1. (20 points)**

(a) (10 points) The form of the partial fraction decomposition of the function is given below:

$$\frac{3x^2 + 3x + 3}{(x - 3)(x^2 + 4)} = \frac{A}{x - 3} + \frac{Bx + C}{x^2 + 4}.$$

Find the coefficients  $A$ ,  $B$  and  $C$ .

**Answer:**

$$\begin{aligned}\frac{3x^2 + 3x + 3}{(x - 3)(x^2 + 4)} &= \frac{A}{x - 3} + \frac{Bx + C}{x^2 + 4} \\ 3x^2 + 3x + 3 &= A(x^2 + 4) + (Bx + C)(x - 3).\end{aligned}$$

Plug in  $x = 3$  :

$$27 + 9 + 3 = 39 = 13A, \Rightarrow A = 3.$$

$$\text{Therefore, } 3x^2 + 3x + 3 = 3(x^2 + 4) + (Bx + C)(x - 3) = (3 + B)x^2 + (C - 3B)x + 12 - 3C.$$

Matching coefficients of appropriate degrees of  $x$ , we get  $3 = 3 + B$  and  $3 = 12 - 3C$ .

So  $B = 0$  and  $C = 3$ .

(b) (10 points) Evaluate the following integral:

$$\int \frac{3x^2 + 3x + 3}{(x - 3)(x^2 + 4)} dx.$$

**Answer:**

From part (a):

$$\begin{aligned}\int \frac{3x^2 + 3x + 3}{(x - 3)(x^2 + 4)} dx \\ = \int \frac{3}{x - 3} + \frac{3}{x^2 + 4} dx.\end{aligned}$$

$$\int \frac{3}{x-3} dx = 3 \ln |x-3| + C.$$

For  $\int \frac{3}{x^2+4} dx$  we will use trig substitution:

$\tan(\theta) = x/2 \Rightarrow 2 \tan(\theta) = x \Rightarrow 2 \sec^2(\theta) d\theta = dx$ . Also,  $\cos(\theta) = \frac{1}{\sqrt{x^2+4}}$ , so  $\cos^2(\theta) = \frac{1}{x^2+4}$ . After substituting,

$$\int \frac{3}{x^2+4} dx = 3 \int \cos^2(\theta) \sec^2(\theta) d\theta = 3 \int d\theta = 3\theta + C = 3 \arctan(x/2) + C.$$

So in total,

$$\int \frac{3}{x-3} + \frac{3}{x^2+4} dx = 3 \ln |x-3| + 3 \arctan(x/2) + C.$$

**2. (20 points)** Consider the solid  $\mathcal{R}$  formed by revolving the function  $y = \sqrt{4-x^2}$  around the  $x$ -axis for  $0 \leq x \leq 2$ .

(a) (10 points) Compute the volume of  $\mathcal{R}$ .

**Answer:**

Using the Disk Method, the volume is

$$\begin{aligned} & \pi \int_0^2 4 - x^2 dx \\ &= \pi \left[ 4x - \frac{x^3}{3} \right]_0^2 \\ &= \frac{16\pi}{3} \end{aligned}$$

(b) (10 points) Compute the surface area of  $\mathcal{R}$ .

**Answer:**

The formula for the surface area is

$$\begin{aligned}
S &= 2\pi \int_0^2 y \sqrt{1 + (y')^2} dx \\
&= 2\pi \int_0^2 \sqrt{4 - x^2} \sqrt{1 + \left(\frac{-x}{\sqrt{4 - x^2}}\right)^2} dx \\
&= 2\pi \int_0^2 \sqrt{4 - x^2} \sqrt{1 + \frac{x^2}{4 - x^2}} dx \\
&= 2\pi \int_0^2 \sqrt{4 - x^2} \sqrt{\frac{4 - x^2}{4 - x^2} + \frac{x^2}{4 - x^2}} dx \\
&= 2\pi \int_0^2 \sqrt{4 - x^2} \sqrt{\frac{4}{4 - x^2}} dx \\
&= 2\pi \int_0^2 2 dx = 2\pi [2x]_0^2 = 8\pi.
\end{aligned}$$

3. (10 points) Compute the following integral:

$$\int x \ln(x) dx$$


**Answer:**

We will do integration by parts:  $u = \ln(x)$ ,  $dv = x dx$ ,  $\Rightarrow du = 1/x dx$ ,  $v = x^2/2$ .

$$\begin{aligned}
\int x \ln(x) dx &= \frac{x^2 \ln(x)}{2} - \int \frac{x}{2} dx \\
&= \frac{x^2 \ln(x)}{2} - \frac{x^2}{4} + C.
\end{aligned}$$

4. (20 points)

Find the arc length  $L$  of the polar curve,  $r = e^{\theta/4}$ , from  $\theta = 0$  to  $\theta = 4\pi$ .



**Answer:**

For future reference note that  $dr/d\theta = e^{\theta/4}/4$ . The polar arc length formula gives

$$\begin{aligned}
 s &= \int_0^{4\pi} \sqrt{r^2 + (dr/d\theta)^2} d\theta \\
 &= \int_0^{4\pi} \sqrt{e^{\theta/2} + e^{\theta/2}/16} d\theta \\
 &= \sqrt{\frac{17}{16}} \int_0^{4\pi} e^{\theta/4} d\theta = \frac{\sqrt{17}}{4} 4e^{\theta/4} \Big|_0^{4\pi} \\
 &= \sqrt{17}(e^\pi - 1).
 \end{aligned}$$

**5. (20 points)**

(a) (10 points) Compute the volume of a region bounded by the curves  $y = x^5 + 1$ ,  $y = 1$  and  $x = 1$  and rotated around the  $y$ -axis.

**Answer:**

Using the shell method we have shells of radius  $x$ , thickness  $dx$  and height  $(x^5 + 1) - 1 = x^5$ . Therefore

$$V = \int_0^1 2\pi x \cdot x^5 dx = 2\pi \frac{x^7}{7} \Big|_0^1 = \frac{2\pi}{7}$$

(b) (10 points) Find the volume of the region bounded by  $y = x^3$ ,  $y = 0$  and  $x = 1$  and rotated around line  $x = 2$ . Use the shell method.

**Answer:**

Using the shell method we have shells of radius  $(2 - x)$ , thickness  $dx$  and height  $x^3$ . Thus the volume is

$$\begin{aligned} V &= \int_0^1 2\pi(2-x)x^3 dx \\ &= \pi \int_0^1 4x^3 - 2x^4 dx \\ &= \pi \left( x^4 - \frac{2x^5}{5} \right) \Big|_0^1 \\ &= \pi \left( 1 - \frac{2}{5} \right) = \frac{3\pi}{5} \end{aligned}$$

6. (20 points) Compute

$$\int \frac{x^2}{(1-4x^2)^{3/2}} dx$$

**Answer:**

Consider the right triangle with hypotenuse 1 and sides  $2x$  and  $\sqrt{1-4x^2}$ . Let  $\theta$  be the angle opposite  $2x$ . Then we have

$$x = \frac{\sin \theta}{2} \qquad dx = \frac{\cos \theta d\theta}{2} \qquad \sqrt{1-4x^2} = \cos \theta.$$


Writing the integral in term of  $\theta$ ,

$$\begin{aligned} \int \frac{x^2}{(1-4x^2)^{3/2}} dx &= \int \frac{(\sin^2 \theta / 4) \cos \theta d\theta}{\cos^3 \theta \cdot 2} \\ &= \frac{1}{8} \int \frac{\sin^2 \theta d\theta}{\cos^2 \theta} \\ &= \frac{1}{8} \int \tan^2 \theta d\theta \\ &= \frac{1}{8} \int (\sec^2 \theta - 1) d\theta \\ &= \frac{1}{8} (\tan \theta - \theta) + c \\ &= \frac{1}{8} \left( \frac{2x}{\sqrt{1-4x^2}} - \arcsin 2x \right) + c \end{aligned}$$

7. (20 points)



(a) (10 points) Find the area of the region both inside the circle  $r = \sin \theta$  and outside the circle  $r = \cos \theta$  (both equations are in polar coordinates). The two circles are shown below. THEY INTERSECT AT THE ORIGIN AND THE POLAR POINT  $(\theta, r) = (\pi/4, \sqrt{2}/2)$ .



**Answer:**

Find the area of the region inside the first circle and outside the second by integrating

$$\begin{aligned} \int_{\pi/4}^{\pi} \frac{r^2}{2} d\theta &= \frac{1}{2} \int_{\pi/4}^{\pi} \sin^2 \theta d\theta = \frac{1}{4} \int_{\pi/4}^{\pi} (1 - \cos 2\theta) d\theta \\ &= \frac{1}{4} \left( \frac{3\pi}{4} - \frac{\sin 2\pi}{2} + \frac{\sin(\pi/2)}{2} \right) = \frac{1}{4} \left( \frac{3\pi}{4} + \frac{1}{2} \right) = \frac{3\pi}{16} + \frac{1}{8} \end{aligned}$$

and subtracting

$$\begin{aligned} \int_{\pi/4}^{\pi/2} \frac{r^2}{2} d\theta &= \frac{1}{2} \int_{\pi/4}^{\pi/2} \cos^2 \theta d\theta = \frac{1}{4} \int_{\pi/4}^{\pi/2} (1 + \cos 2\theta) d\theta \\ &= \frac{1}{4} \left( \frac{\pi}{4} + \frac{\sin \pi}{2} - \frac{\sin(\pi/2)}{2} \right) = \frac{1}{4} \left( \frac{\pi}{4} - \frac{1}{2} \right) \\ &= \frac{\pi}{16} - \frac{1}{8}. \end{aligned}$$

So the area of the region is

$$\left( \frac{3\pi}{16} + \frac{1}{8} \right) - \left( \frac{\pi}{16} - \frac{1}{8} \right) = \frac{\pi}{8} + \frac{1}{4}.$$

(b) (10 points) Compute the equation (in Cartesian coordinates  $x, y$ ) of the tangent line to the circle  $r = \sin \theta$  at the points where it intersects the circle  $r = \cos \theta$

**Answer:**

There are two intersection points, at  $(x, y) = (0, 0)$  and  $(x, y) = (\sqrt{2}/2, \sqrt{2}/2)$ , at which the tangent lines to the upper circle are horizontal with equation  $y = 0$  and vertical with equation  $x = \sqrt{2}/2$  respectively.

**Part B**

**8. (20 points)** Let  $q$  be a positive (greater than 0) real number.

(a) (10 points)

Find the radius of convergence of the series  $\sum_{n=0}^{\infty} q^{2n}(x - \pi)^n$ .

**Answer:**

Applying the ratio test, we have

$$\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{q^{2(n+1)}(x - \pi)^{n+1}}{q^{2n}(x - \pi)^n} \right| = \lim_{n \rightarrow \infty} q^2|x - \pi| = q^2|x - \pi|$$

As  $q^2|x - \pi| < 1$  if and only if  $|x - \pi| < 1/q^2$ , we can conclude that the radius of convergence is  $1/q^2$ .

(b) (10 points) Find the interval of convergence of the series  $\sum_{n=0}^{\infty} q^{2n}(x - \pi)^n$ .

**Answer:**

To determine the interval of convergence, we plug in  $x = \pi \pm 1/q^2$  into the original expression. For  $x = \pi + 1/q^2$ , the series becomes  $\sum_{n=0}^{\infty} q^{2n}(1/q^2)^n = \sum_{n=0}^{\infty} 1$ , which diverges; for  $x = \pi - 1/q^2$ , the series becomes  $\sum_{n=0}^{\infty} q^{2n}(-1/q^2)^n = \sum_{n=0}^{\infty} (-1)^n$ , which also diverges. Hence, the interval of convergence is  $(\pi - 1/q^2, \pi + 1/q^2)$ .

**9. (20 points)**

(a) (10 points) Consider the series  $\sum_{n=1}^{\infty} (-1)^n \sqrt{\frac{1}{n^2} + 1}$ . Is this series conditionally convergent, absolutely convergent, or divergent? Explain your answer.

**Answer:**

The series is divergent, since  $\lim_{n \rightarrow \infty} (-1)^n \sqrt{\frac{1}{n^2} + 1}$  does not exist.

- (b) (10 points) The series  $\sum_{n=1}^{\infty} \frac{(-2)^n}{n}$  converges conditionally. How many terms do you need to estimate the sum with an accuracy of  $1/1000$ ?

[The series actually diverges. IGNORE THIS PROBLEM.]

**Answer:**

**10. (20 points)**

- (a) (10 points) Find a power series expansion of  $f(x) = \frac{1}{x}$  centered at  $x = 1$ .

**Answer:**

Since  $\frac{1}{x} = \frac{1}{1 + (x - 1)} = \frac{1}{1 - (-(x - 1))}$ , we can use the geometric series expansion to get

$$\frac{1}{x} = \sum_{n=0}^{\infty} (-1)^n (x - 1)^n.$$

- (b) (10 points) Use your series from (a) to find a power series expansion of  $\frac{1}{x^2}$  centered at  $x = 1$ .

**Answer:**

As  $\frac{1}{x^2} = -\frac{d}{dx} \left( \frac{1}{x} \right)$ , we can differentiate our series from (a) to get

$$\frac{1}{x^2} = -\sum_{n=1}^{\infty} n(-1)^n (x - 1)^{n-1} = \sum_{n=1}^{\infty} n(-1)^{n-1} (x - 1)^{n-1}.$$

**11. (20 points)**

- (a) (10 points) Show that the following series converges:

$$\sum_{n=1}^{\infty} \frac{1}{n \cdot 5^n}$$

**Answer:**

For all  $n \geq 1$ ,  $\frac{1}{n \cdot 5^n} \leq \frac{1}{5^n}$ .  $\sum_{n=1}^{\infty} \frac{1}{5^n}$  converges because it is a geometric series. Therefore, by the Comparison Test,  $\sum_{n=1}^{\infty} \frac{1}{n \cdot 5^n}$  also converges.

(b) **(5 Points)** Find the Maclaurin power series representation for  $-\ln|1-x|$ . (Hint: What is the Maclaurin series for  $1/(1-x)$ ?)

**Answer:**

$\frac{1}{1-x} = \sum_{n=0}^{\infty} x^n$ . Taking the antiderivative of both sides, we get

$$-\ln|1-x| = \sum_{n=1}^{\infty} \frac{x^n}{n}.$$

(c) **(5 Points)** Find the value of the series of (a) in terms of the natural logarithm.

**Answer:**

Plug in  $\frac{1}{5}$  for  $x$  in the Maclaurin series for  $-\ln|1-x|$  to get

$$\sum_{n=1}^{\infty} \frac{1}{n \cdot 5^n} = -\ln(4/5) = \ln(5/4).$$

This is scratch paper. If you use it to work on a problem, please indicate so on the page where that problem occurs.

---

Second scratch paper page. If you use it to work on a problem, please indicate so on the page where that problem occurs.

---

Third scratch paper page. If you use it to work on a problem, please indicate so on the page where that problem occurs.

---

Fourth scratch paper page. If you use it to work on a problem, please indicate so on the page where that problem occurs.

More scratch paper. If you use it to work on a problem, please indicate so on the page where that problem occurs.



---

And even more scratch paper. If you use it to work on a problem, please indicate so on the page where that problem occurs.