

Formal Group Laws

In this appendix we will give a self-contained account of the relevant aspects of the theory of commutative one-dimensional formal group laws. This theory was developed by various algebraists for reasons having nothing to do with algebraic topology. The bridge between the two subjects is the famous result of Quillen [2] (4.1.6) which asserts that the Lazard ring L (A2.1.8) over which the universal formal group law is defined is naturally isomorphic to the complex cobordism ring. A most thorough and helpful treatment of this subject is given in Hazewinkel [1]. An account of the Lazard ring is also given in Adams [5], while the classification in characteristic p can also be found in Fröhlich [1].

We now outline the main results of Section 1. We define formal group laws (A2.1.1) and homomorphisms between them (A2.1.5) and show that over a field of characteristic 0 every formal group law is isomorphic to the additive one (A2.1.6). The universal formal group law is constructed (A2.1.8) and the structure of the ring L over which it is defined is determined (A2.1.10). This result is originally due to Lazard [1]. Its proof depends on a difficult lemma (A2.1.12) whose proof is postponed to the end of the section.

Then we define p -typical formal group laws (A2.1.17 and A2.1.22) and determine the structure of the p -typical analog of the Lazard ring, V (A2.1.24). This result is due to Carrier [1]; Quillen [2] showed that V is naturally isomorphic to $\pi^*(BP)$ (4.1.12). Using a point of view due to Landweber [1], we determine the structure of algebraic objects LB (A2.1.16) and VT (A2.1.26), which turn out to be isomorphic to $MU_*(MU)$ (4.1.11) and $BP_*(BP)$ (4.1.19), respectively.

All of the results of this section can be found in Adams [5], although our treatment of it differs from his.

In Section 2 we give the explicit generators of V [i.e., of $\pi_*(BP)$] given by Hazewinkel [2] (A2.2.1) and Araki [1] (A2.2.2) and determine the behavior of the right unit η_R on Araki's generators (A2.2.5).

For the Morava theory of Chapter 6 we will need the classification of formal group laws over separably closed fields of characteristic $p > 0$ (A2.2.11) originally due to Lazard [2], and a description of the relevant endomorphism rings (A2.2.17 and A2.2.18) originally due to Dieudonné [1] and Lubin [1].

For a scheme theoretic approach to this subject, see Strickland [?].

1. Universal Formal Group Laws and Strict Isomorphisms

A2.1.1. DEFINITION. *Let R be a commutative ring with unit. A formal group law over R is a power series $F(x, y) \in R[[x, y]]$ satisfying*

- (i) $F(x, 0) = F(0, x) = x$,
- (ii) $F(x, y) = F(y, x)$, and
- (iii) $F(x, F(y, z)) = F(F(x, y), z)$.

Strictly speaking, such an object should be called a commutative one-dimensional formal group law; we omit the first two adjectives as this is the only type of formal group law we will consider. It is known (Lazard [3]) that (ii) is redundant if R has no nilpotent elements.

The reason for this terminology is as follows. Suppose G is a one-dimensional commutative Lie group and $g: \mathbf{R} \rightarrow U \subset G$ is a homomorphism to a neighborhood U of the identity which sends 0 to the identity. Then the group operation $G \times G \rightarrow G$ can be described locally by a real-valued function of two real variables. If the group is analytic then this function has a power series expansion about the origin that satisfies (i)–(iii). These three conditions correspond, respectively, to the identity, commutativity, and associativity axioms of the group. In terms of the power series, the existence of an inverse is automatic, i.e.,

A2.1.2. PROPOSITION. *If F is a formal group law over R then there is a power series $i(x) \in R[[x]]$ (called the formal inverse) such that $F(x, i(x)) = 0$.*

In the Lie group case this power series must of course converge, but in the formal theory convergence does not concern us. Formal group laws arise in more algebraic situations; e.g., one can extract a formal group law from an elliptic curve defined over R ; see Chapter 7 of Silverman [1]. One can also reverse the procedure and get a group out of a formal group law; if R is a complete local ring then $F(x, y)$ will converge whenever x and y are in the maximal ideal, so a group structure is defined on the latter which may differ from the usual additive one.

Before proceeding further note that A2.1.1(i) implies

A2.1.3. PROPOSITION. *If F is a formal group law then*

$$F(x, y) \equiv x + y \pmod{(x, y)^2}. \quad \square$$

A2.1.4. EXAMPLES OF FORMAL GROUP LAWS. (a) $F_a(x, y) = x + y$, the additive formal group law.

(b) $F(x, y) = x + y + uxy$ (where u is a unit in R), the multiplicative formal group law, so named because $1 + uF = (1 + ux)(1 + uy)$.

(c) $F(x, y) = (x + y)/(1 + xy)$.

(d) $F(x, y) = (x\sqrt{1 - y^4} + y\sqrt{1 - x^4})/(1 + x^2y^2)$, a formal group law over $\mathbf{Z}[1/2]$.

The last example is due to Euler and is the addition formula for the elliptic integral

$$\int_0^x \frac{dt}{\sqrt{1 - t^4}}$$

(see Siegel [1, pp. 1-9]). These examples will be studied further below (A2.2.9).

The astute reader will recognize (c) as the addition formula for the hyperbolic tangent function; i.e., if $x = \tanh(u)$ and $y = \tanh(v)$ then $F(x, y) = \tanh(u + v)$. Hence we have

$$\tanh^{-1}(F(x, y)) = \tanh^{-1}(x) + \tanh^{-1}(y)$$

or

$$F(x, y) = \tanh(\tanh^{-1}(x) + \tanh^{-1}(y)),$$

where $\tanh^{-1}(x) = \sum_{i \geq 0} x^{2i+1}/(2i+1) \in R \otimes \mathbf{Q}[[x]]$.

We have a similar situation in (b), i.e.,

$$\log(1 + uF) = \log(1 + ux) + \log(1 + uy),$$

where $\log(l + ux) = \sum_{i>0} (-1)^{i+1} (ux)^i / i \in R \otimes \mathbf{Q}[[x]]$.

This means that the formal group laws of (b) and (c) are isomorphic over \mathbf{Q} to the additive formal group law (a) in the following sense.

A2.1.5. DEFINITION. *Let F and G be formal group laws. A homomorphism from F to G is a power series $f(x) \in R[[x]]$ with constant term 0 such that $f(F(x, y)) = G(f(x), f(y))$. It is an isomorphism if it is invertible, i.e., if $f'(0)$ (the coefficient of x) is a unit in R , and a strict isomorphism if $f'(0) = 1$. A strict isomorphism from F to the addition formal group law $x + y$ is a logarithm for F , denoted by $\log_F(x)$.*

Hence the logarithms for A2.1.4(b) and (c) are

$$\sum_{i>0} \frac{(-u)^{i-1} x^i}{i} \quad \text{and} \quad \tanh^{-1}(x)$$

respectively.

On the other hand, these formal group laws are not isomorphic to the additive one over \mathbf{Z} . To see this for (b), set $u = 1$. Then $F(x, x) = 2x + x^2 \equiv x^2 \pmod{2}$, while $F_a(x, x) = 2x \equiv 0 \pmod{2}$, so the two formal group laws are not isomorphic over $\mathbf{Z}/(2)$. The formal group law of (c) is isomorphic to F_a over $\mathbf{Z}/(2)$, since its logarithm $\tanh^{-1}x$ has coefficients in $\mathbf{Z}/(2)$, but we have $F(F(x, x), x) = (3x + x^3)/(1 + 3x^2) \equiv x^3 \pmod{3}$ while $F_a(F_a(x, x), x) = 3x \equiv 0 \pmod{3}$. Similarly, it can be shown that F and F_a are distinct at every odd prime (see A2.2.9).

A2.1.6. THEOREM. *Let F be a formal group law and let $f(x) \in R \otimes \mathbf{Q}[[x]]$ be given by*

$$f(x) = \int_0^x \frac{dt}{F_2(t, 0)}$$

where $F_2(x, y) = \partial F / \partial y$. Then f is a logarithm for F , i.e., $F(x, y) = f^{-1}(f(x) + f(y))$, and F is isomorphic over $R \otimes \mathbf{Q}$ to the additive formal group law.

PROOF. Let $w = f(F(x, y)) - f(x) - f(y)$. We wish to show $w = 0$. We have $F(F(x, y), z) = F(x, F(y, z))$. Differentiating with respect to z and setting $z = 0$ we get

$$(A2.1.7) \quad F_2(F(x, y), 0) = F_2(x, y)F_2(y, 0).$$

On the other hand, we have $\partial w / \partial y = f'(F(x, y))F_2(y, 0) - f'(y)$, which by the definition of f becomes

$$\frac{\partial w}{\partial y} = \frac{F_2(x, y)}{F_2(F(x, y), 0)} - \frac{1}{F_2(y, 0)} = 0 \quad \text{by A2.1.7.}$$

By symmetry we also have $\partial w / \partial x = 0$, so w is a constant. But f and F both have trivial constant terms, so $w = 0$. \square

Now we wish to consider the universal formal group law. Its construction is easy.

A2.1.8. THEOREM. *There is a ring L (called the Lazard ring) and a formal group law*

$$F(x, y) = \sum a_{i,j} x^i y^j$$

defined over it such that for any formal group law G over any commutative ring with unit R there is a unique ring homomorphism $\theta: L \rightarrow R$ such that $G(x, y) = \sum \theta(a_{i,j})x^i y^j$.

PROOF. Simply set $L = \mathbf{Z}[a_{i,j}]/I$, where I is the ideal generated by the relations among the $a_{i,j}$ required by the definition A2.1.1, i.e., by $a_{1,0} - 1$, $a_{0,1} - 1$, $a_{i,0}$, and $a_{0,i}$ for (i), $a_{i,j} - a_{ji}$ for (ii), and b_{ijk} for (iii), where

$$F(F(x, y), z) - F(x, F(y, z)) = \sum b_{ijk}x^i y^j z^k.$$

Then θ can be defined by the equation it is supposed to satisfy. \square

Determining the structure of L explicitly is more difficult. At this point it is convenient to introduce a grading on L by setting $|a_{i,j}| = 2(i + j - 1)$. Note that if we have $|x| = |y| = -2$ then $F(x, y)$ is a homogeneous expression of degree -2 .

A2.1.9. LEMMA. (a) $L \otimes \mathbf{Q} = \mathbf{Q}[m_1, m_2, \dots]$ with $|m_i| = 2i$ and $F(x, y) = f^{-1}(f(x) + f(y))$ where $f(x) = x + \sum_{i>0} m_i x^{i+1}$.

(b) Let $M \subset L \otimes \mathbf{Q}$ be $\mathbf{Z}[m_1, m_2, \dots]$. Then $\text{im } L \subset M$.

PROOF. (a) By A2.1.6 every formal group law G over a \mathbf{Q} -algebra R has a logarithm $g(x)$ so there is a unique $\phi: \mathbf{Q}[m_1, m_2, \dots] \rightarrow R$ such that $\phi(f(x)) = g(x)$. In particular we have $\phi: \mathbf{Q}[m_1, m_2, \dots] \rightarrow L \otimes \mathbf{Q}$ as well as $\theta: L \otimes \mathbf{Q} \rightarrow \mathbf{Q}[m_1, m_2, \dots]$ with $\theta\phi$ and $\phi\theta$ being identity maps, so θ and ϕ are isomorphisms.

(b) $F(x, y)$ is a power series with coefficients in M , so the map from L to $L \otimes \mathbf{Q}$ factors through M . \square

Now recall that if R is a graded connected ring (e.g., $L \otimes \mathbf{Q}$) the group of indecomposables QR is I/I^2 where $I \subset R$ is the ideal of elements of positive degree.

A2.1.10. THEOREM (Lazard [1]). (a) $L = \mathbf{Z}[x_1, x_2, \dots]$ with $|x_i| = 2i$ for $i > 0$.

(b) x_i can be chosen so that its image in $QL \otimes \mathbf{Q}$ is

$$\begin{cases} pm_i & \text{if } i = p^k - 1 \text{ for some prime } p \\ m_i & \text{otherwise.} \end{cases} \quad \square$$

(c) L is a subring of M [A2.1.9(b)].

The proof of this is not easy and we will postpone the hardest part of it (A2.1.12) to the end of this section. The difficulty is in effect showing that L is torsion-free. Without proving A2.1.12 we can determine $L/\text{torsion}$ with relative ease. We will not give F in terms of the x_i , nor will the latter be given explicitly. Such formulas can be found, however, in Hazewinkel [3] and in Section 5 of Hazewinkel [1].

Before stating the hard lemma we need the following exercise in binomial coefficients.

A2.1.11. PROPOSITION. Let u_n be the greatest common divisor of the numbers $\binom{n}{i}$ for $0 < i < n$. Then

$$u_n = \begin{cases} p & \text{if } n = p^k \text{ for some prime } p \\ 1 & \text{otherwise.} \end{cases} \quad \square$$

Now we are ready for the hard lemma. Define homogeneous symmetric polynomials $B_n(x, y)$ and $C_n(x, y)$ of degree n for all $n > 0$ by

$$B_n(x, y) = (x + y)^n - x^n - y^n$$

$$C_n(x, y) = \begin{cases} B_n/p & \text{if } n = p^k \text{ for some prime } p \\ B_n & \text{otherwise.} \end{cases}$$

It follows from A2.1.11 that $C_n(x, y)$ is integral and that it is not divisible by any integer greater than one.

A2.1.12. COMPARISON LEMMA (Lazard [1]). *Let F and G be two formal group laws over R such that $F \equiv G \pmod{(x, y)^n}$. Then $F \equiv G + aC_n \pmod{(x, y)^{n+1}}$ for some $a \in R$. \square*

The proof for general R will be given at the end of this section. For now we give a proof for torsion-free R .

In this case we lose no information by passing to $R \otimes \mathbf{Q}$, where we know (A2.1.6) that both formal group laws have logarithms, say $f(x)$ and $g(x)$, respectively. Computing $\pmod{(x, y)^{n+1}}$ we have

$$f(x) \equiv g(x) + bx^n \quad \text{for some } b \in R \otimes \mathbf{Q} \quad \text{so} \quad f^{-1}(x) = g^{-1}(x) - bx^n$$

and

$$\begin{aligned} F - G &= f^{-1}(f(x) + f(y)) - g^{-1}(g(x) + g(y)) \\ &\equiv g^{-1}(g(x) + g(y) + b(x^n + y^n)) - b(x + y)^n - g^{-1}(g(x) + g(y)) \\ &\equiv g^{-1}(g(x) + g(y)) + b(x^n + y^n) - b(x + y)^n - g^{-1}(g(x) + g(y)) \\ &\equiv -bB_n(x, y). \end{aligned}$$

Since this must lie in R it must have the form $aC_n(x, y)$, completing the proof for torsion-free R .

A2.1.13. LEMMA. (a) *In $QL \otimes \mathbf{Q}$, $a_{i,j} = -\binom{i+j}{j}m_{i+j-1}$.*

(b) *QL is torsion-free.*

PROOF. (a) Over $L \otimes \mathbf{Q}$ we have $\sum m_{n-1}(\sum a_{i,j}x^i y^j)^n = \sum m_{n-1}(x^n + y^n)$. Using A2.1.3 to pass to $QL \otimes \mathbf{Q}$ we get

$$\sum a_{i,j}x^i y^j + \sum_{n>1} m_{n-1}(x + y)^n = \sum_{n>0} m_{n-1}(x^n + y^n),$$

which gives the desired formula.

(b) Let $Q_{2n}L$ denote the component of QL in degree $2n$, and let R be the graded ring $\mathbf{Z} \oplus Q_{2n}L$. Let F be the formal group law over R induced by the obvious map $\theta: L \rightarrow R$, and let G be the additive formal group law over R . Then by A2.1.12, $F(x, y) \equiv x + y + aC_{n+1}(x, y)$ for $a \in Q_{2n}L$. It follows that $Q_{2n}L$ is a cyclic group generated by a . By (a) $Q_{2n}L \otimes \mathbf{Q} = \mathbf{Q}$, so $\mathbf{Q}_{2n}L = \mathbf{Z}$ and QL is torsion-free. \square

It follows from the above that L is generated by elements x_i whose images in $QL \otimes \mathbf{Q}$ are $u_i m_i$, where u_i is as in A2.1.11, i.e., that L is a quotient of $\mathbf{Z}[x_i]$. By A2.1.9 it is the quotient by the trivial ideal, so A2.1.10 is proved.

Note that having A2.1.12 for torsion-free R implies that $L/\text{torsion}$ is as claimed.

The reader familiar with Quillen's theorem (4.1.6) will recognize L as $\pi_*(MU) = MU_*$. We will now define an object which is canonically isomorphic to $\pi_*(MU \wedge MU) = MU_*(MU)$. This description of the latter is due to Landweber [1].

A2.1.14. DEFINITION. *Let R be a commutative ring with unit. Then $FGL(R)$ is the set of formal group laws over R (A2.1.1) and $SI(R)$ is the set of triples (F, f, G) where $F, G \in FGL(R)$ and $f: F \rightarrow G$ is a strict isomorphism (A2.1.5), i.e., $f(x) \in R[[x]]$ with $f(0) = 0$, $f'(0) = 1$, and $f(F(x, y)) = G(f(x), f(y))$. We call such a triple a matched pair*

A2.1.15. PROPOSITION. *$FGL(-)$ and $SI(-)$ are covariant functors on the category of commutative rings with unit. $FGL(-)$ is represented by the Lazard ring L and $SI(-)$ is represented by the ring $LB = L \otimes \mathbf{Z}[b_1, b_2, \dots]$. In the grading introduced above, $|b_i| = 2i$.*

PROOF. All but the last statement are obvious. Note that a matched pair (F, f, G) is determined by F and f and that f can be any power series of the form $f(x) = x + \sum_{i>0} f_i x^{i+1}$. Hence such objects are in 1-1 correspondence with ring homomorphisms $\theta: LB \rightarrow R$ with $\theta(b_i) = f_i$. \square

Now LB has some additional structure which we wish to describe. Note that $FGL(R)$ and $SI(R)$ are the sets of objects and morphisms, respectively, of a groupoid, i.e., a small category in which every morphism is an equivalence. Hence these functors come equipped with certain natural transformations reflecting this structure. The most complicated is the one corresponding to composition of morphisms, which gives a natural (in R) map from a certain subset of $SI(R) \times SI(R)$ to $SI(R)$. This structure also endows (L, LB) with the structure of a Hopf algebroid (A1.1.1). Indeed that term was invented by Haynes Miller with this example in mind. We now describe this structure.

A2.1.16. THEOREM. *In the Hopf algebroid (L, LB) defined above $\varepsilon: LB \rightarrow L$ is defined by $\varepsilon(b_i) = 0$; $\eta_L: L \rightarrow LB$ is the standard inclusion while $\eta_R: L \otimes \mathbf{Q} \rightarrow LB \otimes \mathbf{Q}$ is given by*

$$\sum_{i \geq 0} \eta_R(m_i) = \sum_{i \geq 0} m_i \left(\sum_{j \geq 0} c(b_j) \right)^{i+1},$$

where $m_0 = b_0 = 1$; $\sum_{i \geq 0} \Delta(b_i) = \sum_{j \geq 0} (\sum_{i \geq 0} b_i)^{j+1} \otimes b_j$; and $c: LB \rightarrow LB$ is determined by $c(m_i) = \eta_R(m_i)$ and $\sum_{i \geq 0} c(b_i) \left(\sum_{j \geq 0} b_j \right)^{i+1} = 1$.

These are the structure formulas for $MU_*(MU)$ (4.1.11).

PROOF. ε and η_L are obvious. For c , if $f(x) = \sum b_i x^{i+1}$ then $f^{-1}(x) = \sum c(b_i) x^{i+1}$. Expanding $f^{-1}(f(1)) = 1$ gives the formula for $c(b_i)$. For η_R , let $\log x = \sum m_i x^{i+1}$ and $\text{mog } x = \sum \eta_R(m_i) x^{i+1}$ be the logarithms for F and G , respectively. Then we have

$$f^{-1}(G(x, y)) = F(f^{-1}(x), f^{-1}(y))$$

so

$$\log(f^{-1}(G(x, y))) = \log(f^{-1}(x)) + \log(f^{-1}(y)).$$

We also have

$$\text{mog}(G(x, y)) = \text{mog}(x) + \text{mog}(y)$$

for which we deduce

$$\text{mog}(x) = \log f^{-1}(x).$$

Setting $x = 1$ gives the formula for η_R . For Δ let $f_1(x) = b'_i x^{i+1}$, $f_2(x) = \sum b'_i x^{i+1}$, and $f(x) = f_2(f_1(x))$. Then expanding and setting $x = 1$ gives $\sum b_i = \sum b'_i (\sum b'_j)^{i+1}$. Since f_2 follows f_1 this gives the formula for Δ . \square

Note that (L, LB) is split (A1.1.22) since Δ defines a Hopf algebra structure on $B = \mathbf{Z}[b_i]$.

Next we will show how the theory simplifies when we localize at a prime p , and this will lead us to BP_* and $BP_*(BP)$.

A2.1.17. DEFINITION. A formal group law over a torsion-free $\mathbf{Z}_{(p)}$ -algebra is p -typical if its logarithm has the form $\sum_{i \geq 0} \ell_i x^{p^i}$ with $\ell_0 = 1$.

Later (A2.1.22) we will give a form of this definition which works even when the $\mathbf{Z}_{(p)}$ -algebra R has torsion. Assuming this can be done, we have

A2.1.18. THEOREM (Cartier [1]). Every formal group law over a $\mathbf{Z}_{(p)}$ -algebra is canonically strictly isomorphic to a p -typical one.

Actually A2.1.17 is adequate for proving the theorem because it suffices to show that the universal formal group law is isomorphic over $L \otimes \mathbf{Z}_{(p)}$ to a p -typical one.

The following notation will be used repeatedly.

A2.1.19. DEFINITION. Let F be a formal group law over R . If x and y are elements in an R -algebra A which also contains the power series $F(x, y)$, let

$$x +_F y = F(x, y).$$

This notation may be iterated, e.g., $x +_F y +_F z = F(F(x, y), z)$. Similarly, $x -_F y = F(x, i(y))$ (A2.1.2). For nonnegative integers n , $[n]_F(x) = F(x, [n-1]_F(x))$ with $[0]_F(x) = 0$. (The subscript F will be omitted whenever possible.) $\sum^F(\)$ will denote the formal sum of the indicated elements.

A2.1.20. PROPOSITION. If the formal group law F above is defined over a K -algebra R where K is a subring of \mathbf{Q} , then for each $r \in K$ there is a unique power series $[r]_F(x)$ such that

- (a) if r is a nonnegative integer, $[r]_F(x)$ is the power series defined above,
- (b) $[r_1 + r_2]_F(x) = F([r_1]_F(x), [r_2]_F(x))$,
- (c) $[r_1 r_2]_F(x) = [r_1]_F([r_2]_F(x))$.

PROOF. Let $[-1]_F(x) = i(x)$ (A2.1.2), so $[r]_F(x)$ is defined by (b) for all $r \in \mathbf{Z}$. We have $[r]_F(x) \equiv rx \pmod{x^2}$, so if $d \in \mathbf{Z}$ is invertible in K , the power series $[d]_F(x)$ is invertible and we can define $[d^{-1}]_F(x) = [d]_F^{-1}(x)$. \square

Now we suppose q is a natural number which is invertible in R . Let

$$(A2.1.21) \quad f_q(x) = [1/q] \left(\sum_{i=1}^q {}^F \zeta^i x \right)$$

where ζ is a primitive q th root of unity. A priori this is a power series over $R[\zeta]$, but since it is symmetric in the ζ^i it is actually defined over R .

If R is torsion-free and $\log(x) = \sum_{i \geq 0} m_i x^{i+1}$, we have

$$\begin{aligned} \log(f_q(x)) &= \frac{1}{q} \sum_{i=1}^q \log(\zeta^i x) \\ &= \frac{1}{q} \sum_{i=1}^q \sum_{j \geq 0} m_j x^{j+1} \zeta^{i(j+1)} \\ &= \frac{1}{q} \sum_{j \geq 0} m_j x^{j+1} \sum_{i=1}^q \zeta^{i(j+1)}. \end{aligned}$$

The expression $\sum_{i=1}^q \zeta^{i(j+1)}$ vanishes unless $(j+1)$ is divisible by q , in which case its value is q . Hence, we have

$$\log(f_q(x)) = \sum_{j > 0} m_{qj-1} x^{qj}.$$

If F is p -typical for $p \neq q$, this expression vanishes, so we make

A2.1.22. DEFINITION. A formal group law F over a $\mathbf{Z}_{(p)}$ -algebra is p -typical if $f_q(x) = 0$ for all primes $q \neq p$.

Clearly this is equivalent to our earlier definition A2.1.17 for torsion-free R .

To prove Cartier's theorem (A2.1.18) we claim that it suffices to construct a strict isomorphism $f(x) = \sum f_i x^i \in L \otimes \mathbf{Z}_{(p)}[[x]]$ from the image of F over $L \otimes \mathbf{Z}_{(p)}$ to a p -typical formal group law F' . Then if G is a formal group law over a $\mathbf{Z}_{(p)}$ -algebra R induced by a homomorphism $\theta: L \otimes \mathbf{Z}_{(p)} \rightarrow R$, $g(x) = \sum \theta(f_i) x^i \in R[[x]]$ is a strict isomorphism from G to a p -typical formal group law G' .

Recall that if $\text{mog}(x)$ is the logarithm for F' then

$$\text{mog}(x) = \log(f^{-1}(x)).$$

We want to use the $f_q(x)$ for various primes $q \neq p$ to concoct an $f^{-1}(x)$ such that

$$\log(f^{-1}(x)) = \sum_{i \geq 0} m_{p^i-1} x^{p^i}.$$

It would not do to set

$$f^{-1}(x) = x -_F \sum_{q \neq p}^F f_q(x)$$

because if n is a product of two or more primes $\neq p$ then a negative multiple of $M_{n-1} x^n$ would appear in $\log f^{-1}(x)$. What we need is the Möbius function $\mu(n)$ defined on natural numbers n by

$$\mu(n) = \begin{cases} 0 & \text{if } n \text{ is divisible by a square} \\ (-1)^r & \text{if } n \text{ is the product of } r \text{ distinct primes.} \end{cases}$$

Note that $\mu(1) = 1$ and $\mu(q) = -1$ if q is prime. Then we define $f(x)$ by

$$(A2.1.23) \quad f^{-1}(x) = \sum_{p \nmid q}^F [\mu(q)]_F (f_q(x)).$$

[Note also that $f_1(x) = x$.] The sum is over all natural numbers q not divisible by p . This infinite formal sum is well defined because $f_q(x) \equiv 0 \pmod{(x^q)}$.

Now

$$\log(f^{-1}(x)) = \sum_{p \nmid q} \mu(q) \sum_{j>0} m_{qj-1} x^{qj} = \sum_{n>0} \left(\sum_{\substack{p \nmid q \\ q|n}} \mu(q) \right) m_{n-1} x^n.$$

It is elementary to verify that

$$\sum_{\substack{p \nmid q \\ q|n}} \mu(q) = \begin{cases} 1 & \text{if } n = p^k \\ 0 & \text{otherwise.} \end{cases}$$

It follows that F' has logarithm

$$(A2.1.24) \quad \text{mog}(x) = \sum_{i \geq 0} m_{p^i-1} x^{p^i},$$

so F' is p -typical. This completes the proof of A2.1.18.

Now we will construct the universal p -typical formal group law.

A2.1.25. THEOREM. *Let $V = \mathbf{Z}_{(p)}[v_1, v_2, \dots]$ with $|v_n| = 2(p^n - 1)$. Then there is a universal p -typical formal group law F defined over V ; i.e., for any p -typical formal group law G over a commutative $\mathbf{Z}_{(p)}$ -algebra R , there is a unique ring homomorphism $\theta: V \rightarrow R$ such that $G(x, y) = \theta(F(x, y))$. Moreover the homomorphism from $L \otimes \mathbf{Z}_{(p)}$ to V corresponding (A2.1.8) to this formal group law is surjective, i.e., V is isomorphic to a direct summand $L \otimes \mathbf{Z}_{(p)}$. \square*

We will give an explicit formula for the v_n 's in terms of the log coefficients m_{p^n-1} below (A2.2.2). In 4.1.12 it is shown that V is canonically isomorphic to $\pi_*(BP)$.

PROOF. Recall that the canonical isomorphism f above corresponds to an endomorphism ϕ of $L \otimes \mathbf{Z}_{(p)}$ given by

$$\phi(m_i) = \begin{cases} m_i & \text{if } i = p^k - 1 \\ 0 & \text{otherwise.} \end{cases}$$

This ϕ is idempotent, i.e., $\phi^2 = \phi$ and its image is a subring $V \subset L \otimes \mathbf{Z}_{(p)}$ over which the universal p -typical formal group law is defined. An argument similar to the proof of Lazard's theorem A2.1.9 shows that V has the indicated structure. \square

Now we will construct a ring VT canonically isomorphic to $BP_*(BP)$ and representing the set of p -typical matched pairs (F, f, G) (A2.1.14), i.e., matched pairs with F and G p -typical. The power series f must be chosen carefully to ensure that G is p -typical, and this choice depends on F . There is no such thing as a " p -typical power series," i.e., one that sends any p -typical F to a p -typical G . To characterize the appropriate f we have

A2.1.26. LEMMA. *Let F be a p -typical formal group law over a $\mathbf{Z}_{(p)}$ -algebra R . Let $f(x)$ be an isomorphism (A2.1.5) from F to a formal group law G . Then G is p -typical if*

$$f^{-1}(x) = \sum_{i \geq 0}^F t_i x^{p^i}$$

for $t_i \in R$ with t_0 a unit in R .

PROOF. For a prime number $\neq p$ let

$$h_q(x) = [q^{-1}]_G \left(\sum_{i=1}^q \zeta^i x \right)$$

where ζ is a primitive p th root of unity. By A2.1.22 we need to show that $h_q(x) = 0$ for all $q \neq p$ iff f is as specified. From the relation

$$G(x, y) = f(F(f^{-1}(x), f^{-1}(y)))$$

we deduce

$$f^{-1}(h_q(x)) = [q^{-1}]_F \left(\sum_{j=1}^q f^{-1}(\zeta^j x) \right).$$

Now for isomorphism $f(x)$ there are unique $c_i \in R$ such that

$$f^{-1}(x) = \sum F_{i>0} c_i x^i$$

with c_1 a unit in R . Hence we have

$$\begin{aligned} f^{-1}(h_q(x)) &= [q^{-1}]_F \left(\sum_{i,j}^F c_i \zeta^{ij} x^i \right) \\ &= [q^{-1}]_F \left(\sum_{q \nmid i}^F \sum_j^F \zeta^j c_i x^i \right) +_F [q^{-1}] +_F \left(\sum_i^F [q]_F (c_{qi} x^{qi}) \right) \\ &= \sum_{q \nmid i}^F f_q(c_i x^i) +_F \sum_i^F c_{qi} x^{qi} = \sum_{i>0} c_{qi} x^{qi}. \end{aligned}$$

This expression vanishes for all $q \neq p$ iff $c_{qi} = 0$ for all $i > 0$ and $q \neq p$, i.e., iff f is as specified. \square

It follows immediately that $VT = V \otimes \mathbf{Z}_{(p)}[t_1, t_2, \dots]$ as a ring since for a strict isomorphism $t_0 = 1$. The rings V and VT represent the sets of objects and morphisms in the groupoid of strict isomorphisms of p -typical formal group laws over a $\mathbf{Z}_{(p)}$ -algebra. Hence (V, VT) , like (L, LB) , is a Hopf algebroid (A1.1.1) and it is isomorphic to $(BP_*, BP_*(BP))$. Its structure is as follows.

A2.1.27. THEOREM. *In the Hopf algebroid (V, VT) (see A1.1.1)*

- (a) $V = \mathbf{Z}_{(p)}[v_1, v_2, \dots]$ with $|v_n| = 2(p^n - 1)$,
- (b) $VT = V \otimes \mathbf{Z}_{(p)}[t_1, t_2, \dots]$ with $|t_n| = 2(p^n - 1)$, and
- (c) $\eta_L: V \rightarrow VT$ is the standard inclusion and $\varepsilon: VT \rightarrow V$ is defined by $\varepsilon(t_i) = 0$, $\varepsilon(v_i) = v_i$.

Let $\ell_i \in V \otimes \mathbf{Q}$ denote the image of $m_{p^i-1} \in L \otimes \mathbf{Q}$ (see A2.1.9). Then

- (d) $\eta_R: V \rightarrow VT$ is determined by $\eta_R(\ell_n) = \sum_{0 \leq i \leq n} \ell_i t_{n-i}^{p^i}$ where $\ell_0 = t_0 = 1$,
- (e) Δ is determined by $\sum_{i,j \geq 0} \ell_i \Delta(t_j)^{p^i} = \sum_{i,k,j \geq 0} \ell_i t_j^{p^i} \otimes t_k^{p^{i+j}}$, and
- (f) c is determined by $\sum_{i,j,k \geq 0} \ell_i t_j^{p^i} c(t_k)^{p^{i+j}} = \sum_{i \geq 0} \ell_i$.
- (g) The forgetful functor from p -typical formal group laws to formal group laws induces a surjection of Hopf algebroids (A1.1.19) $(L \otimes \mathbf{Z}_{(p)}, LB \otimes \mathbf{Z}_{(p)}) \rightarrow (V, VT)$.

Note that (e) and (f) are equivalent to

$$\sum_{i \geq 0}^F \Delta(t_i) = \sum_{i,j \geq 0}^F t_i \otimes t_j^{p^i} \quad \text{and} \quad \sum_{i,j \geq 0}^F t_i c(t_j)^{p^i} = 1,$$

respectively.

It can be shown that unlike (L, LB) (A2.1.16), (V, VT) is not split (A1.1.22).

PROOF. Part (a) was proved in A2.1.23, (b) follows from A2.1.23, and (c) is obvious, as is (g).

For (d) let f be a strict isomorphism between p -typical formal group law F and G with logarithms $\log(x)$ and $\text{mog}(x)$, respectively. If $f(x)$ satisfies

$$f^{-1}(x) = \sum_{i \geq 0}^F t_i x^{p^i}$$

and

$$\log(x) = \sum_{i \geq 0} \ell_i x^{p^i}$$

then by definition of η_R

$$\text{mog}(x) = \sum_{i \geq 0} \eta_R(\ell_i) x^{p^i}.$$

We have (see the proof of A2.1.16)

$$\begin{aligned} \text{mog}(x) &= \log(f^{-1}(x)) = \log\left(\sum_{i \geq 0}^F t_i x^{p^i}\right) \\ &= \sum_{i \geq 0} \log(t_i x^{p^i}) = \sum_{i, j \geq 0} \ell_i t_j^{p^i} x^{p^{i+j}} \end{aligned}$$

and (d) follows.

For (e) let $F \xrightarrow{f_1} G \xrightarrow{f_2} H$ be strict isomorphisms of p -typical formal group laws with

$$f_1^{-1}(x) = \sum_{i \geq 0}^F t'_i x^{p^i} \quad \text{and} \quad f_2^{-1}(x) = \sum_{j \geq 0}^G t''_j x^{p^j}.$$

If we set $f = f_2 \circ f_1$, with

$$f^{-1}(x) = \sum_{i \geq 0}^F t_i x^{p^i}$$

then a formula for t_i in terms of t'_i and t''_i will translate to a formula for $\Delta(t_i)$.

We have

$$\begin{aligned} f^{-1}(x) &= f_1^{-1}(f_2^{-1}(x)) = f_1^{-1}\left(\sum_{j \geq 0}^G t''_j x^{p^j}\right) \\ &= \sum_j^F f_1^{-1}(t''_j x^{p^j}) = \sum_{i, j}^F t'_i (t''_j x^{p^j})^{p^i}. \end{aligned}$$

This gives

$$\sum_i^F \Delta(t_i) = \sum_{i, j}^F t_i \otimes t_j^{p^i}$$

as claimed.

For (f) let $f: F \rightarrow G$ be as above. Then

$$f(x) = \sum^G c(t_j) x^{p^j}$$

so

$$\begin{aligned} x &= f^{-1}(f(x)) = f^{-1}\left(\sum_j^G c(t_j)x^{p^j}\right) \\ &= \sum_j^F f^{-1}(c(t_j)x^{p^j}) = \sum_{i,j}^F t_i(c(t_j)x^{p^j})^{p^i} \end{aligned}$$

setting $x = 1$ gives (f). □

Our only remaining task is to prove Lazard's comparison lemma A2.1.12. The proof below is due to Fröhlich [1]. The lemma states that if F and G are formal group laws with $F \equiv G \pmod{(x, y)^n}$ then

$$F \equiv G + aC_n(x, y) \pmod{(x, y)^{n+1}},$$

where

$$C_n(x, y) = \begin{cases} \frac{(x+y)^n - x^n - y^n}{p} & \text{if } n = p^k \text{ for some prime } p \\ (x+y)^n - x^n - y^n & \text{otherwise.} \end{cases}$$

Let $\Gamma(x, y)$ be the degree n component of $F - G$.

A2.1.28. LEMMA. $\Gamma(x, y)$ above is a homogeneous polynomial satisfying

- (i) $\Gamma(x, y) = \Gamma(y, x)$,
- (ii) $\Gamma(x, 0) = \Gamma(0, x) = 0$,
- (iii) $\Gamma(x, y) + \Gamma(x + y, z) = \Gamma(x, y + z) + \Gamma(y, z)$.

PROOF. Parts (i) and (ii) follow immediately A2.1.1(ii) and (i), respectively. For (iii) let $G(x, y) = x + y + G'(x, y)$. Then mod $(x, y, z)^{n+1}$ we have

$$\begin{aligned} F(F(x, y), z) &\equiv G(F(x, y), z) + \Gamma(F(x, y), z) \\ &\equiv F(x, y) + z + G'(F(x, y), z) + \Gamma(x + y, z) \\ &\equiv G(x, y) + \Gamma(x, y) + z + G'(G(x, y), z) + \Gamma(x + y, z) \\ &\equiv G(G(x, y), z) + \Gamma(x, y) + \Gamma(x + y, z). \end{aligned}$$

Similarly,

$$F(x, F(y, z)) = G(x, G(y, z)) + \Gamma(x, y + z) + \Gamma(y, z)$$

from which (iii) follows. □

It suffices to show that any such Γ must be a multiple of C_n .

A2.1.29. LEMMA. Let R be a field of characteristic $p > 0$. Then any $\Gamma(x, y)$ over R as above is a multiple of $C_n(x, y)$.

PROOF. It is easy to verify that C_n satisfies the conditions of A2.1.28, so it suffices to show that the set of all such Γ is one-dimensional vector space. Let $\Gamma(x, y) = \sum a_i x^i y^{n-i}$. Then from A2.1.28 we have

$$a_0 = a_n = 0, \quad a_i = a_{n-i},$$

and

$$(A2.1.30) \quad a_i \binom{n-i}{j} = a_{i+j} \binom{i+j}{j} \quad \text{for } 0 < i, i+j < n.$$

The case $n = 1$ is trivial so we write $n = sp^k$ with either $s = p$ or $s > 1$ and $s \not\equiv 0 \pmod p$. We will prove the lemma by showing $a_i = 0$ if $i \not\equiv 0 \pmod{p^k}$ and that a_{cp^k} is a fixed multiple of a_{p^k} .

If $i \not\equiv 0 \pmod{p^k}$ we can assume by symmetry that $i < (s - 1)p^k$ and write $i = cp^k - j$ with $0 < c < s$ and $0 < j < p^k$. Then A2.1.30 gives

$$a_i \binom{(s - c)p^k + j}{j} = a_{cp^k} \binom{cp^k}{j},$$

i.e., $a_i = 0$.

To show a_{cp^k} is determined by a_{p^k} for $c < s$ let $i = p^k$ and $j = (c - 1)p^k$. Then A2.1.30 gives

$$a_{p^k} \binom{(s - 1)p^k}{(c - 1)p^k} = a_{cp^k} \binom{cp^k}{(c - 1)p^k},$$

i.e.,

$$a_{p^k} \binom{s - 1}{c - 1} = a_{cp^k} c.$$

This determines a_{cp^k} provided $c \not\equiv 0 \pmod p$. Since $c < s$ we are done for the case $s = p$. Otherwise $a_{cp^k} = a_{(s-c)p^k}$ by symmetry and since $s \not\equiv 0 \pmod p$ either c or $s - c$ is $\not\equiv 0 \pmod p$. \square

Note that A2.1.29 is also true for fields of characteristic 0; this can be deduced immediately from A2.1.30. Alternatively, we have already proved A2.1.12, which is equivalent to A2.1.29, for torsion-free rings.

The proof of A2.1.29 is the last hard computation we have to do. Now we will prove the analogous statement for $R = \mathbf{Z}/(p^m)$ by induction on m . We have

$$\Gamma(x, y) = aC_n(x, y) + p^{m-1}\Gamma'(x, y),$$

where Γ' satisfies A2.1.28 mod p . Hence by A2.1.29 $\Gamma'(x, y) = bC_n(x, y)$ so

$$\Gamma(x, y) = (a + bp^{m-1})C_n(x, y)$$

as claimed.

To prove A2.1.29 (and hence A2.1.12) for general R note that the key ingredient A2.1.30 involves only the additive structure of R ; i.e., we only have to compute in a finitely generated abelian group A containing the coefficient of Γ . We have to show that symmetry and A2.1.30 imply that the coefficients a_i are fixed in relation to each other as are the coefficients of C_n . We have shown that this is true for $A = \mathbf{Z}$ (from the case $R = \mathbf{Q}$) and $A = \mathbf{Z}/(p^m)$. It is clear that if it is true for groups A_1 and A_2 then it is true for $A_1 \oplus A_2$, so it is true for all finitely generated abelian groups A . This completes the proof of A2.1.12.

2. Classification and Endomorphism Rings

In order to proceed further we need an explicit choice of the generators v_n . The first such choice was given by Hazewinkel [2], which was circulating in preprint form six years before it was published. The same generators for $p = 2$ were defined earlier still by Liulevicius [3]. A second choice, which we will use, was given by Araki [1].

Hazewinkel's generators are defined by

$$(A2.2.1) \quad p\ell_n = \sum_{0 \leq i < n} \ell_i v_{n-i}^p$$

which gives, for example,

$$\begin{aligned}\ell_1 &= \frac{v_1}{p}, & \ell_2 &= \frac{v_2}{p} + \frac{v_1^{1+p}}{p^2}, \\ \ell_3 &= \frac{v_3}{p} + \frac{v_1 v_2^p + v_2 v_1^{p^2}}{p^2} + \frac{v_1^{1+p+p^2}}{p^3}.\end{aligned}$$

Of course, it is nontrivial to prove that these v_n are contained in and generate V . Araki's formula is nearly identical,

$$(A2.2.2) \quad p\ell_n = \sum_{0 \leq i \leq n} \ell_i v_{n-i}^{p^i}$$

where $v_0 = p$. These v_n can be shown to agree with Hazewinkel's mod (p) . They give messier formulas for ℓ_n , e.g.,

$$\begin{aligned}\ell_1 &= \frac{v_1}{p - p^p}, & (p - p^{p^2})\ell_2 &= v_2 + \frac{v_1^{1+p}}{p - p^p}, \\ (p - p^{p^3})\ell_3 &= v_3 + \frac{v_1 v_2^p}{p - p^p} + \frac{v_2 v_1^{p^2}}{p - p^{p^2}} + \frac{v_1^{1+p+p^2}}{(p - p^p)(p - p^{p^2})},\end{aligned}$$

but a nicer formula (A2.2.5) for η_R .

A2.2.3. THEOREM (Hazewinkel [2], Araki [1]). *The sets of elements defined by A2.2.1 and A2.2.2 are contained in and generate V as a ring, and they are congruent mod (p) .*

PROOF. We first show that Araki's elements generate V . Equation A2.2.2 yields

$$\sum_{i \geq 0} p\ell_i x^{p^i} = \sum_{i, j \geq 0} \ell_i v_j^{p^i} x^{p^{i+j}}.$$

Applying \exp (the inverse of \log) to both sides gives

$$(A2.2.4) \quad [p]_F(x) = \sum_{i \geq 0}^F v_i x^{p^i},$$

which proves the integrality of the v_n , i.e., that $v_n \in V$. To show that they generate V it suffices by A2.1.10 to show $v_n = pu_n \ell_n$ in $QV \otimes \mathbf{Q}$, where u_n is a unit in $\mathbf{Z}_{(p)}$. Reducing A2.2.2 modulo decomposables gives

$$p\ell_n = v_n + \ell_n p^{p^n}$$

so the result follows.

We now denote Hazewinkel's generators of A2.2.1 by w_i . Then A2.2.1 gives

$$p \log x - px = \sum_{i > 0} \log w_i x^{p^i}$$

or

$$px = p \log x - \sum_{i > 0} \log w_i x^{p^i}.$$

Exponentiating both sides gives

$$\begin{aligned} \exp px &= [p](x) -_F \sum_{i>0}^F w_i x^{p^i} \\ &= px +_F \sum_{i>0}^F v_i x^{p^i} -_F \sum_{i>0}^F w_i x^{p^i} \text{ by A2.2.4.} \end{aligned}$$

If we can show that $(\exp px)/p$ is integral then the above equation will give

$$\sum_{i>0}^F v_i x^{p^i} \equiv \sum_{i>0}^F w_i x^{p^i} \pmod{(p)}$$

and hence $v_i \equiv w_i \pmod{(p)}$ as desired.

To show that $(\exp px)/p$ is integral simply note that its formal inverse is $(\log px)/p = \sum \ell_i p^{i-1} x^{p^i}$, which is integral since $p^i \ell_i$ is. \square

From now on v_n will denote the Araki generator defined by A2.2.2 or equivalently by A2.2.4. The following formula for $\eta_R(v_n)$ first appeared in Ravenel [1], where it was stated mod (p) in terms of the Hazewinkel generators; see also Moriera [2].

A2.2.5. THEOREM. *The behavior of η_R on v_n is defined by*

$$\sum_{i,j \geq 0}^F t_i \eta_R(v_j)^{p^i} = \sum_{i,j \geq 0}^F v_i t_j^{p^i}.$$

PROOF. Applying η_R to A2.2.2 and reindexing we get by A2.1.27(d)

$$\sum p \ell_i t_j^{p^i} = \sum \ell_i t_j^{p^i} \eta_R(v_k)^{p^{i+j}}.$$

Substituting A2.2.2 on the left-hand side and reindexing gives

$$\sum \ell_i v_j^{p^i} t_k^{p^{i+j}} = \sum \ell_i t_j^{p^i} \eta_R(v_k)^{p^{i+j}}.$$

Applying the inverse of log to this gives the desired formula. \square

This formula will be used to prove the classification theorem A2.2.11 below. Computational corollaries of it are given in Section 4.3.

We now turn to the classification in characteristic p . We will see that formal group laws over a field are characterized up to isomorphism over the separable algebraic closure by an invariant called the height (A2.2.7). In order to define it we need

A2.2.6. LEMMA. *Let F be a formal group law over a commutative \mathbf{F}_p -algebra R and let $f(x)$ be a nontrivial endomorphism of F (A2.1.5). Then for some n , $f(x) = g(x^{p^n})$ with $g'(0) \neq 0$. In particular f has leading term ax^{p^n} .*

For our immediate purpose we only need the statement about the leading term, which is easier to prove. The additional strength of the lemma will be needed below (A2.2.19). The argument we use can be adapted to prove a similar statement about a homomorphism to another formal group law G .

PROOF. Suppose inductively we have shown that $f(x) = f_i(x^{p^i})$, this being trivial for $i = 0$, and suppose $f'_i(0) = 0$, as otherwise we are done. Define $F^{(i)}(x, y)$

$$F(x, y)^{p^i} = F^{(i)}(x^{p^i}, y^{p^i}).$$

It is straightforward to show that $F^{(i)}$ is also a formal group law. Then we have

$$\begin{aligned} f_i(F^{(i)}(x^{p^i}, y^{p^i})) &= f_i(F(x, y)^{p^i}) = f(F(x, y)) \\ &= F(f(x), f(y)) = F(f_i(x^{p^i}), f_i(y^{p^i})) \end{aligned}$$

so

$$f_i(F^{(i)}(x, y)) = F(f_i(x), f_i(y)).$$

Differentiating with respect to y and setting $y = 0$ we get

$$f'_i(F^{(i)}(x, 0))F_2^{(i)}(x, 0) = F_2(f_i(x), f_i(0))f'_i(0).$$

Since $f'_i(0) = 0$, $F_2^{(i)}(x, 0) \neq 0$, and $F^{(i)}(x, 0) = x$, this gives us

$$f'_i(x) = 0 \quad \text{so} \quad f_i(x) = f_{i+1}(x^p).$$

We repeat this process until we get an $f_n(x)$ with $f'_n(0) \neq 0$ and set $g = f_n$. \square

A2.2.7. DEFINITION. A formal group law F over a commutative \mathbf{F}_p -algebra R has height n if $[p]_F(x)$ has leading term ax^{p^n} . If $[p]_F(x) = 0$ then F has height ∞ .

A2.2.8. LEMMA. The height of a formal group law is an isomorphism invariant.

PROOF. Let f be an isomorphism from F to G . Then

$$f([p]_F(x)) = [p]_G(f(x));$$

since $f(x)$ has leading term ux for u a unit in R and the result follows. \square

A2.2.9. EXAMPLES. Just for fun we will compute the heights of the mod (p) reductions of the formal group laws in A2.1.4.

(a) $[p]_F(x) = 0$ for all p so F has height ∞ .

(b) $[p]_F(x) = u_{p-1}x^p$ so F has height 1.

(c) As remarked earlier, F is isomorphic over $\mathbf{Z}_{(2)}$ to the additive formal group law, so its height at $p = 2$ is ∞ . Its logarithm is

$$\sum_{i \geq 0} \frac{x^{2i+1}}{2i+1}$$

so for each odd prime p we have $\ell_1 = m_{p-1} = 1/p$, so $v_1 \neq 0 \pmod p$ by A2.2.2, so the height is 1 by A2.2.4 and A2.2.7.

(d) Since F is not defined over $\mathbf{Z}_{(2)}$ (as can be seen by expanding it through degree 5) it does not have a mod 2 reduction. To compute its logarithm we have

$$F_2(x, 0) = \sqrt{1 - x^4}$$

so by A2.1.6

$$\begin{aligned} \log(x) &= \int_0^x \frac{dt}{\sqrt{1-t^4}} \\ &= \sum_{i \geq 0} \binom{-1/2}{i} \frac{(-1)^i x^{4i+1}}{4i+1} \\ &= \sum_{i \geq 0} \binom{(2i-1)/2}{i} \frac{x^{4i+1}}{4i+1} \\ &= \sum_{i \geq 0} \frac{1 \cdot 3 \cdot 5 \cdots (2i-1) x^{4i+1}}{2^i i! (4i+1)} \\ &= \sum_{i \geq 0} \frac{(2i)! x^{4i+1}}{2^{2i} (i!)^2 (4i+1)}. \end{aligned}$$

Now if $p \equiv 1 \pmod{4}$, we find that $\ell_1 = m_{p-1}$ is a unit (in $\mathbf{Z}_{(p)}$) multiple of $1/p$, so as in (c) the height is 1. However, if $p \equiv -1 \pmod{4}$, $v_1 = \ell_1 = 0$ so so the height is at least 2. We have

$$\ell_2 = m_{p^2-1} = \frac{(2i)!}{4(i!)^2 p^2} \quad \text{where } i = \frac{p^2-1}{4}.$$

Since

$$\frac{p^2-1}{2} = \frac{p(p-1)}{2} + \frac{p-1}{2},$$

$(2i)!$ is a unit multiple of $p^{(p-1)/2}$; since

$$\frac{p^2-1}{4} = p \binom{p-3}{4} + \frac{3p-1}{4}$$

$(i)!$ is a unit multiple of $p^{(p-3)/4}$. It follows that ℓ_2 is a unit multiple of $1/p$, so $v_2 \not\equiv 0 \pmod{p}$ and the height is 2.

It is known that the formal group law attached to a nonsingular elliptic curve always has height 1 or 2. (See Corollary 7.5 of Silverman [1]).

Now we will specify a formal group law of height n for each n .

A2.2.10. DEFINITION. $F_\infty(x, y) = x + y$. For a natural number h let F_n be the p -typical formal group law (of height n) induced by the homomorphism $\theta: V \rightarrow R$ (A2.1.25) defined by $\theta(v_n) = 1$ and $\theta(v_i) = 0$ for $i \neq n$.

A2.2.11. THEOREM (Lazard [2]). Let K be a separably closed field of characteristic $p > 0$. A formal group law G over K of height n is isomorphic to F_n .

PROOF. By Cartier's theorem (A2.1.18) we can assume G is p -typical (A2.1.22) and hence induced by a homomorphism $\theta: V \rightarrow K$ (A2.1.24). If $n = \infty$ then by A2.2.4 $\theta(v_n) = 0$ for all n and $G = F_\infty$. For n finite we have $\theta(v_i) = 0$ for $i < n$ and $\theta(v_n) \neq 0$. Let $F = F_n$. We want to construct an isomorphism $f: F \rightarrow G$ with $f^{-1}(x) = \sum_{i \geq 0}^F t_i x^{p^i}$. It follows from A2.2.5 that these t_i must satisfy

$$(A2.2.12) \quad \sum_{i,j}^F t_i \theta(v_j)^{p^i} x^{p^{i+j}} = \sum_j^F t_j^{p^n} x^{p^{n+j}}$$

since the homomorphism from V inducing F is given in A2.2.10, and the $\eta_R(v_j)$ in A2.2.5 correspond to $\theta(v_j)$. Here we are not assuming $t_0 = 1$; the proof of A2.2.5 is still valid if $t_0 \neq 1$.

Equating the coefficient of x^{p^n} in A2.2.12, we get $t_0\theta(v_n) = t_0^{p^n}$, which we can solve for t_0 since K is separably closed. Now assume inductively that we have solved A2.2.12 for t_0, t_1, \dots, t_{i-1} . Then equating coefficients of $x^{p^{i+n}}$ gives

$$t_i\theta(v_n)^{p^i} + c = t_i^{p^n}$$

for some $c \in K$. This can also be solved for t_i , completing the proof. \square

Our last objective in this section is to describe the endomorphism rings of the formal group laws F_n of A2.2.10.

A2.2.13. LEMMA. *Let F be a formal group law over a field K of characteristic $p > 0$ and let E be the set of endomorphisms of F .*

(a) *E is a ring under composition and formal sum, i.e., the sum of two endomorphisms $f(x)$ and $g(x)$ is $f(x) +_F g(x)$.*

(b) *E is a domain.*

(c) *E is a \mathbf{Z}_p -algebra (where \mathbf{Z}_p denotes the p -adic integers) which is a free \mathbf{Z}_p -module if F has finite height, and an \mathbf{F}_p -vector space if F has infinite height.*

PROOF.

(a) We need to verify the distributive law for these two operations. Let $f(x)$, $g(x)$, and $h(x)$ be endomorphisms. Then

$$f(g(x) +_F f(x)) = f(g(x)) +_F f(h(x))$$

so

$$f(g + h) = (fg) + (fh) \quad \text{in } E.$$

Similarly,

$$(g +_F h)(f(x)) = g(f(x)) +_F h(f(x))$$

so

$$(g + h)f = (gf) + (hf) \quad \text{in } E.$$

(b) Suppose $f(x)$ and $g(x)$ having leading terms ax^{p^n} and bx^{p^m} , respectively, with $a, b \neq 0$ (A2.2.6). Then $f(g(x))$ has leading term $ab^{p^m}x^{p^{m+n}}$, so $fg \neq 0$ in E .

(c) We need to show that $[a]_F(x)$ is defined for $a \in \mathbf{Z}_p$. We can write $a = \sum a_i p^i$ with $a_i \in \mathbf{Z}$. Then we can define

$$[a]_F(x) = \sum^F [a_i]_F([p^i]_F(x))$$

because the infinite formal sum on the right is in $K[[x]]$ since $[p^i]_F(x) \equiv 0$ modulo x^{p^i} . If $h < \infty$ then $[a]_F(x) \neq 0$ for all $0 \neq a \in \mathbf{Z}_p$, so E is torsion-free by (b). If $h = \infty$ then $[p]_F(x) = 0$ so E is an \mathbf{F}_p -vector space. \square

Before describing our endomorphism rings we need to recall some algebra.

A2.2.14. LEMMA. Let p be a prime and $q = p^i$ for some $i > 0$.

(a) There is a unique field \mathbf{F}_p with q elements.

(b) Each $x \in \mathbf{F}_q$ satisfies $x^q - x = 0$.

(c) \mathbf{F}_{p^m} is a subfield of \mathbf{F}_{p^n} iff $m \mid n$. The extension is Galois with Galois group $Z/(m/n)$ generated by the Frobenius automorphism $x \mapsto x^{p^m}$.

(d) $\overline{\mathbf{F}}_p$ the algebraic closure of \mathbf{F}_p and of each \mathbf{F}_q , is the union of all the \mathbf{F}_q . Its Galois group is $\widehat{\mathbf{Z}} = \varprojlim \mathbf{Z}/(m)$, the profinite integers, generated topologically by the Frobenius automorphism $x \mapsto x^p$. The subgroup $m\mathbf{Z}$ of index m is generated topologically by $x \mapsto x^{p^m}$ and fixes the field \mathbf{F}_{p^m} . \square

A proof can be found, for example, in Lang [1, Section VII.5]

Now we need to consider the Witt rings $W(\mathbf{F}_q)$, which can be obtained as follows. Over \mathbf{F}_p the polynomial $x^q - x$ is the product of irreducible factors of degrees at most n (where $q = p^n$) since it splits over \mathbf{F}_q , which is a degree n extension of \mathbf{F}_p . Let $h(x) \in \mathbf{Z}_p[x]$ be a lifting of an irreducible factor of degree n of $x^q - x$. Then let $W(\mathbf{F}_q) = \mathbf{Z}_p[x]/(h(x))$. It is known to be independent of the choices made and to have the following properties.

A2.2.15. LEMMA. (a) $W(\mathbf{F}_q)$ is a \mathbf{Z}_p -algebra and a free \mathbf{Z}_p -module of rank n , where $q = p^n$ [e.g., $W(\mathbf{F}_p) = \mathbf{Z}_p$].

(b) $W(\mathbf{F}_q)$ is a complete local ring with maximal ideal (p) and residue field \mathbf{F}_q .

(c) Each $w \in W(\mathbf{F}_q)$ can be written uniquely as $w = \sum_{i \geq 0} w_i p^i$ with $w_i^q - w_i = 0$ for each i .

(d) The Frobenius automorphism of \mathbf{F}_q lifts to an automorphism σ of $W(\mathbf{F}_q)$ defined by

$$w^\sigma = \sum_{i \geq 0} w_i^p p^i.$$

σ generates the Galois group $\mathbf{Z}/(n)$ of $W(\mathbf{F}_q)$ over \mathbf{Z}_p .

(e) $W(\mathbf{F}_q) = \varprojlim W(\mathbf{F}_q)/(p^i)$, so it is a compact topological ring.

(f) The group of units $W(\mathbf{F}_q)^\times$ is isomorphic to $W(\mathbf{F}_q) \oplus \mathbf{F}_q^\times$, where $\mathbf{F}_q^\times \cong \mathbf{Z}/(q-1)$, for $p > 2$, and to $W(\mathbf{F}_q) \oplus \mathbf{F}_q^\times \oplus \mathbf{Z}/(2)$ for $p = 2$, the extra summand being generated by -1 .

(g) $W(\mathbf{F}_q) \otimes \mathbf{Q} = \mathbf{Q}_p[x]/(h(x))$, the unramified degree n extension of \mathbf{Q}_p , the field of p -adic numbers.

A proof can be found in Mumford [1, Lecture 26] and in Serre [1, Section 11.5.6]. We will sketch the proof of (f). For $p > 2$ there is a short exact sequence

$$1 \rightarrow W(\mathbf{F}_q) \xrightarrow{i} W(\mathbf{F}_q)^\times \xrightarrow{j} \mathbf{F}_q^\times \rightarrow 1$$

where j is mod (p) reduction and $i(w) = \exp pw = \sum_{i \geq 0} (pw)^i / i!$ [this power series converges in $W(\mathbf{F}_q)$]. To get a splitting $\mathbf{F}_q^\times \rightarrow W(\mathbf{F}_q)^\times$ we need to produce $(q-1)$ th roots of unity in $W(\mathbf{F}_q)$, i.e., roots of the equation $x^q - x = 0$. [This construction is also relevant to (c).]

These roots can be produced by a device known as the Teichmüller construction. Choose a lifting u of a given element in \mathbf{F}_q , and consider the sequence $\{u, u^q, u^{q^2}, \dots\}$. It can be shown that it converges to a root of $x^q - x = 0$ which is independent of the choice of u .

For $p = 2$ the power series $\exp 2w$ need not converge, so we consider instead the short exact sequence

$$1 \rightarrow W(\mathbf{F}_q) \xrightarrow{i} W(\mathbf{F}_q)^\times \xrightarrow{j} W(\mathbf{F}_q)/(4)^\times \rightarrow 1,$$

where j is reduction mod (4) and $i(w) = \exp 4w$, which always converges. This sequence does not split. We have $W(\mathbf{F}_q)/(4)^\times \cong \mathbf{F}_q \oplus \mathbf{F}_q^\times$. Since $W(\mathbf{F}_q) \otimes \mathbf{Q}$ is a field, $W(\mathbf{F}_q)^\times$ can have no elements of order 2 other than ± 1 , so the other elements of order 2 in $W(\mathbf{F}_q)/(4)^\times$ lift to elements in $W(\mathbf{F}_q)^\times$ with nontrivial squares.

Next we describe the noncommutative \mathbf{Z}_p -algebra E_n , which we will show to be isomorphic to the endomorphism ring of F_n , for finite n .

A2.2.16. LEMMA. *Let E_n be the algebra obtained from $W(\mathbf{F}_q)$ by adjoining an indeterminate S and setting $S^n = p$ and $Sw = w^\sigma S$ for $w \in W(\mathbf{F}_q)$. Then*

- (a) E_n is a free \mathbf{Z}_p module of rank n^2 .
- (b) Each element $e \in E_n$ can be expressed uniquely as $\sum_{i \geq 0} e_i S^i$ with $e_i^q - e_i = 0$.
- (c) E_n is generated as a \mathbf{Z}_p -algebra by S and a primitive $(q-1)$ th root of unity ω with relations $S^n - p = 0$, $S\omega = \omega^p S = 0$, and $h(w) = 0$, where $h(x)$ is an irreducible degree n factor of $x^q - x$ over \mathbf{Z}_p .
- (d) E_n is the maximal order in $D_n = E_n \otimes \mathbf{Q}$ which is a division algebra with center \mathbf{Q}_p and invariant $1/n$.

The proofs of (a), (b), and (c) are elementary. To see that D_n is a division algebra, note that any element in D_n can be multiplied by some power of S to give an element in E_n which is nonzero mod (S) . It is elementary to show that such an element is invertible.

The invariant referred to in (d) is an element in \mathbf{Q}/\mathbf{Z} which classifies division algebras over \mathbf{Q}_p . Accounts of this theory are given in Serre [1, Chapters XII and XIII] Cassels and Fröhlich [1, pp. 137–139], Hazewinkel [1, Sections 20.2.16 and 23.1.4]. We remark that for $0 < i < n$ and i prime to n a division algebra with invariant i/n has a description similar to that of D_n except that S^n is p^i instead of p .

Our main results on endomorphism rings are as follows.

A2.2.17. THEOREM (Dieudonné [1] and Lubin [1]). *Let K be a field of characteristic p containing \mathbf{F}_q , with $q = p^n$. Then the endomorphism ring of the formal group law F_n (A2.2.10) over K is isomorphic to E_n . The generators ω and S [A2.2.16(c)] correspond to endomorphisms $\bar{\omega}x$ and x^p , respectively.*

A2.2.18. THEOREM. *Let R be a commutative \mathbf{F}_p -algebra. Then the endomorphism ring of the additive formal group law F_∞ over R is the noncommutative power series ring $R\langle\langle S \rangle\rangle$ in which $Sa = a^p S$ for $a \in R$. The elements a and S correspond to the endomorphisms ax and x^p , respectively.*

PROOF OF A2.2.18. An endomorphism $f(x)$ of F_∞ must satisfy $f(x+y) = f(x) + f(y)$. This is equivalent to $f(x) = \sum_{i \geq 0} a_i x^{p^i}$ for $a_i \in R$. The relation $Sa = a^p S$ corresponds to $(ax)^p = a^p x^p$. \square

There is an amusing connection between this endomorphism ring and the Steenrod algebra. Theorem A2.2.18 implies that the functor which assigns to each commutative \mathbf{F}_p -algebra R the strict automorphism group of the additive formal group

law is represented by the ring

$$P = \mathbf{F}_p[a_1, a_1, \dots]$$

since $a_0 = 1$ in this case. The group operation is represented by a coproduct $\Delta: P \rightarrow P \otimes P$. To compute Δa_n let $f_1(x) = \sum a'_j x^{p^k}$, $f_2(x) = \sum a''_k x^{p^k}$, and $f(x) = f_2(f_1(x)) = \sum a_i x^{p^i}$ with $a'_0 = a''_0 = a_0 = 1$. Then we have

$$f(x) = \sum a''_k \left(\sum a'_j x^{p^j} \right)^{p^k} = \sum a''_k (a'_j)^{p^j} x^{p^{j+k}}.$$

It follows that

$$\Delta a_n = \sum_{0 \leq i \leq n} a_{n-i}^{p^i} \otimes a_i \quad \text{with } a_0 = 1,$$

i.e., P is isomorphic to the dual of the algebra of Steenrod reduced powers.

Before proving A2.2.17 we need an improvement of A2.2.6.

A2.2.19. LEMMA. *Let F be a formal group law over a field K of characteristic $p > 0$, and let $f(x)$ be an endomorphism of F . Then*

$$f(x) = \sum_{i \geq 0}^F a_i x^{p^i}$$

for some $a_i \in K$.

PROOF. Suppose inductively we have $f(x) = \sum_{i=0}^{m-1} a_i x^{p^i} +_F f_m(x^{p^m})$, this being trivial for $m = 0$. Then set $a_m = f'_m(0)$ and consider the power series

$$g(x^{p^m}) = f_m(x^{p^m}) -_F a_m x^{p^m}.$$

By A2.2.13 this is an endomorphism and we have $g'_m(0) = 0$, so by A2.2.6 $g(x^{p^m}) = f_{m+1}(x^{p^{m+1}})$, completing the inductive step and the proof. \square

A2.2.17 will follow easily from the following.

A2.2.20. LEMMA. *Let $E(F_n)$ be the endomorphism ring of F_n (A2.2.10) over a field K containing \mathbf{F}_q where $q = p^n$. Then*

- (a) if $f(x) = \sum_{i=0}^{F_n} a_i x^{p^i}$ is in $E(F_n)$, then each $a_i \in \mathbf{F}_q$;
- (b) for $a \in \mathbf{F}_q$, $ax \in E(F_n)$;
- (c) $x^p \in E(F_n)$; and
- (d) $E(F_n)/(p) = E_n/(p) = \mathbf{F}_q \langle S \rangle / (S^n)$ with $Sa = a^p S$.

PROOF. (a) By the definition of F_n (A2.2.10) and A2.2.14 we have

$$(A2.2.21) \quad [p](x) = x^{p^n}.$$

Any endomorphism f commutes with $[p]$ so by A2.2.19 we have

$$[p](f(x)) = [p] \left(\sum_{i=0}^F a_i x^{p^i} \right) = \sum_{i=0}^F [p](a_i x^{p^i}) = \sum_{i=0}^F a_i^{p^n} x^{p^{i+n}}.$$

This must equal

$$f([p](x)) = \sum_{i=0}^F a_i ([p](x))^{p^i} = \sum_{i=0}^F a_i x^{p^{i+n}}.$$

Hence $a_i^{p^n} = a_i$ for all i and $a_i \in \mathbf{F}_q$.

(b) It suffices to prove this for $K = \mathbf{F}_q$. \mathbf{F}_n can be lifted to a formal group law \tilde{F}_n over $w(\mathbf{F}_q)$ (A2.2.15) by the obvious lifting of $\theta: V \rightarrow \mathbf{F}_q$, to $W(\mathbf{F}_q)$. It

suffices to show that ωx is an endomorphism of \tilde{F}_n if $\omega^q - \omega = 0$. By A2.2.2 \tilde{F}_n has a logarithm of the form

$$\log(x) = \sum a_i x^{q^i}$$

so $\log(\omega x) = \omega \log(x)$ and ωx is an endomorphism.

(c) This follows from the fact that F_n is defined over \mathbf{F}_p , so $F_n(x^p, y^p) = F_n(x, y)^p$.

(d) By A2.2.21, (b) and (c), $f(x) \in pE(F_n)$ iff $a_i = 0$ for $i < n$. It follows that for $f(x), g(x) \in E(F_n)$, $f \equiv g \pmod{(p)E(F_n)}$ iff $f(x) \equiv g(x) \pmod{(x^q)}$. Now our lifting \tilde{F}_n of F_n above has $\log x \equiv x \pmod{(x^q)}$, so $F_n(x, y) \equiv x + y \pmod{(x, y)^q}$. It follows that $E(F_n)/(p)$ is isomorphic to the corresponding quotient of $E(F_\infty)$ over \mathbf{F}_q , which is as claimed by A2.2.17. \square

PROOF OF A2.2.17. . By A2.2.16(c) E_n is generated by ω and S . The corresponding elements are in $E(F_n)$ by A2.2.20(b) and (c). The relation $S\omega = \omega^p S$ corresponds as before to the fact that $(\bar{\omega}x)^p = \bar{\omega}^p x^p$, where $\bar{\omega}$ is mod (p) reduction of ω . Hence we have a homomorphism $\lambda: E_n \rightarrow E(F_n)$ which is onto by A2.2.19. We know [A2.2.13(c)] that $E(F_n)$ is a free \mathbf{Z}_p -module. It has rank n^2 by A2.2.20(d), so λ is 1-1 by A2.2.16(a). \square

Bibliography

Adams, J. F.

- [1] *On the groups $J(x)$, IV*, *Topology* **5** (1966), 21–71.
- [2] *On the groups $J(x)$, II*, *Topology* **3** (1965), 137–171.
- [3] *On the structure and applications of the Steenrod algebra*, *Comm. Math. Helv.* **32** (1958), 180–214.
- [4] *Stable homotopy and generalised homology, Part III*, Univ. of Chicago Press, Illinois and London, 1974.
- [5] *Stable homotopy and generalised homology, Part II*, Univ. of Chicago Press, Illinois and London, 1974.
- [6] *Lectures on generalised cohomology*, *Lecture Notes in Math.*, vol. 99, Springer-Verlag, Berlin, 1969.
- [7] *Stable homotopy theory*, *Lecture Notes in Math.*, vol. 3, Springer-Verlag, Berlin, 1966.
- [8] *On Chern characters and the structure of the unitary group*, *Proc. Cambridge Philos. Soc.* **57** (1961), 189–199.
- [9] *Infinite loop spaces*, Princeton University Press, Princeton, N.J., 1978.

Adams, J. F. and Priddy, S. B.

- [10] *Uniqueness of BSO*, *Math. Proc. Cambridge Philos. Soc.* **80** (1978), 475–509.

Adams, J. F. and Margolis, H. R.

- [11] *Modules over the Steenrod algebra*, *Topology* **10** (1971), 271–282.

Adams, J. F.

- [12] *On the non-existence of element of Hopf invariant one*, *Ann. of Math.* **72** (1960), 20–104.

Adams, J. F. and Atiyah, M. F.

- [13] *K-theory and the Hopf invariant*, *Quart. J. Math. Oxford Ser. (2)* **17** (1966), 31–38.

Adams, J. F.

- [14] *Operations of the n th kind in K-theory, and what we don't know about $\mathbb{R}P^\infty$* , *New developments in topology. Proc. Sympos. Algebraic Topology (Oxford, 1972)*, London Math. Soc. Lecture Ser., vol. 11, Cambridge Univ. Press, London, 1974, pp. 1–9.
- [15] *The Kahn–Priddy theorem*, *Proc. Cambridge Philos. Soc.* **73** (1973), 45–55.
- [16] *Vector fields on spheres*, *Ann. of Math.* **75** (1962), 603–632.
- [17] *A periodicity theorem in homological algebra*, *Proc. Cambridge Philos. Soc.* **62** (1966), 365–377.

Adams, J. F., Gunawardena, J. H., and Miller, H. R.

- [18] *The Segal conjecture for elementary abelian p -groups*, *Topology* **24** (1985), no. 4, 435–460.

Aikawa, T.

- [1] *3-dimensional cohomology of the mod p Steenrod algebra*, *Math. Scand* **47** (1980), 91–115.

Anderson, D. W. and Davis, D. W.

- [1] *A vanishing theorem in homological algebra*, *Comm. Math. Helv.* **48** (1973), 318–327.

Anderson, D. W. and Hodckin, L.

- [2] *The K-theory of Eilenberg–Mac Lane complexes*, *Topology* **7** (1968), 317–329.

Ando, M., Hopkins, M. J., and Strickland, N. P.

- [1] *Elliptic spectra, the Witten genus and the theorem of the cube*, *Invent. Math.* **146** (2001), no. 3, 595–687.

Araki, S.

- [1] *Typical formal groups in complex cobordism and K-theory*, Kinokuniya Book-Store Co. Ltd., Tokyo, 1973.

Aubry, M.

- [1] *Calculs de groupes d'homotopie stables de la sphère, par la suite spectrale d'Adams–Novikov*, Math. Z. **185** (1984), no. 1, 45–91.

Baas, N. A.

- [1] *On bordism theory of manifolds with singularity*, Math. Scand. **33** (1973), 279–302.

Bahri, A. P. and Mahowald, M. E.

- [1] *A direct summand in $H^*(MO(8), \mathbf{Z}_2)$* , Proc. Amer. Math. Soc. **78** (1980), no. 2, 295–298.

Barratt, M. G., Mahowald, M. E., and Tangora, M. C.

- [1] *Some differentials in the Adams spectral sequence–II*, Topology **9** (1970), 309–316.

Barratt, M. G., Jones, J. D. S., and Mahowald, M. E.

- [2] *Relations amongst Toda brackets and Kervaire invariant in dimension 62*, J. London Math. Soc. **30** (1985), 533–550.

Behrens, M. and Pemmaraju, S.

- [1] *On the existence of the self map v_2^9 on the Smith-Toda complex $v(1)$ at the prime 3*, Online at <http://www.math.uchicago.edu/~mbehrens/>, to appear in the proceedings of the Northwestern University Algebraic Topology Conference, March 2002.

Bendersky, M., Curtis, E. B., and Miller, H. R.

- [1] *The unstable Adams sequence for generalized homology*, Topology **17** (1978), 229–248.

Bendersky, M.

- [2] *Some calculations in the unstable Adams–Novikov spectral sequence*, Publ. Res. Inst. Math. Sci., Kyoto University **16** (1980), no. 3, 739–766.

Bendersky, M., Curtis, E. B., and Ravenel, D. C.

- [3] *EHP sequence in BP theory*, Topology **21** (1982), 373–391.

Bott, R.

- [1] *The stable homotopy of the classical groups*, Ann. of Math. (2) **70** (1959), 313–337.
 [2] *The space of loops on a Lie group*, Michigan Math. J. **5** (1958), 35–61.

Bousfield, A. K. and Kan, D. M.

- [1] *Homotopy limits, completions and localizations*, Lecture Notes in Math., vol. 304, Springer-Verlag, Berlin, 1972.

Bousfield, A. K., Curtis, E. B., Kan, D. M., Quillen, D. G., Rector, D. L., and Schlesinger, J.

- [2] *The mod p lower central series and the Adams spectral sequence*, Topology **5** (1966), 331–342.

Bousfield, A. K. and Kan, D. M.

- [3] *The homotopy spectral sequence of a space with coefficients in a ring*, Topology **11** (1972), 79–100.

Bousfield, A. K. and Curtis, E. B.

- [4] *A spectral sequence for the homotopy of nice spaces*, Trans. Amer. Math. Soc. **151** (1970), 457–479.

Browder, W.

- [1] *The Kervaire invariant of framed manifolds and its generalization*, Ann. of Math. **90** (1969), 157–186.

Brown, E. H. and Peterson, F. P.

- [1] *A spectrum whose \mathbf{Z}_p cohomology is the algebra of reduced p -th powers*, Topology **5** (1966), 149–154.

Brown, E. H.

- [2] *Cohomology theories*, Ann. of Math. **75** (1962), 467–484.

Brown, E. H. and Gitler, S.

- [3] *A spectrum whose cohomology is a certain cyclic module over the steenrod algebra*, Topology **12** (1973), 283–295.

- Bruner, R. R., May, J. P., McClure, J. E., and Steinberger, M.
 [1] *H_∞ ring spectra and their applications*, Lecture Notes in Math., Springer-Verlag, Berlin, 1986.
- Bruner, R. R.
 [2] *A new differential in the Adams spectral sequence*, *Topology* **23** (1984), 271–276.
 [3] *Cohomology of modules over the mod 2 Steenrod algebra*, Online at <http://www.math.wayne.edu/rrb/cohom/>.
- Carlsson, G.
 [1] *Equivariant stable homotopy and Segal's Burnside ring conjecture*, *Ann. of Math.* **120** (1984), 189–224.
- Cartan, H. and Eilenberg, S.
 [1] *Homological algebra*, Princeton Univ. Press, Princeton, New Jersey, 1956.
- Cartier, P.
 [1] *Modules associés à un groupe formel commutatif. Courbes typiques*, *C. R. Acad. Sci. Paris Sér. A-B* **265** (1967), A129–A132.
- Cassels, J. W. S. and Fröhlich, A.
 [1] *Algebraic number theory*, Academic Press, New York, 1967.
- Cohen, F., Moore, J. C., and Neisendorfer, J.
 [1] *The double suspension and exponents of the homotopy groups of spheres*, *Ann. of Math.* **110** (1979), 549–565.
- Cohen, R. L. and Goerss, P.
 [2] *Secondary cohomology operations that detect homotopy classes*, *Topology* **23** (1984), 177–194.
- Cohen, R. L.
 [3] *Odd primary infinite families in stable homotopy theory*, *Mem. Amer. Math. Soc.* **30** (1981), no. 242, viii+92.
- Conner, P. E. and Floyd, E. E.
 [1] *The relation of cobordism to K-theories*, Lecture Notes in Math., vol. 28, Springer-Verlag, Berlin, 1966.
 [2] *Differentiable periodic maps*, Academic Press Inc., Publishers, New York, 1964.
- Conner, P. E. and Smith, L.
 [3] *On the complex bordism of finite complexes*, *Inst. Hautes Études Sci. Publ. Math.* (1969), no. 37, 117–221.
- Conner, P. E.
 [4] *Differentiable periodic maps*, Lecture Notes in Math., Springer, Berlin, 1979.
- Curtis, E. B.
 [1] *Simplicial homotopy theory*, *Adv. in Math.* **6** (1971), 107–209.
- Davis, D. M. and Mahowald, M. E.
 [1] v_1 - and v_2 -periodicity in stable homotopy theory, *Amer. J. Math.* **103** (1981), 615–659.
- Davis, D. M.
 [2] *An infinite family in the cohomology of the Steenrod algebra*, *J. Pure Appl. Algebra* **21** (1981), 145–150.
 [3] *On the cohomology of $MO\langle 8 \rangle$* , *Symposium on Algebraic Topology in honor of José Adem (Oaxtepec, 1981)*, *Amer. Math. Soc.*, Providence, R.I., 1982, pp. 91–104.
- Davis, D. M. and Mahowald, M. E.
 [4] *The nonrealizability of the quotient $A//A_2$ of the Steenrod algebra*, *Amer. J. Math.* **104** (1982), 1211–1216.
 [5] *Ext over the subalgebra A_2 of the Steenrod algebra for stunted projective spaces*, *Current trends in algebraic topology, Part 1 (London, Ont., 1981)*, *Amer. Math. Soc.*, Providence, RI, 1982, pp. 297–342.
- Davis, D. M.
 [6] *The splitting of $BO\langle 8 \rangle \wedge bo$ and $MO\langle 8 \rangle \wedge bo$* , *Trans. Amer. Math. Soc.* **276** (1983), 671–683.

Devinatz, E. S. and Hopkins, M. J.

- [1] *The action of the Morava stabilizer group on the Lubin-Tate moduli space of lifts*, Amer. J. Math. **117** (1995), no. 3, 669–710.

Devinatz, E. S., Hopkins, M. J., and Smith, J. H.

- [2] *Nilpotence and stable homotopy theory*, Annals of Mathematics **128** (1988), 207–242.

Dieudonné, J.

- [1] *Groupes de Lie et hyperalgèbres de Lie sur un corps de caractéristique $p > 0$. VII*, Math. Ann. **134** (1957), 114–133.

Dold, A. and Thom, R.

- [1] *Quasifaserungen und unendliche symmetrische Produkte*, Ann. of Math. (2) **67** (1958), 239–281.

Eckmann, B.

- [1] *Gruppentheoretischer Beweis des Satzes von Hurwitz–Radon über die Komposition quadratischer Formen*, Comment. Math. Helv. **15** (1942), 358–366.

Eilenberg, S. and Moore, J. C.

- [1] *Foundations of relative homological algebra*, Mem. Amer. Math. Soc. **55** (1965), 39.

Eilenberg, S. and MacLane, S.

- [2] *Relation between homology and homotopy groups of spaces*, Ann. of Math. (2) **46** (1945), 480–509.

Elmendorf, A. D., Kriz, I., Mandell, M. A., and May, J. P.

- [1] *Rings, modules, and algebras in stable homotopy theory*, Mathematical Surveys and Monographs, vol. 47, American Mathematical Society, Providence, RI, 1997, With an appendix by M. Cole.

Floyd, E. E.

- [1] *Actions of $(\mathbf{Z}_p)^k$ without stationary points*, Topology **10** (1971), 327–336.

Fröhlich, A.

- [1] *Formal groups*, Lecture Notes in Math., Springer-Verlag, Berlin, 1968.

Giambalvo, V. and Pengelley, D. J.

- [1] *The homology of M spin*, Math. Proc. Cambridge Philos. Soc. **95** (1984), 427–436.

Giambalvo, V.

- [2] *On $\langle 8 \rangle$ -cobordism*, Illinois J. Math. **15** (1971), 533–541.

Gorbounov, V. and Symonds, P.

- [1] *Toward the homotopy groups of the higher real K -theory EO_2* , Homotopy theory via algebraic geometry and group representations (Evanston, IL, 1997), Contemp. Math., vol. 220, Amer. Math. Soc., Providence, RI, 1998, pp. 103–115.

Gray, B. W.

- [1] *On the sphere of origin of infinite families in the homotopy groups of spheres*, Topology **8** (1969), 219–232.

Gunawardena, J. H. C.

- [1] *Segal's conjecture for cyclic groups of (odd) prime order*, J. T. Knight Prize Essay, Cambridge, 1979.

Harper, J. R. and Miller, H. R.

- [1] *On the double suspension homomorphism at odd primes*, Trans. Amer. Math. Soc. **273** (1982), 319–331.

Hatcher, A.

- [1] *Stable homotopy groups of spheres*, Online at <http://www.math.cornell.edu/~hatcher/stemfigs/stems.html>.

Hazewinkel, M.

- [1] *Formal groups and applications*, Academic Press, New York, 1978.
 [2] *Constructing formal groups. I. The local one dimensional case*, J. Pure Appl. Algebra **9** (1976/77), no. 2, 131–149.

- [3] *Constructing formal groups. II. The global one dimensional case*, J. Pure Appl. Algebra **9** (1976/77), no. 2, 151–161.
- [4] *Constructing formal groups. I. The local one dimensional case*, J. Pure Appl. Algebra **9** (1976/77), no. 2, 131–149.
- Henn, H.-W.
- [1] *Centralizers of elementary abelian p -subgroups and mod- p cohomology of profinite groups*, Duke Math. J. **91** (1998), no. 3, 561–585.
- Higgins, P. J.
- [1] *Notes on categories and groupoids*, Van Nostrand Reinhold Co., London, 1971, Van Nostrand Reinhold Mathematical Studies, No. 32.
- Hilton, P. J. and Stammbach, U.
- [1] *A course in homological algebra*, Springer-Verlag, New York, 1971, Graduate Texts in Mathematics, Vol. 4.
- Hirzebruch, F.
- [1] *Topological methods in algebraic geometry*, Third enlarged edition. New appendix and translation from the second German edition by R. L. E. Schwarzenberger, with an additional section by A. Borel. Die Grundlehren der Mathematischen Wissenschaften, Band 131, Springer-Verlag New York, Inc., New York, 1966.
- Hopf, H.
- [1] *Über die Abbildungen von Sphären auf Sphären niedrigerer Dimension*, Fund. Math. **25** (1935), 427–440.
- Hopkins, M. J., Kuhn, N. J., and Ravenel, D. C.
- [1] *Generalized group characters and complex oriented cohomology theories*, J. Amer. Math. Soc. **13** (2000), no. 3, 553–594 (electronic).
- Hopkins, M. J. and Mahowald, M. A.
- [2] *From elliptic curves to homotopy theory*, Preprint in Hopf archive at <http://hopf.math.purdue.edu/Hopkins-Mahowald/eo2homotopy>.
- Hopkins, M. J. and Smith, J. H.
- [3] *Nilpotence and stable homotopy theory. II*, Ann. of Math. (2) **148** (1998), no. 1, 1–49.
- Hurewicz, W.
- [1] *Beiträge zur Topologie der Deformationen*, Nederl. Akad. Wetensch. Proc. Ser. A **38** (1935), 112–119, 521–528.
- [2] *Beiträge zur Topologie der Deformationen*, Nederl. Akad. Wetensch. Proc. Ser. A **39** (1936), 117–126, 215–224.
- James, I. M.
- [1] *Reduced product spaces*, Ann. of Math. (2) **62** (1955), 170–197.
- [2] *Whitehead products and vector fields on spheres*, Proc. Cambridge Philos. Soc. **53** (1957), 817–820.
- [3] *Spaces associated with Stiefel manifolds*, Proc. London Math. Soc. (3) **9** (1959), 115–140.
- Johnson, D. C., Miller, H. R., Wilson, W. S., and Zahler, R. S.
- [1] *Boundary homomorphisms in the generalized Adams spectral sequence and the nontriviality of infinitely many γ_t in stable homotopy*, Conference on homotopy theory (Evanston, Ill., 1974), Soc. Mat. Mexicana, México, 1975, pp. 47–63.
- Johnson, D. C. and Wilson, W. S.
- [2] *Projective dimension and Brown–Peterson homotopy*, Topology **12** (1973), 327–353.
- [3] *BP-operations and Morava’s extraordinary K -theories*, Math. Z. **144** (1975), 55–75.
- Johnson, D. C. and Yosimura, Z.
- [4] *Torsion in Brown–Peterson homology of and Hurewicz homomorphisms*, Osaka. J. Math. **17** (1980), 117–136.
- Johnson, D. C. and Wilson, W. S.
- [5] *The Brown–Peterson homology of elementary p -groups*, Amer. J. Math. **107** (1985), 427–453.

Kahn, D. S.

- [1] Cup – i products and the Adams spectral sequence, *Topology* **9** (1970), 1–9.

Kahn, D. S. and Priddy, S. B.

- [2] *The transfer and stable homotopy*, *Math. Proc. Cambridge Philos. Soc.* **83** (1978), 103–111.

Kambe, T., Matsunaga, H., and Toda, H.

- [1] *A note on stunted lens space*, *J. Math. Kyoto Univ.* **5** (1966), 143–149.

Kochman, S. O.

- [1] *The symplectic cobordism ring. I*, *Mem. Amer. Math. Soc.* **24** (1980), no. 228, ix+206.
 [2] *The symplectic cobordism ring. II*, *Mem. Amer. Math. Soc.* **40** (1982), no. 271, vii+170.
 [3] *The symplectic cobordism ring III*, To appear.
 [4] *A chain functor for bordism*, *Trans. Amer. Math. Soc.* **239** (1978), 167–190.
 [5] *Uniqueness of Massey products on the stable homotopy of spheres*, *Can. J. Math.* **32** (1980), 576–589.

Kriz, I.

- [1] *Morava K -theory of classifying spaces: some calculations*, *Topology* **36** (1997), no. 6, 1247–1273.

Kriz, I. and Lee, K. P.

- [2] *Odd-degree elements in the Morava $K(n)$ cohomology of finite groups*, *Topology Appl.* **103** (2000), no. 3, 229–241.

Kuhn, N. J.

- [1] *A Kahn–Priddy sequence and a conjecture of G. W. Whitehead*, *Math. Proc. Cambridge Philos. Soc.* **92** (1982), 467–483.

Landweber, P. S.

- [1] *$BP_*(BP)$ and typical formal groups*, *Osaka J. Math.* **12** (1975), 357–363.
 [2] *Cobordism operations and Hopf algebras*, *Trans. Amer. Math. Soc.* **129** (1967), 94–110.
 [3] *Homological properties of comodules over $MU_*(MU)$ and $BP_*(BP)$* , *Amer. J. Math.* **98** (1976), 591–610.
 [4] *Annihilator ideals and primitive elements in complex cobordism*, *Illinois J. Math.* **17** (1973), 273–284.
 [5] *Assosiated prime ideals and Hopf algebras*, *J. Pure Appl. Algebra* **3** (1973), 43–58.
 [6] *Complex bordism of classifying spaces*, *Proc. Amer. Math. Soc.* **27** (1971), 175–179.
 [7] *Invariant regular ideals in Brown–Peterson homology*, *Duke Math. J.* **42** (1975), 499–505.

Landweber, P. S., Ravenel, D. C., and Stong, R. E.

- [8] *Periodic cohomology theories defined by elliptic curves*, *The Čech Centennial* (Providence, Rhode Island) (Cenkl, M. and Miller, H., eds.), *Contemporary Mathematics*, vol. 181, American Mathematical Society, 1995, pp. 317–338.

Lang, S.

- [1] *Algebra*, Addison-Wesley Publishing Co., Inc., Reading, Mass., 1965.

Lannes, J.

- [1] *Applications dont la source est un classifiant*, *Proceedings of the International Congress of Mathematicians*, Vol. 1, 2 (Zürich, 1994) (Basel), Birkhäuser, 1995, pp. 566–573.
 [2] *Sur les espaces fonctionnels dont la source est le classifiant d'un p -groupe abélien élémentaire*, *Inst. Hautes Études Sci. Publ. Math.* (1992), no. 75, 135–244, With an appendix by Michel Zisman.

Lannes, J. and Schwartz, L.

- [3] *À propos de conjectures de Serre et Sullivan*, *Invent. Math.* **83** (1986), no. 3, 593–603.

Lazard, M.

- [1] *Lois de groupes et analyseurs*, *Ann. Écoles Norm. Sup.* **72** (1955), 299–400.
 [2] *Sur les groupes de Lie formels à un paramètre*, *Bull. Soc. Math. France* **83** (1955), 251–274.
 [3] *La non-existence des groupes de Lie formels non abéliens à un paramètre*, *C. R. Acad. Sci. Paris* **239** (1954), 942–945.
 [4] *Groupes analytiques p -adiques*, *Inst. Hautes Études Sci. Publ. Math.* (1965), no. 26, 389–603.

Li, H. H. and Singer, W. M.

- [1] *Resolution of modules over the Steenrod algebra and the classical theory of invariants*, Math. Z. **181** (1982), 269–286.

Lin, W. H.

- [1] *On conjecture of Mahowald, Segal, and Sullivan*, Math. Proc. Cambridge Philos. Soc. **87** (1980), 449–458.

Lin, W. H., Davis, D. M., Mahowald, M. E., and Adams, J. F.

- [2] *Calculation of Lin's Ext groups*, Math. Proc. Cambridge Philos. Soc. **87** (1980), 459–469.

Lin, W. H.

- [3] *Algebraic Kahn–Priddy theorem*, Pacific J. Math. **96** (1981), 435–455.

Liulevicius, A. L.

- [1] *A proof of Thom's theorem*, Comment. Math. Helv. **37** (1962/1963), 121–131.

Liulevicius, A.

- [2] *The factorization of cyclic reduced powers by secondary cohomology operations*, Proc. Nat. Acad. Sci. U.S.A. **46** (1960), 978–981.
 [3] *On the algebra $BP_*(BP)$* , Symposium on Algebraic Topology (Battelle Seattle Res. Center, Seattle, Wash., 1971), Lecture Notes in Math, vol. 249, Springer, Berlin, 1971, pp. 47–53.

Lubin, J.

- [1] *One-parameter formal Lie groups over p -adic integer rings*, Ann. of Math. (2) **80** (1964), 464–484.
 [2] *Correction to: "One-parameter formal Lie groups over p -adic integer rings"*, Ann. of Math. (2) **84** (1966), 372.

Lubin, J. and Tate, J.

- [3] *Formal moduli for one parameter formal Lie groups*, Bull. Soc. Math. France **94** (1966), 49–60.

Mac Lane, S.

- [1] *Homology*, Springer-Verlag, Berlin and New York, 1963.

MacLane, S.

- [1] *Categories for the working mathematician*, Springer-Verlag, New York, 1971, Graduate Texts in Mathematics, Vol. 5.

Mahowald, M. E.

- [1] *The metastable homotopy of S^n* , no. 72, Mem. Amer. Math. Soc., Providence, R.I., 1967.

Mahowald, M.

- [2] *Description homotopy of the elements in the image of the J -homomorphism*, Manifolds—Tokyo 1973 (Proc. Internat. Conf., Tokyo, 1973) (Tokyo), Univ. Tokyo Press, 1975, pp. 255–263.
 [3] *On the double suspension homomorphism*, Trans. Amer. Math. Soc. **214** (1975), 169–178.

Mahowald, M. E.

- [4] *The image of J in the EHP sequence*, Ann. of Math. **116** (1982), 65–112.

Mahowald, M. and Milgram, R. J.

- [5] *Operations which detect Sq_4 in connective K -theory and their applications*, Quart. J. Math. Oxford Ser. (2) **27** (1976), no. 108, 415–432.

Mahowald, M. E.

- [6] *A new infinite family in ${}_2\pi_*^S$* , Topology **16** (1977), 249–256.
 [7] *Some remarks on the Kervaire invariant problem from the homotopy point of view*, Proc. Sympos. Pure Math. **22** (1971), 165–169.

Mahowald, M. E. and Tangora, M. C.

- [8] *On secondary operations which detect homotopy classes*, Bol. Soc. Mat. Mexicana **12** (1967), 71–75.
 [9] *Some differentials in the Adams spectral sequence*, Topology **6** (1967), 349–369.

Mahowald, M. E.

- [10] *Some homotopy classes generated by η_j* , Algebraic topology, Aarhus 1978 (Proc. Sympos., Univ. Aarhus, Aarhus, 1978), Lectures Notes in Math., Springer, Berlin, 1979, pp. 23–37.
- [11] *The construction of small ring spectra*, Geometric applications of homotopy theory (Proc. Conf., Evanston, Ill., 1977), II, Lectures Notes in Math., Springer, Berlin, 1978, pp. 234–239.
- [12] *The primary v_2 -periodic family*, Math. Z. **177** (1981), 381–393.
- [13] *The order of the image of the J -homomorphism*, Bull. Amer. Math. Soc. **76** (1970), 1310–1313.

Mahowald, M. E. and Tangora, M. C.

- [14] *An infinite subalgebra of $\text{Ext}_A(\mathbf{Z}_2, \mathbf{Z}_2)$* , Trans. Amer. Math. Soc. **132** (1968), 263–274.

Mahowald, M. E.

- [15] *bo resolutions*, Pacific J. Math. **192** (1981), 365–383.

Mäkinen, J.

- [1] *Boundary formulae for reduced powers in the Adams spectral sequence*, Ann. Acad. Sci. Fenn. Ser. A I (1973), no. 562, 42.

Margolis, H. R., Priddy, S. B., and Tangora, M. C.

- [1] *Another systematic phenomenon in the cohomology of the Steenrod algebra*, Topology **10** (1971), 43–46.

Massey, W. S.

- [1] *Algebraic topology: an introduction*, Springer-Verlag, New York, 1977.
- [2] *Exact couples in algebraic topology*, Ann. of Math. **56** (1952), 363–396.
- [3] *Some higher order cohomology operations*, Sympos. Internat. Alg. Top., Universidad Nacional Autónoma de México and UNESCO, 1958, pp. 145–154.

Maunder, C. R. F.

- [1] *On the differentials in the Adams spectral sequence for the stable homotopy groups of spheres. II*, Proc. Cambridge Philos. Soc. **61** (1965), 855–868.
- [2] *On the differentials in the Adams spectral sequence for the stable homotopy groups of spheres. I*, Proc. Cambridge Philos. Soc. **61** (1965), 53–60.

May, J. P.

- [1] *The cohomology of restricted Lie algebras and of Hopf algebras; application to the Steenrod algebra*, Princeton, Univ., 1964, Thesis.
- [2] *The cohomology of the restricted Lie algebras and of Hopf algebras*, J. Algebra **3** (1966), 123–146.
- [3] *Matrix Massey products*, J. Algebra **12** (1969), 533–568.
- [4] *The Steenrod algebra and its associated graded algebra*, University of Chicago preprint, 1974.
- [5] *A general algebraic approach to Steenrod operations*, The Steenrod Algebra and its Applications (Proc. Conf. to Celebrate N. E. Steenrod's Sixtieth Birthday, Battelle Memorial Inst., Columbus, Ohio, 1970), Lecture Notes in Mathematics, Vol. 168, Springer, Berlin, 1970, pp. 153–231.
- [6] 1982, Private communication.

Milgram, R. J.

- [1] *The Steenrod algebra and its dual for connective K -theory*, Reunion sobre teoria de homotopia, Northern Univ., Soc. Mat. Mexicana, México, 1974, pp. 127–158.
- [2] *Group representations and the Adams spectral sequence*, Pacific J. Math. **41** (1972), 157–182.

Miller, H. R., Ravenel, D. C., and Wilson, W. S.

- [1] *The periodic phenomena in the Adams–Novikov spectral sequence*, Ann. of Math. **106** (1977), 169–516.

Miller, H. R.

- [2] *Some algebraic aspects of the Adams–Novikov spectral sequence*, Princeton Univ., 1974, Thesis.

Miller, H. R. and Wilson, W. S.

- [3] *On the Novikov's Ext^1 modulo an invariant prime ideal*, Topology **15** (1976), 131–141.

- [4] *On relations between Adams spectral sequences, with an application to the stable homotopy of a Moore space*, J. Pure Appl. Algebra **20** (1981), 287–312.
- Miller, H. R. and Ravenel, D. C.
- [5] *Morava stabilizer algebras and the localization of Novikov's E_2 -term*, Duke Math. J. **44** (1977), no. 2, 433–447.
- Miller, H. R.
- [6] *An algebraic analogue of a conjecture of G. W. Whitehead*, Proc. Amer. Math. Soc. **84** (1982), 131–137.
- [7] *A localization theorem in homological algebra*, Math. Proc. Cambridge Philos. Soc. **84** (1978), 73–84.
- Miller, H. R. and Wilkerson, C.
- [8] *Vanishing lines for modules over the Steenrod algebra*, J. Pure Appl. Algebra **22** (1981), 293–307.
- Miller, H. and Wilkerson, C.
- [9] *On the Segal conjecture for periodic groups*, Proc. Northwestern Homotopy Theory Conf., Contemp. Math., 1983, pp. 233–246.
- Miller, H. R.
- [10] *The Sullivan conjecture on maps from classifying spaces*, Ann. of Math. (2) **120** (1984), no. 1, 39–87.
- Milnor, J. W.
- [1] *Morse theory*, Ann. of Math., Princeton University Press, Princeton, N.J., 1963.
- [2] *The Steenrod algebra and its dual*, Ann. of Math. **67** (1958), 150–171.
- Milnor, J. W. and Moore, J. C.
- [3] *On the structure of Hopf algebra*, Ann. of Math. **81** (1965), 211–264.
- Milnor, J. W.
- [4] *On the cobordism ring Ω^* and a complex analogue*, Amer. J. Math. **82** (1960), 505–521.
- Milnor, J. W. and Stasheff, J. D.
- [5] *Characteristic classes*, Ann. of Math. Studies, Princeton University Press, Princeton, N. J., 1974.
- Mironov, O. K.
- [1] *Existence of multiplicative structures in the theory of cobordism with singularities*, Izv. Akad. Nauk SSSR Ser. Mat. **39** (1975), no. 5, 1065–1092, 1219.
- Mischenko, A. S.
- [1] Appendix 1 in Novikov [1].
- Moore, J. C. and Peterson, F. P.
- [1] *Nearly Frobenius algebras, Poincaré algebras and their modules*, J. Pure Appl. Algebra **3** (1973), 83–93.
- Morava, J.
- [1] *A product for the odd primary bordism of manifolds with singularities*, Topology **18** (1979), 177–186.
- [2] *Completions of complex cobordism*, Geometric applications of homotopy theory (Proc. Conf., Evanston, Ill., 1977), II, Lecture Notes in Math., Springer, Berlin, 1978, pp. 349–361.
- [3] *Noetherian localisations of categories of cobordism comodules*, Ann. of Math. **121** (1985), 1–39.
- Moreira, M. R. F.
- [1] *Primitives of BP_*BP modulo an invariant prime ideal*, Amer. J. Math. **100** (1975), 1247–1273.
- [2] *Formal groups and the BP spectrum*, J. Pure Appl. Algebra **18** (1960), 79–89.
- [3] *The last distinguished primitive of BP_*BP modulo an invariant prime ideal*, to appear.
- Moss, R. M. F.
- [1] *On the compositon pairing of Adams spectral sequences*, Proc. London Math. Soc. (3) **18** (1968), 179–182.

Mahowald, M., Ravenel, D., and Shick, P.

- [1] *The triple loop space approach to the telescope conjecture*, Homotopy methods in algebraic topology (Boulder, CO, 1999), Contemp. Math., vol. 271, Amer. Math. Soc., Providence, RI, 2001, pp. 217–284.

Mumford, D.

- [1] *Lectures on curves on an algebraic surface*, With a section by G. M. Bergman. Annals of Mathematics Studies, No. 59, Princeton University Press, Princeton, N.J., 1966.

Nakamura, O.

- [1] *On the squaring operations in the May spectral sequence*, Mem. Fac. Sci. Kyushu Univ. Ser. A **26** (1972), 293–308.
 [2] *On the cohomology of the mod p Steenrod algebra*, Bull. Sci. Engrg. Dig. Univ. Ryukyus Math. Natur. Sci. **18** (1957), 9–58.
 [3] *Some differentials in the mod 3 Adams spectral sequence*, Bull. Sci. Engrg. Div. Univ. Ryukyus Math. Natur. Sci. (1975), no. 19, 1–25.

Nassau, C.

- [1] *Steenrod algebra cohomology*, Online at <http://www.nullhomotopie.de/maths/index.html>.

Neisendorfer, J. A.

- [1] *3-primary exponents*, Math. Proc. Cambridge Philos. Soc. **90** (1981), no. 1, 63–83.

Times, N. Y.

- [1] New York Times, Editorial Page, June 2 1976.

Nishida, G.

- [1] *The nilpotency of elements of the stable homotopy groups of spheres*, J. Math. Soc. Japan **25** (1973), 707–732.

Novikov, S. P.

- [1] *The methods of algebraic topology from the viewpoint of cobordism theories*, Izv. Akad. Nauk. SSSR. Ser. Mat. **31** (1967), 855–951 (Russian).
 [2] *Some problems in the topology of manifolds connected with the theory of Thom spaces*, Dokl. Akad. Nauk. SSSR. **132** (1960), 1031–1034 (Russian).
 [3] *Homotopy properties of Thom complexes*, Mat. Sb. (N.S.) **57 (99)** (1962), 407–442.

Oka, S.

- [1] *The stable homotopy groups of spheres. I*, Hiroshima Math. J. **1** (1971), 305–337.
 [2] *The stable homotopy groups of spheres. II*, Hiroshima Math. J. **2** (1972), 99–161.
 [3] *The stable homotopy groups of spheres. III*, Hiroshima Math. J. **5** (1975), no. 3, 407–438.
 [4] *A new family in the stable homotopy groups of spheres*, Hiroshima Math. J. **5** (1975), 87–114.
 [5] *A new family in the stable homotopy groups of spheres. II*, Hiroshima Math. J. **6** (1976), no. 2, 331–342.
 [6] *Realizing some cyclic BP_* -modules and applications to stable homotopy groups of spheres*, Hiroshima Math. J. **7** (1977), 427–447.
 [7] *Ring spectra with few cells*, Japan. J. Math. **5** (1979), 81–100.

Oka, S. and Toda, H.

- [8] *3-primary β -family in stable homotopy*, Hiroshima Math. J. **5** (1975), 447–460.

Oka, S. and Shimomura, K.

- [9] *On products of the β -element in the stable homotopy of spheres*, Hiroshima Math. J. **12** (1982), 611–626.

Oka, S.

- [10] *Small ring spectra with p -rank of the stable homotopy of spheres*, Proc. Northwestern Univ. Homotopy Theory Conf., Contemp. Math., vol. 19, 1983, pp. 282–308.

Palmieri, J. H.

- [1] *Quillen stratification for the Steenrod algebra*, Ann. of Math. (2) **149** (1999), no. 2, 421–449.
 [2] *Stable homotopy over the Steenrod algebra*, Mem. Amer. Math. Soc. **151** (2001), no. 716, xiv+172.

Pengelley, D. J.

- [1] *The homotopy type of MSU*, Amer. J. Math. **104** (1982), 1101–1123.
- [2] *The homology of MSO and MSU as A^* comodule algebras and the cobordism ring*, J. London Math. Soc. **25** (1982), 467–472.

Priddy, S. B.

- [1] *A cellular construction of BP and other irreducible spectra*, Math. Z. **173** (1980), no. 1, 29–34.

Quillen, D. G.

- [1] *The Adams conjecture*, Topology **10** (1971), 1–10.
- [2] *On the formal group laws of unoriented and complex cobordism theory*, Bull. Amer. Math. Soc. **75** (1969), 1293–1298.

Quillen, D.

- [3] *The spectrum of an equivariant cohomology ring. I, II*, Ann. of Math. (2) **94** (1971), 549–572; *ibid.* (2) **94** (1971), **573–602**.

Ravenel, D. C.

- [1] *The structure of BP_*BP modulo an invariant prime ideal*, Topology **15** (1976), 149–153.

Ravenel, D. C. and Wilson, W. S.

- [2] *The Hopf ring for complex cobordism*, J. Pure Appl. Algebra **9** (1977), 241–280.
- [3] *The Morava K -theories of Eilenberg–Mac Lane spaces and the Conner–Floyd conjecture*, Amer. J. Math. **102** (1980), 691–748.

Ravenel, D. C.

- [4] *The Segal conjecture for cyclic groups and its consequences*, Amer. J. Math. **106** (1984), 415–446.
- [5] *The structure of Morava stabilizer algebras*, Inrvnt. Math. **37** (1976), 109–120.
- [6] *The cohomology of the Morava stabilizer algebras*, Math. Z. **152** (1977), 287–297.
- [7] *The nonexistence of odd primary Arf invariant elements in stable homotopy*, Math. Proc. Cambridge Philos. Soc. **83** (1987), 429–433.
- [8] *Localisation with respect to certain periodic homology theories*, Amer. J. Math. **106** (1984), 351–414.
- [9] *The geometric realization of the chromatic resolution*, Algebraic topology and algebraic K -theory (Princeton, N.J., 1983), Princeton Univ. Press, Princeton, NJ, 1987, pp. 168–179.
- [10] *Formal A -modules and the Adams–Novikov spectral sequence*, J. Pure Appl. Algebra **32** (1984), 327–345.
- [11] *The Adams–Novikov E_2 -term for a complex with p -cells*, Amer. J. Math. **107** (1985), no. 4, 933–968.
- [12] *The method of infinite descent in stable homotopy theory. I*, Recent progress in homotopy theory (Baltimore, MD, 2000), Contemp. Math., vol. 293, Amer. Math. Soc., Providence, RI, 2002, pp. 251–284.
- [13] *Nilpotence and periodicity in stable homotopy theory*, Annals of Mathematics Studies, vol. 128, Princeton University Press, Princeton, 1992.

Rezk, C.

- [1] *Notes on the Hopkins–Miller theorem*, Homotopy theory via algebraic geometry and group representations (Evanston, IL, 1997), Contemp. Math., vol. 220, Amer. Math. Soc., Providence, RI, 1998, pp. 313–366.

Richter, W.

- [1] *Lambda algebra unstable composition products and the Lambda EHP sequence*, Online at <http://hopf.math.purdue.edu/cgi-bin/generate?/Richter/Richter-Lambda-EHP>.

Schwartz, L.

- [1] *Unstable modules over the Steenrod algebra and Sullivan’s fixed point set conjecture*, Chicago Lectures in Mathematics, University of Chicago Press, Chicago, IL, 1994.

Science

- [1] Science, June, 7 1976.

Serre, J.-P.

- [1] *Local fields*, Springer-Verlag, New York, 1979.
- [2] *Homologie singulière des espace fibrés*, Ann. of Math. **54** (1951), 425–505.
- [3] *Groupes d'homotopie et classes de groupes abélien*, Ann. of Math. **58** (1953), 258–294.

Shimada, N. and Yagita, N.

- [1] *Multiplication in the complex bordism theory with with singularities*, Publ. Res. Inst. Math. Sci. **12** (1976–1977), no. 1, 259–293.

Shimada, N. and Iwai, A.

- [2] *On the cohomology of some Hopf algebras*, Nagoya Math. J. **30** (1967), 103–111.

Shimada, N. and Yamanoshita, T.

- [3] *On the triviality of the mod p Hopf invariant*, Japan. J. Math. **31** (1961), 1–24.

Shimomura, K.

- [1] *Novikov's Ext^2 at the prime 2*, Hiroshima Math. J. **11** (1981), 499–513.
- [2] *Chromatic E_1 -terms – up to April 1995*, J. Fac. Educ. Tottori Univ. (Nat. Sci.) **44** (1995), 1–6.

Shimomura, K. and Wang, X.

- [3] *The Adams–Novikov E_2 -term for $\pi_*(L_2S^0)$ at the prime 2*, Math. Z. **241** (2002), no. 2, 271–311.
- [4] *The homotopy groups $\pi_*(L_2S^0)$ at the prime 3*, Topology **41** (2002), no. 6, 1183–1198.

Shimomura, K. and Yabe, A.

- [5] *The homotopy groups $\pi_*(L_2S^0)$* , Topology **34** (1995), no. 2, 261–289.

Shimomura, K.

- [6] *The Adams–Novikov E_2 -term for computing $\pi_*(L_2V(0))$ at the prime 2*, Topology Appl. **96** (1999), no. 2, 133–152.
- [7] *The homotopy groups of the L_2 -localized mod 3 Moore spectrum*, J. Math. Soc. Japan **52** (2000), no. 1, 65–90.
- [8] *On the Adams–Novikov spectral sequence and products of β -elements*, Hiroshima Math. J. **16** (1986), no. 1, 209–224.

Siegel, C. L.

- [1] *Topics in complex function theory. Vol. I: Elliptic functions and uniformization theory*, Wiley (Interscience), New York, 1969.

Silverman, J. H.

- [1] *The arithmetic of elliptic curves*, Graduate Texts in Math., vol. 106, Springer-Verlag, New York, 1986.

Singer, W. M.

- [1] *A new chain complex for the homology of the Steenrod algebra*, Math. Proc. Cambridge Philos. Soc. **90** (1981), 279–292.
- [2] *Invariant theory and the Lambda algebra*, Trans. Amer. Math. Soc. **280** (1983), 573–693.
- [3] *The algebraic EHP sequence*, Trans. Amer. Math. Soc. **201** (1975), 367–382.
- [4] *Squaring operations and the algebraic EHP sequence*, J. Pure Appl. Algebra **6** (1975), 13–29.
- [5] *Extension theory for connected Hopf algebras*, J. Algebra **21** (1972), 1–16.
- [6] *Steenrod squares in spectral sequences. I, II*, Trans. Amer. Math. Soc. **175** (1973), I, 327–336; II, 337–353.

Smith, L.

- [1] *On realizing complex bordism modules*, Amer. J. Math. **92** (1970), 793–856.
- [2] *On realizing complex bordism modules, IV, Applications to the stable homotopy groups of spheres*, Amer. J. Math. **99** (1971), 418–436.

Snaith, V. P.

- [1] *Stable decomposition of $\Omega^n \Sigma^n X$* , J. London Math. Soc. (2) **7** (1974), 577–583.

Spanier, E. H.

- [1] *Algebraic topology*, McGraw-Hill, New York, 1966.

Steenrod, N. E. and Epstein, D. B. A.

- [1] *Cohomology operations*, Ann. of Math. Studies, no. 50, Princeton University Press, Princeton, N. J., 1962.

Stong, R. E.

- [1] *Notes on cobordism theory*, Princeton University Press, Princeton, N.J., 1968.
 [2] *Determination of $H^*(BO(k, \dots, \infty), \mathbb{Z}_2)$ and $H^*(BU(k, \dots, \infty), \mathbb{Z}_2)$* , Trans. Amer. Math. Soc. **107** (1963), 526–544.
 [3] *Cobordism of maps*, Topology **5** (1966), 245–258.

Strickland, N. P.

- [1] *Formal schemes and formal groups*, Homotopy invariant algebraic structures (Baltimore, MD, 1998), Contemp. Math., vol. 239, Amer. Math. Soc., Providence, RI, 1999, pp. 263–352.

Sullivan, D. P.

- [1] *Genetics of homotopy theory and the Adams conjecture*, Ann. of Math. **100** (1974), 1–79.

Sweedler, M. E.

- [1] *Hopf algebras*, Benjamin, New York, 1969.

Switzer, R. M.

- [1] *Algebraic topology—homotopy and homology*, Springer-Verlag, New York, 1975.

Tangora, M. C.

- [1] *On the cohomology of the Steenrod algebra*, Math. Z. **116** (1970), 18–64.
 [2] *Some remarks on the lambda algebras*, Geometric applications of homotopy theory (Proc. Conf., Evanston, Ill., 1977), II, Lecture Notes in Math., Springer, Berlin, 1978, pp. 476–487.
 [3] *Generating Curtis tables*, Algebraic topology (Proc. Conf., Univ. British Columbia, Vancouver, B.C., 1977), Lecture Notes in Math., Springer, Berlin, 1978, pp. 243–253.
 [4] *Computing the homology of the lambda algebra*, Mem. Amer. Math. Soc. **58** (1985), no. 337, v+163.
 [5] *Some extension questions in the Adams spectral sequence*, Proc. Adv. Study Inst. Algebra Top., Aarhus, Mat. Inst., 1970, pp. 578–587.
 [6] *Some homotopy groups mod 3*, Conference on homotopy theory (Evanston, Ill., 1974), Soc. Mat. Mexicana, México, 1975, pp. 227–245.

Thom, R.

- [1] *Quelques propriétés globales des variétés différentiables*, Comm. Math. Helv. **28** (1954), 109–181.

Thomas, E. and Zahler, R. S.

- [1] *Nontriviality of the stable homotopy element γ_1* , J. Pure and Applied Algebra **4** (1974), 189–203.

Toda, H.

- [1] *On spectra realizing exterior parts of the Steenrod algebra*, Topology **10** (1971), 53–65.
 [2] *Extended p -th powers of complexes and applications to homotopy theory*, Proc. Japan Acad. **44** (1968), 198–203.
 [3] *An important relation in the homotopy groups of spheres*, Proc. Japan. Acad. **43** (1967), 893–942.
 [4] *p -primary components of homotopy groups. I. Exact sequences in Steenrod algebra; II. Mod p Hopf invariant; III. Stable groups of the sphere*, Mem. Coll. Sci. Univ. Kyoto. Ser. A. Math. **31** (1958), 129–142, 143–160, 191–210.
 [5] *p -primary components of homotopy groups. IV. Compositions and toric constructions*, Mem. Coll. Sci. Univ. Kyoto. Ser. A. Math. **32** (1959), 297–332.
 [6] *Composition methods in homotopy groups of spheres*, Ann. of Math. Studies, no. 49, Princeton University Press, Princeton, N.J., 1962.
 [7] *Algebra of stable homotopy of \mathbb{Z}_p -spaces and applications*, J. Math. Kyoto Univ. **11** (1971), 197–251.
 [8] *Unstable 3-primary homotopy groups of spheres*, Econoinformatics **29** (2003), 1–77.

tom Dieck, T.

- [1] *Actions of finite abelian groups without stationary points*, Topology **9** (1970), 359–366.

Wang, J. S. P.

- [1] *On the cohomology of the mod 2 Steenrod algebra and the non-existence of elements of Hopf invariant one*, Illinois J. Math. **11** (1967), 480–490.

Welcher, P. J.

- [1] *Symmetric products and the stable Hurewicz homomorphism*, Illinois J. Math. **24** (1980), 527–544.
 [2] *Symmetric fiber spectra $K(n)$ -homology acyclicity*, Indiana J. Math. **30** (1981), 801–812.

Wellington, R. J.

- [1] *The computation of Ext groups for modules over the Steenrod algebra*, To appear.

Wilson, W. S.

- [1] *The Ω -spectrum for cohomology. Part I*, Comment. Math. Helv. **48** (1973), 45–55.
 [2] *The Ω -spectrum for cohomology. Part II*, Amer. J. Math. **97** (1975), 123.
 [3] *Brown–Peterson homology, an introduction and sampler*, Regional Conference series in Math. (Rhode Island), no. 48, AMS, Providence, 1980.

Würgler, U.

- [1] *On products in a family of cohomology theories associated to the invariant prime ideals of $\pi_*(BP)$* , Comment. Math. Helv. **52** (1977), no. 4, 457–481.
 [2] *A splitting theorem for certain cohomology theories associated to $BP^*(-)$* , Manuscripta Math. **29** (1979), no. 1, 93–111.
 [3] *On the relation of Morava K -theories to Brown–Peterson homology*, Topology and algebra (Proc. Colloq., Eidgenöss. Tech. Hochsch., Zurich, 1977), Monograph. Enseign. Math., vol. 26, Univ. Genève, Geneva, 1978, pp. 269–280.
 [4] *Cobordism theories of unitary manifolds with singularities and formal group laws*, Math. Z. **150** (1976), no. 3, 239–260.

Yagita, N.

- [1] *On the algebraic structure of cobordism operations with singularities*, J. London Math. Soc. **16** (1977), 131–141.
 [2] *On the Steenrod algebra of Morava K -theory*, J. London Math. Soc. **22** (1980), 423–438.
 [3] *The exact functor theorem for BP_*/I_n -theory*, Proc. Japan Acad. **52** (1976), 1–3.

Yosimura, Z.

- [1] *Projective dimension of Brown–Peterson homology with modulo (p, v_1, \dots, v_{n-1}) coefficients*, Osaka J. Math. **13** (1976), 289–309.

Zachariou, A.

- [1] *A subalgebra of $\text{Ext}_A(\mathbf{Z}_2, \mathbf{Z}_2)$* , Bull. Amer. Math. Soc. **73** (1967), 647–648.

Zahler, R.

- [1] *The Adams–Novikov spectral sequence for the spheres*, Ann. of Math. **96** (1972), 480–504.
 [2] *Fringe families in stable homotopy*, Trans. Amer. Math. Soc. **224** (1976), 243–253.