



# **An introduction to elliptic cohomology and topological modular forms**

*Wayne State University Topology Seminar  
February 27, 2007*

Doug Ravenel

# What is elliptic cohomology?

**Definition 1** *For a ring  $R$ , an  $R$ -valued genus on a class of closed manifolds is a function  $\varphi$  that assigns to each manifold  $M$  an element  $\varphi(M) \in R$  such that*

# What is elliptic cohomology?

**Definition 1** *For a ring  $R$ , an  $R$ -valued genus on a class of closed manifolds is a function  $\varphi$  that assigns to each manifold  $M$  an element  $\varphi(M) \in R$  such that*

$$(i) \quad \varphi(M_1 \amalg M_2) = \varphi(M_1) + \varphi(M_2)$$

# What is elliptic cohomology?

**Definition 1** *For a ring  $R$ , an  $R$ -valued genus on a class of closed manifolds is a function  $\varphi$  that assigns to each manifold  $M$  an element  $\varphi(M) \in R$  such that*

$$(i) \quad \varphi(M_1 \amalg M_2) = \varphi(M_1) + \varphi(M_2)$$

$$(ii) \quad \varphi(M_1 \times M_2) = \varphi(M_1)\varphi(M_2)$$

# What is elliptic cohomology?

**Definition 1** *For a ring  $R$ , an  $R$ -valued genus on a class of closed manifolds is a function  $\varphi$  that assigns to each manifold  $M$  an element  $\varphi(M) \in R$  such that*

- (i)  $\varphi(M_1 \amalg M_2) = \varphi(M_1) + \varphi(M_2)$*
- (ii)  $\varphi(M_1 \times M_2) = \varphi(M_1)\varphi(M_2)$*
- (iii)  $\varphi(M) = 0$  if  $M$  is a boundary.*

# What is elliptic cohomology?

**Definition 1** *For a ring  $R$ , an  $R$ -valued genus on a class of closed manifolds is a function  $\varphi$  that assigns to each manifold  $M$  an element  $\varphi(M) \in R$  such that*

- (i)  $\varphi(M_1 \amalg M_2) = \varphi(M_1) + \varphi(M_2)$*
- (ii)  $\varphi(M_1 \times M_2) = \varphi(M_1)\varphi(M_2)$*
- (iii)  $\varphi(M) = 0$  if  $M$  is a boundary.*

*Equivalently,  $\varphi$  is a homomorphism from the appropriate cobordism ring  $\Omega$  to  $R$ .*

# What is elliptic cohomology?

**Definition 2** *A 1-dimensional formal group law over  $R$  is a power series  $F(x, y) \in R[[x, y]]$  satisfying*

# What is elliptic cohomology?

**Definition 2** *A 1-dimensional formal group law over  $R$  is a power series  $F(x, y) \in R[[x, y]]$  satisfying*

$$(i) \quad F(x, 0) = F(0, x) = x.$$



# What is elliptic cohomology?

**Definition 2** *A 1-dimensional formal group law over  $R$  is a power series  $F(x, y) \in R[[x, y]]$  satisfying*

(i)  $F(x, 0) = F(0, x) = x.$

(ii)  $F(y, x) = F(x, y)$

# What is elliptic cohomology?

**Definition 2** *A 1-dimensional formal group law over  $R$  is a power series  $F(x, y) \in R[[x, y]]$  satisfying*

- (i)  $F(x, 0) = F(0, x) = x$ .*
- (ii)  $F(y, x) = F(x, y)$*
- (iii)  $F(x, F(y, z)) = F(F(x, y), z)$ .*

# What is elliptic cohomology?

A theorem of Quillen says that in the complex case (where  $\Omega = MU_*$ , the complex cobordism ring),  $\varphi$  is equivalent to a 1-dimensional formal group law over  $R$ .

# What is elliptic cohomology?

A theorem of Quillen says that in the complex case (where  $\Omega = MU_*$ , the complex cobordism ring),  $\varphi$  is equivalent to a 1-dimensional formal group law over  $R$ . More precisely, there is a 1-dimensional formal group law  $G$  over  $MU_*$  having a certain universal property. It is defined as follows.

# What is elliptic cohomology?

A theorem of Quillen says that in the complex case (where  $\Omega = MU_*$ , the complex cobordism ring),  $\varphi$  is equivalent to a 1-dimensional formal group law over  $R$ . More precisely, there is a 1-dimensional formal group law  $G$  over  $MU_*$  having a certain universal property. It is defined as follows.

Let  $\alpha$  be a complex line bundle over a space  $X$ . It has a Conner-Floyd Chern class  $c_1(\alpha) \in MU^2(X)$ .

# What is elliptic cohomology?

A theorem of Quillen says that in the complex case (where  $\Omega = MU_*$ , the complex cobordism ring),  $\varphi$  is equivalent to a 1-dimensional formal group law over  $R$ . More precisely, there is a 1-dimensional formal group law  $G$  over  $MU_*$  having a certain universal property. It is defined as follows.

Let  $\alpha$  be a complex line bundle over a space  $X$ . It has a Conner-Floyd Chern class  $c_1(\alpha) \in MU^2(X)$ . Given two such line bundles  $\alpha_1$  and  $\alpha_2$ , we have

$$c_1(\alpha_1 \otimes \alpha_2) = G(c_1(\alpha_1), c_1(\alpha_2))$$

where  $G$  is the desired formal group law.

# What is elliptic cohomology?

This formal group law  $G$  over  $MU_*$  has the following universal property: Any formal group law  $F$  over a ring  $R$  is induced from  $G$  via a homomorphism

$$\theta : MU_* \rightarrow R.$$

# What is elliptic cohomology?

This formal group law  $G$  over  $MU_*$  has the following universal property: Any formal group law  $F$  over a ring  $R$  is induced from  $G$  via a homomorphism

$$\theta : MU_* \rightarrow R.$$

It follows that an  $R$ -valued genus on complex manifolds is equivalent to a 1-dimensional formal group law over  $R$ .



# What is elliptic cohomology?

This formal group law  $G$  over  $MU_*$  has the following universal property: Any formal group law  $F$  over a ring  $R$  is induced from  $G$  via a homomorphism

$$\theta : MU_* \rightarrow R.$$

It follows that an  $R$ -valued genus on complex manifolds is equivalent to a 1-dimensional formal group law over  $R$ .

It is also known that the functor

$$X \mapsto MU_*(X) \otimes_{\varphi} R$$

is a homology theory if  $\varphi$  satisfies certain conditions spelled out in Landweber's Exact Functor Theorem.

# What is elliptic cohomology?

Now suppose  $E$  is an elliptic curve defined over  $R$ . It is a 1-dimensional algebraic group, and choosing a local parameter at the identity leads to a formal group law  $\widehat{E}$ , the formal completion of  $E$ . Thus we can apply the machinery above and get an  $R$ -valued genus.

# What is elliptic cohomology?

Now suppose  $E$  is an elliptic curve defined over  $R$ . It is a 1-dimensional algebraic group, and choosing a local parameter at the identity leads to a formal group law  $\widehat{E}$ , the formal completion of  $E$ . Thus we can apply the machinery above and get an  $R$ -valued genus.

For example, the *Jacobi quartic*, defined by the equation

$$y^2 = 1 - 2\delta x^2 + \epsilon x^4,$$

is an elliptic curve over the ring

$$R = \mathbf{Z}[1/2, \delta, \epsilon].$$

# What is elliptic cohomology?

The resulting formal group law is the power series expansion of

$$F(x, y) = \frac{x\sqrt{1 - 2\delta y^2 + \epsilon y^4} + y\sqrt{1 - 2\delta x^2 + \epsilon x^4}}{1 - \epsilon x^2 y^2};$$

this calculation is originally due to Euler. The resulting genus is known to satisfy Landweber's conditions, and this leads to one definition of elliptic cohomology.

# What are modular forms?

Recall that an elliptic curve is determined by a lattice in  $\mathbf{C}$  generated by 1 and a complex number  $\tau$  in the upper half plane  $H$ . There is an action of the group  $SL_2(\mathbf{Z})$  on  $H$  given by

$$\tau \mapsto \tau' = \frac{a\tau + b}{c\tau + d} \quad \text{for} \quad \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in SL_2(\mathbf{Z}).$$

# What are modular forms?

Recall that an elliptic curve is determined by a lattice in  $\mathbf{C}$  generated by 1 and a complex number  $\tau$  in the upper half plane  $H$ . There is an action of the group  $SL_2(\mathbf{Z})$  on  $H$  given by

$$\tau \mapsto \tau' = \frac{a\tau + b}{c\tau + d} \quad \text{for} \quad \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in SL_2(\mathbf{Z}).$$

An easy calculation shows that  $\tau'$  determines the same lattice as  $\tau$ .

# What are modular forms?

Recall that an elliptic curve is determined by a lattice in  $\mathbf{C}$  generated by 1 and a complex number  $\tau$  in the upper half plane  $H$ . There is an action of the group  $SL_2(\mathbf{Z})$  on  $H$  given by

$$\tau \mapsto \tau' = \frac{a\tau + b}{c\tau + d} \quad \text{for} \quad \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in SL_2(\mathbf{Z}).$$

An easy calculation shows that  $\tau'$  determines the same lattice as  $\tau$ . *This means that elliptic curves are parametrized by the orbits under this action.*

# What are modular forms?

A *modular form of weight  $k$*  is a meromorphic function  $g$  defined on the upper half plane satisfying

$$g\left(\frac{a\tau + b}{c\tau + d}\right) = (c\tau + d)^k g(\tau).$$



# What are modular forms?

A *modular form of weight  $k$*  is a meromorphic function  $g$  defined on the upper half plane satisfying

$$g\left(\frac{a\tau + b}{c\tau + d}\right) = (c\tau + d)^k g(\tau).$$

Here is an example. Let

$$G_k(\tau) := \sum'_{m,n \in \mathbf{Z}} \frac{1}{(m\tau + n)^k}$$

where the sum is over all nonzero lattice points. This vanishes if  $k$  is odd and is known to converge for  $k > 2$ .

# What are modular forms?

Note that

$$\begin{aligned} G_k \left( \frac{a\tau + b}{c\tau + d} \right) &= \sum'_{m,n \in \mathbf{Z}} \left( \frac{c\tau + d}{m(a\tau + b) + n(c\tau + d)} \right)^k \\ &= (c\tau + d)^k G_k(\tau). \end{aligned}$$

so  $G_k$  is a modular form of weight  $k$ .

# What are modular forms?

Now let  $q = e^{2\pi i\tau}$ . In terms of it we have

$$G_k(\tau) = 2(2\pi i)^k \frac{B_k}{2k!} \left( 1 - \frac{2k}{B_k} \sum_{n=1}^{\infty} \sigma_{k-1}(n) q^n \right)$$

# What are modular forms?

Now let  $q = e^{2\pi i\tau}$ . In terms of it we have

$$G_k(\tau) = 2(2\pi i)^k \frac{B_k}{2k!} \left( 1 - \frac{2k}{B_k} \sum_{n=1}^{\infty} \sigma_{k-1}(n) q^n \right)$$

where  $B_k$  is the  $k$ th Bernoulli number defined by

$$\frac{x}{e^x - 1} =: \sum_{k \geq 0} B_k \frac{x^k}{k!},$$

# What are modular forms?

Now let  $q = e^{2\pi i\tau}$ . In terms of it we have

$$G_k(\tau) = 2(2\pi i)^k \frac{B_k}{2k!} \left( 1 - \frac{2k}{B_k} \sum_{n=1}^{\infty} \sigma_{k-1}(n) q^n \right)$$

where  $B_k$  is the  $k$ th Bernoulli number defined by

$$\frac{x}{e^x - 1} =: \sum_{k \geq 0} B_k \frac{x^k}{k!},$$

and

$$\sigma_{k-1}(n) := \sum_{d|n} d^{k-1}.$$

# What are modular forms?

It is convenient to normalize  $G_k$  by defining the Eisenstein series

$$E_k(\tau) := 1 - \frac{2k}{B_k} \sum_{n=1}^{\infty} \sigma_{k-1}(n) q^n.$$

# What are modular forms?

It is convenient to normalize  $G_k$  by defining the Eisenstein series

$$E_k(\tau) := 1 - \frac{2k}{B_k} \sum_{n=1}^{\infty} \sigma_{k-1}(n) q^n.$$

It turns out that

$$E_k(\tau) = \sum_{(m,n)=1} \frac{1}{(m\tau + n)^k},$$

where the sum is over pairs of integers that are relatively prime.

# What are modular forms?

Let

$$\Delta := \frac{E_4(\tau)^3 - E_6(\tau)^2}{1728}$$

$$= q \prod_{n=1}^{\infty} (1 - q^n)^{24}$$

and  $j := E_4^3 / \Delta$ .

$\Delta$  is called the discriminant, and the modular function  $j$  of weight 0 is a complex analytic isomorphism between  $H/SL_2(\mathbf{Z})$  and the Riemann sphere.



# What are modular forms?

It is known that the ring of all modular forms with respect to  $\Gamma = SL_2(\mathbf{Z})$  is

$$M_*(\Gamma) = \mathbf{C}[E_4, E_6],$$

with  $(\Delta)$  being the ideal of forms that vanish at  $i\infty$ , which are called *cuspidal forms*.

# The Weierstrass curve

The *Weierstrass equation* for an elliptic curve in affine form is

$$y^2 + a_1xy + a_3y = x^3 + a_2x^2 + a_4x + a_6.$$

# The Weierstrass curve

The *Weierstrass equation* for an elliptic curve in affine form is

$$y^2 + a_1xy + a_3y = x^3 + a_2x^2 + a_4x + a_6.$$

It is known that any elliptic curve is isomorphic to one of this form.

# The Weierstrass curve

The Eisenstein series are related to the  $a_k$  by

$$E_4 = b_2^2 - 24b_4$$

and

$$E_6 = -b_2^3 + 36b_2b_4 - 216b_6$$

where

$$b_2 = a_1^2 + 4a_2$$

$$b_4 = a_1a_3 + 2a_4$$

$$b_6 = a_3^2 + 4a_6.$$

# The Weierstrass curve

Then

$$\Delta = \frac{E_4^3 - E_6^2}{1728}$$

is the discriminant of the Weierstrass equation, meaning that the curve is smooth iff  $\Delta$  is a unit in the ground ring.

# The Weierstrass curve

Then

$$\Delta = \frac{E_4^3 - E_6^2}{1728}$$

is the discriminant of the Weierstrass equation, meaning that the curve is smooth iff  $\Delta$  is a unit in the ground ring.

This means there is a formal group law defined over the ring

$$Ell_* = \mathbf{Z}[a_1, a_2, a_3, a_4, a_6][\Delta^{-1}]$$

and the resulting genus is Landweber exact.

# The Weierstrass curve

Then

$$\Delta = \frac{E_4^3 - E_6^2}{1728}$$

is the discriminant of the Weierstrass equation, meaning that the curve is smooth iff  $\Delta$  is a unit in the ground ring.

This means there is a formal group law defined over the ring

$$Ell_* = \mathbf{Z}[a_1, a_2, a_3, a_4, a_6][\Delta^{-1}]$$

and the resulting genus is Landweber exact. Thus we get a spectrum  $Ell$  with  $\pi_*(Ell) = Ell_*$ .

# The Weierstrass curve

Consider the affine coordinate change

$$y \mapsto y + r \quad \text{and} \quad x \mapsto x + sy + t$$



# The Weierstrass curve

Consider the affine coordinate change

$$y \mapsto y + r \quad \text{and} \quad x \mapsto x + sy + t$$

It can be thought of as an action of an affine group  $G$  of  $3 \times 3$  matrices given by

$$\begin{bmatrix} 1 & s & t \\ 0 & 1 & r \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} = \begin{bmatrix} x + sy + t \\ y + r \\ 1 \end{bmatrix}$$

# The Weierstrass curve

Under it we get

$$a_6 \mapsto a_6 + a_4 r + a_3 t + a_2 r^2 + a_1 r t + t^2 - r^3$$

$$a_4 \mapsto a_4 + a_3 s + 2 a_2 r + a_1 (r s + t) + 2 s t - 3 r^2$$

$$a_3 \mapsto a_3 + a_1 r + 2 t$$

$$a_2 \mapsto a_2 + a_1 s - 3 r + s^2$$

$$a_1 \mapsto a_1 + 2 s.$$

# The spectrum $tmf$

Very roughly speaking, there is an action of the group  $G$  on the spectrum  $Ell$  inducing the above action on its homotopy groups.

# The spectrum $tmf$

Very roughly speaking, there is an action of the group  $G$  on the spectrum  $Ell$  inducing the above action on its homotopy groups.

The homotopy fixed point set of this action is  $tmf$ .

# The spectrum $tmf$

Very roughly speaking, there is an action of the group  $G$  on the spectrum  $Ell$  inducing the above action on its homotopy groups.

The homotopy fixed point set of this action is  $tmf$ .

There is a spectral sequence converging to  $\pi_*(tmf)$  with

$$E_2 = H^*(G; Ell_*).$$

# The spectrum $tmf$

Very roughly speaking, there is an action of the group  $G$  on the spectrum  $Ell$  inducing the above action on its homotopy groups.

The homotopy fixed point set of this action is  $tmf$ .

There is a spectral sequence converging to  $\pi_*(tmf)$  with

$$E_2 = H^*(G; Ell_*).$$

The modular forms  $E_4$  and  $E_6$  are both invariant under this action, and

$$H^0(G; Ell_*) = \mathbf{Z}[E_4, E_6][\Delta^{-1}]$$

# The spectrum $tmf$

The coordinate change above can be used to define a Hopf algebroid  $(A, \Gamma)$  with

$$A = \mathbf{Z}[a_1, a_2, a_3, a_4, a_6]$$

$$\Gamma = A[r, s, t]$$

and right unit  $\eta_R : A \rightarrow \Gamma$  given by the formulas above.

# The spectrum $tmf$

The coordinate change above can be used to define a Hopf algebroid  $(A, \Gamma)$  with

$$\begin{aligned} A &= \mathbf{Z}[a_1, a_2, a_3, a_4, a_6] \\ \Gamma &= A[r, s, t] \end{aligned}$$

and right unit  $\eta_R : A \rightarrow \Gamma$  given by the formulas above. It was first described by Hopkins and Mahowald in *From elliptic curves to homotopy theory*. Its Ext group is the cohomology group mentioned above. Tilman Bauer has written a nice account of this calculation.