

**TOWARD HIGHER CHROMATIC ANALOGS OF ELLIPTIC
COHOMOLOGY AND TOPOLOGICAL MODULAR FORMS**

WAYNE STATE UNIVERSITY TOPOLOGY SEMINAR

MARCH 1, 2007

DOUG RAVENEL

WHAT DOES “CHROMATIC” MEAN?

The stable homotopy category localized at a prime p can be studied via a series of increasingly complicated localization functors L_n for $n \geq 0$, which detect “ v_n -periodic” phenomena.

- L_0 is rationalization. Rational stable homotopy theory is very well understood. It detects only the 0-stem in the stable homotopy groups of spheres.
- L_1 is localization with respect to K -theory. It detects the image of J and the α family in the stable homotopy groups of spheres. The Lichtenbaum-Quillen conjecture is a statement about L_1 of algebraic K -theory.
- L_2 is equivalent to localization with respect to elliptic cohomology as defined above. It detects the β family in the stable homotopy groups of spheres. Davis’ nonimmersion theorem for real projective spaces was proved using related methods. The theory of topological modular forms of Hopkins *et al* is a refinement of elliptic cohomology.
- For $n > 2$ there is no comparable geometric definition of L_n , which can only be constructed by less illuminating algebraic methods related to BP -theory. It detects higher Greek letter families in the stable homotopy groups of spheres. The n th Morava K -theory is closely related to it.

THE HEIGHT OF A FORMAL GROUP LAW

Definition 1. Let F be 1-dimensional formal group law. For a positive integer m , the m -series is defined inductively by

$$[m]_F(x) = F(x, [m-1]_F(x))$$

where $[1]_F(x) = x$. Over a field k of characteristic p , the p -series is either 0 or has the form

$$[p]_F(x) = ax^{p^n} + \dots$$

for some nonzero $a \in k$. The **height** of F is the integer n . If $[p]_F(x) = 0$ (which happens when $F(x, y) = x + y$), the height is defined to be ∞ .

EXAMPLES OF m -SERIES

- For the additive formal group law, $[m](x) = mx$.
- For the multiplicative formal group law,

$$\begin{aligned} [m](x) &= (1+x)^m - 1 \\ &= mx + \binom{m}{2}x^2 + \dots \end{aligned}$$

- For $F(x, y) = \frac{x+y}{1-xy}$, we have

$$[m](x) = \frac{\sum_i \binom{m}{2i+1} x^{2i+1}}{\sum_i (-1)^i \binom{m}{2i} x^{2i}}$$

EXAMPLES OF HEIGHTS

- The multiplicative formal group law (which is associated with K -theory) has height 1.
- The formal group law associated with an elliptic curve is known to have height at most 2.
- v_n -periodic phenomena (the n th layer in the chromatic tower) are related to formal group laws of height n .

QUESTION How can we attach formal group laws of height > 2 to geometric objects (such as algebraic curves) and use them get insight into cohomology theories that go deeper into the chromatic tower?

PROGRAM:

- Let C be a curve of genus g over some ring R .
- Its Jacobian $J(C)$ is an abelian variety of dimension g .
- $J(C)$ has a formal completion $\widehat{J}(C)$ which is a g -dimensional formal group law.
- If $\widehat{J}(C)$ has a 1-dimensional summand, then Quillen's theorem gives us a genus (in the sense of the talk or 2/27) associated with the curve C .

CAVEAT

Note that a 1-dimensional summand of the formal completion $\widehat{J}(C)$ is *not* the same thing as 1-dimensional factor of the Jacobian $J(C)$. The latter would be an elliptic curve, whose formal completion can have height at most 2. There is a theorem that says if an abelian variety A has a 1-dimensional formal summand of height n for $n > 2$, then the dimension of A (and the genus of the curve, if A is a Jacobian) is at least n .

ARTIN-SCHREIER CURVES

Theorem 2. Let $C(p, f)$ be the curve over \mathbf{F}_p defined by the affine equation

$$y^e = x^p - x \quad \text{where } e = p^f - 1.$$

(Assume that $f > 1$ when $p = 2$.) Then its Jacobian has a 1-dimensional formal summand of height $(p-1)f$.

The resulting genus is *not* Landweber exact, so this does not lead to a cohomology theory.

THE LUBIN-TATE LIFTING

Theorem 3. Let $C'(p, f)$ be the curve over the ring $E_* = \mathbf{Z}_p[[u_1, \dots, u_{h-1}]][[u, u^{-1}]]$ defined by

$$y^e = x^p - u^m x + \sum_{i=0}^{h-2} u_{i+1} x^{p-1-[i/f]} y^{p^{f-1}-p^{i-[i/f]f}}$$

where $m = (p-1)e$.

Then its Jacobian has a formal 1-dimensional subgroup isomorphic to the Lubin-Tate [LT65] lifting of the formal group law above. The resulting genus satisfies Landweber's exactness criteria, so we get a cohomology theory.

MY DREAM CURVE

Conjecture 4. Let $C''(p, f)$ be the curve over the ring

$$R = \mathbf{Z}_p[[u, u^{-1}]][[a_{(p-s)e-qt} : s, t \geq 0, es + pt < pe]]$$

defined by

$$y^e = x^p - u^m x + \sum_{s,t} a_{(p-s)e-pt} x^s y^t$$

where $|u| = 2$ and $|a_i| = 2i$.

Then its Jacobian has a formal 1-dimensional subgroup, subject to certain divisibility conditions among the a_i for $f > 2$. The resulting genus also satisfies Landweber's exactness criteria.

This last curve is amenable to change of coordinates and possibly to a calculation generalizing that of Hopkins-Mahowald for tmf .

PROPERTIES OF $C(p, f)$

Recall that $C(p, f)$ is the Artin-Schreier curve over \mathbf{F}_p defined by the affine equation

$$y^e = x^p - x, \quad \text{where } e = p^f - 1,$$

and its Jacobian has a 1-dimensional formal summand of height $h = (p - 1)f$.

- Its genus is $(p - 1)(e - 1)/2$. (Thus it is zero in the excluded case $(p, f) = (2, 1)$.)
- It has an action by the group

$$G = \mathbf{F}_p \mu_m \quad \text{where } m = (p - 1)e$$

and μ_m is the group of m th roots of unity, given by

$$(x, y) \mapsto (\zeta^e x + a, \zeta y)$$

for $a \in \mathbf{F}_p$ and $\zeta \in \mu_m$. This group is a maximal finite subgroup of the h th Morava stabilizer group, and it acts appropriately on the 1-dimensional formal summand.

- For $f = 1$ (and $p > 2$) Theorem 3 was proved by Gorbunov-Mahowald [GM00].

EXAMPLES OF THESE CURVES

- $C(2, 2)$ and $C(3, 1)$ are elliptic curves whose formal group laws have height 2.
- $C(2, 3)$ has genus 3 and a 1-dimensional formal summand of height 3.
- $C(2, 4)$ and $C(3, 2)$ each has genus 7 and a 1-dimensional formal summand of height 4.

REMARKS

- Theorem 2 was known to and cited by Manin in 1963 [Man63]. Most of what is needed for the proof can be found in Katz's 1979 Bombay Colloquium paper [Kat81] and in Koblitz' Hanoi notes [Kob80].
- The original proof rests on the determination of the zeta function of the curve by Davenport-Hasse in 1934 [HD34], and on some properties of Gauss sums proved by Stickelberger in 1890 [Sti90]. The method leads to complete determination of $\widehat{J}(C(p, f))$.
- We have reproved Theorem 2 using Honda's theory of commutative formal group laws developed in the early '70s. This proof does not rely on knowledge of the zeta function and can be modified to prove Theorem 3 and presumably Conjecture 4.

HONDA THEORY:

- Given a power series $f(x_1, x_2, \dots)$ in several variables over \mathbf{Z}_p or \mathbf{Q}_p , let $T^n(f)$ be the power series obtained from f by replacing each variable by its p^n th power. This leads to an action of the ring $\mathbf{Z}_p[T]$ on the power series ring $R = \mathbf{Q}_p[[x_1, x_2, \dots]]$. Similarly a vector of d such power series admits an action by the matrix ring $M_d(\mathbf{Z}_p[T])$.
- Suppose we have a d -dimensional formal group law F over \mathbf{Z}_p . F is characterized by its logarithm f , which is a vector of d power series in d variables over the field \mathbf{Q}_p . Given a matrix $H = \sum_i C_i T^i$ with $C_i \in M_d(\mathbf{Z}_p)$, define

$$(H * f)(x_1, \dots, x_d) = \sum_i C_i f(x_1^{p^i}, \dots, x_d^{p^i}).$$

Definition 5. We say that H is a **Honda matrix** for F (or for the vector f) and that F is of type H , if $H \equiv pI_d$ modulo T (I_d is the $d \times d$ identity matrix) and $(H * f)(x) \equiv 0$ modulo (p) . Two such matrices are said to be **equivalent** if they differ by unit multiplication on the left.

Theorem 6 (Honda, 1970 [Hon70]). *The strict isomorphism classes of d -dimensional formal group laws over \mathbf{Z}_p correspond bijectively to the equivalence classes of matrices*

$$H \in M_d(\mathbf{Z}_p)\langle\langle T \rangle\rangle$$

congruent to pI_d modulo degree 1. H and f are related by the formula

$$f(x) = (H^{-1} * p)(x).$$

EXAMPLES OF HONDA MATRICES

- For $d = 1$, let H be the 1×1 matrix with entry $h = p - T^n$ for a positive integer n . Then

$$f(x) = \sum_{i \geq 0} \frac{x^{p^{ni}}}{p^i}$$

and F is the formal group law for the Morava K-theory $K(n)_*$.

- Let $A = \mathbf{Z}_p[[u_1, u_2, \dots, u_{n-1}]]$ for a positive integer m , and let $Tu_i = u_i^p$. Let H be the 1×1 matrix with entry

$$h = p - T^n - \sum_{0 < i < n} u_i T^i.$$

Then $f(x)$ is the logarithm for the Lubin-Tate lifting of the formal group law above.

HONDA THEORY FOR CURVES

Question: How can we find the Honda matrix for the formal completion of the Jacobian of an algebraic curve?

Theorem 7 (Honda, 1973 [Hon73]). *Let C be a curve of genus g over \mathbf{Z}_p with smooth reduction modulo p , let*

$$\{\omega_1, \dots, \omega_g\}$$

be a basis for the space of holomorphic 1-forms of C written as power series in a local parameter y , and let

$$\psi_i = \int_0^y \omega_i.$$

Then if H is a Honda matrix for the vector (ψ_1, \dots, ψ_g) , it is also one for $\widehat{J}(C)$, the formal completion of the Jacobian $J(C)$.

Note that ψ above is a vector of power series in one variable over \mathbf{Q}_p , while the logarithm of $\widehat{J}(C)$ is a vector of power series in g variables. The theorem asserts that they have the same Honda matrix.

HONDA THEORY FOR $C(p, f)$

Recall that our curve $C(p, f)$ is defined by the affine equation

$$y^e = x^p - x \quad \text{where } e = p^f - 1.$$

Its genus is $g = (e-1)(p-1)/2$. A basis for the holomorphic 1-forms for $C(p, f)$ is

$$\{\omega_{j,k} : j, k \geq 0, ej + pk < 2g - 1\},$$

where

$$\omega_{j,k} = \frac{x^j y^k dy}{1 - px^{p-1}}.$$

We denote the integral of its expansion in terms of y by ψ_{ej+k+1} , and we have

$$\psi_{ej+k+1} = \sum_{i \geq 0} \binom{pi+j}{i} \frac{y^{mi+ej+k+1}}{mi+ej+k+1}.$$

This enables us to prove Theorem 2.

THE HONDA MATRIX FOR $C(2, 3)$

For $C(2, 3)$ (where $g = 3$ and $m = 7$), the integrals have the form

$$\begin{aligned}\psi_1 &\in y\mathbf{Q}_2[[y^7]] \\ \psi_2 &\in y^2\mathbf{Q}_2[[y^7]] \\ \psi_3 &\in y^3\mathbf{Q}_2[[y^7]]\end{aligned}$$

More explicitly

$$\psi_k = \sum_{i \geq 0} \binom{2i}{i} \frac{y^{7i+k}}{7i+k}.$$

This means that

$$\begin{aligned}T\psi_1 &\in y^2\mathbf{Q}_2[[y^7]] \\ T\psi_2 &\in y^4\mathbf{Q}_2[[y^7]] \\ T\psi_3 &\in y^6\mathbf{Q}_2[[y^7]] \\ T^2\psi_1 &\in y^4\mathbf{Q}_2[[y^7]] \\ T^2\psi_2 &\in y^8\mathbf{Q}_2[[y^7]] \subset y\mathbf{Q}_2[[y^7]] \\ T^2\psi_3 &\in y^{12}\mathbf{Q}_2[[y^7]] \subset y^5\mathbf{Q}_2[[y^7]]\end{aligned}$$

This implies that the Honda matrix has the form

$$H = \begin{bmatrix} h_{1,1}(T^3) & T^2h_{1,2}(T^3) & 0 \\ Th_{2,1}(T^3) & h_{2,2}(T^3) & 0 \\ 0 & 0 & h_{3,3}(T^3) \end{bmatrix}$$

where

$$h_{i,j}(T^3) = \sum_{k \geq 0} h_{i,j,k} T^{3k}$$

with $h_{i,j,k} \in \mathbf{Z}_{(2)}$ and $h_{i,i,0} = 2$.

This means that the 3-dimensional formal group law has a 1-dimensional summand. Since

$$\begin{aligned}\psi_3 &= \frac{y^3}{3} + \frac{y^{10}}{5} + \frac{6y^{17}}{17} + \frac{5y^{24}}{6} + \dots \\ &\equiv y^3 + y^{10} + \frac{y^{24}}{2} + \dots \pmod{2},\end{aligned}$$

$h_{3,3}$ is roughly $2 - T^3$, and the 1-dimensional summand has height 3.

THE HONDA MATRIX FOR $C(3, 2)$

For $C(3, 2)$ (where $g = 7$ and $m = 16$), we get integrals ψ_k for

$$k \in S = \{1, 2, 3, 4, 5, 9, 10\}.$$

Explicitly,

$$\psi_k = \sum_{i \geq 0} \binom{3i + \lfloor k/8 \rfloor}{i} \frac{y^{16i+k}}{16i+k}.$$

A similar computation shows that ψ_5 corresponds to a 1-dimensional formal summand. The argument boils down to seeing how the orbits O of $\mathbf{Z}/(16)$ under multiplication by 3 intersect the set S above. Each such intersection corresponds to a formal summand whose dimension is the cardinality of $O \cap S$ and whose height is the cardinality of O . One such orbit is $\{5, 15, 13, 7\}$, whose intersection with S is the singleton $\{5\}$.

We find that

$$\psi_5 \equiv -y^5 + y^{21} - y^{117} + y^{261} + \frac{2y^{405}}{3} + \dots \pmod{3},$$

which leads to a Honda eigenvalue of roughly $3 - T^4$, so the height of the 1-dimensional formal summand is 4 as claimed.

REFERENCES

- [GM00] V. Gorbounov and M. Mahowald. Formal completion of the Jacobians of plane curves and higher real K -theories. *J. Pure Appl. Algebra*, 145(3):293–308, 2000.
- [HD34] H. Hasse and H. Davenport. Die Nullstellensatz der Kongruenz zeta-funktionen in gewissen zyklischen Fällen. *J. Reine Angew. Math.*, 172:151–182, 1934.
- [HM] M. J. Hopkins and M. A. Mahowald. From elliptic curves to homotopy theory. Preprint in Hopf archive.
- [Hon70] Taira Honda. On the theory of commutative formal groups. *J. Math. Soc. Japan*, 22:213–246, 1970.
- [Hon73] Taira Honda. On the formal structure of the Jacobian variety of the Fermat curve over a p -adic integer ring. In *Symposia Mathematica, Vol. XI (Convegno di Geometria, IN-DAM, Rome, 1972)*, pages 271–284. Academic Press, London, 1973.
- [Kat81] Nicholas M. Katz. Crystalline cohomology, Dieudonné modules, and Jacobi sums. In *Automorphic forms, representation theory and arithmetic (Bombay, 1979)*, volume 10 of *Tata Inst. Fund. Res. Studies in Math.*, pages 165–246. Tata Inst. Fundamental Res., Bombay, 1981.
- [Kob80] Neal Koblitz. *p -adic analysis: a short course on recent work*, volume 46 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 1980.
- [Lan76] P. S. Landweber. Homological properties of comodules over $MU_*(MU)$ and $BP_*(BP)$. *American Journal of Mathematics*, 98:591–610, 1976.
- [LRS95] P. S. Landweber, D. C. Ravenel, and R. E. Stong. Periodic cohomology theories defined by elliptic curves. In Mila Cenk and Haynes Miller, editors, *The Čech Centennial*, volume 181 of *Contemporary Mathematics*, pages 317–338, Providence, Rhode Island, 1995. American Mathematical Society.
- [LT65] J. Lubin and J. Tate. Formal complex multiplication in local fields. *Annals of Mathematics*, 81:380–387, 1965.
- [Man63] Ju. I. Manin. Theory of commutative formal groups over fields of finite characteristic. *Uspehi Mat. Nauk*, 18(6 (114)):3–90, 1963.
- [Qui69] D. G. Quillen. On the formal group laws of oriented and unoriented cobordism theory. *Bulletin of the American Mathematical Society*, 75:1293–1298, 1969.
- [Sti90] L. Stickelberger. Über eine Verallgemeinerung der Kreistheilung. *Mathematische Annalen*, 37:321–367, 1890.