

# On the Non-Existence of Kervaire Invariant One Manifolds

M. Hill<sup>1</sup>, M. Hopkins<sup>2</sup>, D. Ravenel<sup>3</sup>

<sup>1</sup>University of Virginia

<sup>2</sup>Harvard University

<sup>3</sup>University of Rochester

Isle of Skye, June 2009

# Main Result

## Theorem (H.-Hopkins-Ravenel)

*There are smooth Kervaire invariant one manifolds only in dimensions 2, 6, 14, 30, 62, and maybe 126.*

Exemplars:

- 2  $S^1 \times S^1$
- 6  $SU(2) \times SU(2)$
- 14  $S(\mathbb{O}) \times S(\mathbb{O})$
- 30 (Bökstedt) Related to  $E_6/(U(1) \times Spin(10))$
- 62 Possibly a similar construction.

# Geometry and History

1930s Pontryagin proves

$$\{\text{framed } n - \text{manifolds}\} / \text{cobordism} \cong \pi_n^S.$$

Tries to use surgery to reduce to spheres & misses an obstruction.

1950s Kervaire-Milnor show can always reduce to case of spheres

Except possibly in dimension  $4k + 2$ , where there is an obstruction: Kervaire Invariant.

# Adams Spectral Sequence

$$[X, Y] \rightsquigarrow \text{Hom}_{\mathcal{A}}(H^*(Y), H^*(X))$$

Have a SS with

$$E_2 = \text{Ext}_{\mathcal{A}}(H^*(Y), H^*(X))$$

and converging to  $[X, Y]$ .

- (Adem)  $\text{Ext}^1(\mathbb{F}_2, \mathbb{F}_2)$  is generated by classes  $h_i$ ,  $i \geq 0$ .
- $h_j$  survives the Adams SS if  $\mathbb{R}^{2^j}$  admits a division algebra structure.

# Browder's Reformulation

## Theorem (Browder 1969)

- 1 *There are no smooth Kervaire invariant one manifolds in dimensions not of the form  $2^{j+1} - 2$ .*
- 2 *There is such a manifold in dimension  $2^{j+1} - 2$  iff  $h_j^2$  survives the Adams spectral sequence.*

Adams showed that  $h_j$  itself survives only if  $j < 4$

$$d_2(h_{j+1}) = h_0 h_j^2.$$

## Previous Progress

$h_1^2$ ,  $h_2^2$ , and  $h_3^2$  classically exist.

Theorem (Mahowald-Tangora)

*The class  $h_4^2$  survives the Adams SS.*

Theorem (Barratt-Jones-Mahowald)

*The class  $h_5^2$  survives the Adams SS.*

Theorem (H.-Hopkins-Ravenel)


*For  $j \geq 7$ ,  $h_j^2$  does not survive the Adams SS.*

# General Outline

There are four main steps

- 1 Reduce to a simpler homotopy computation which faithfully sees the Kervaire classes
- 2 Rigidify the problem to get more structure and less wiggle-room
- 3 Show homotopy is automatically zero in dimension  $-2$
- 4 Show homotopy is periodic with period  $2^8$

# Reduction to Simpler Cases





Reduction is purely algebraic!

- 1 Lifting from Adams to Adams-Novikov is well understood.
- 2 Reduction from Adams-Novikov to homotopy fixed points is formal deformation theory.

So good choice of  $R$  gives us something that is

- easily computable
- strong enough to detect the classes.

## Why Go Equivariant?

- Homotopy fixed point spectral sequence is still too complicated.
- Simplify computation by adding extra structure: equivariance.
- Here have fixed points, rather than homotopy fixed points.
- And there are spheres for every real representation.

### Example

*If  $G = \mathbb{Z}/2$ , then have  $S^{p_2} = \mathbb{C}^+$  and  $S^2$ .*

# Important Representations

Focus now on  $G = \mathbb{Z}/8$ .

$RO(\mathbb{Z}/8)$  is rank 5 over  $\mathbb{Z}$ , generated by 1-dim reps:

- trivial rep  $1$
- sign rep  $\sigma$

and 2-dim reps:  $L = \mathbb{C}, L^2, L^3$ .

We care only about  $\rho_8 = 1 \oplus \sigma \oplus L \oplus L^2 \oplus L^3$ . Plus the regular reps for subgroups.


# What is $R$ ?

- 1 Begin with  $MU$  with  $\mathbb{Z}/2$  given by complex conjugation.
- 2 “induce” up to a  $\mathbb{Z}/8$  spectrum:


$$\begin{array}{c} \overline{(-)} \\ \downarrow \qquad \qquad \qquad \uparrow \\ MU \otimes MU \otimes MU \otimes MU \\ \uparrow \qquad \uparrow \qquad \uparrow \qquad \uparrow \end{array}$$

- 3 The “fixed points” for the  $\mathbb{Z}/8$ -action is geometric.
- 4 Inverting an equivariant class  $\Delta$  makes the fixed points and homotopy fixed points agree.

# Advantages of the Slice SS



# Advantages of the Slice SS



# Basic Idea of Slices

Want to decompose  $X$  into computable pieces.

Similar to Postnikov tower.

Key difference: **don't use all spheres!**

## Acceptable Ones

- 1  $S^{k\rho_8}, S^{k\rho_8-1}$
- 2  $\mathbb{Z}/8 \otimes_{\mathbb{Z}/4} S^{k\rho_4}$
- 3  $\mathbb{Z}/8 \otimes_{\mathbb{Z}/2} S^{k\rho_2}$
- 4  $\mathbb{Z}/8 \otimes S^k$

## Unacceptable Ones


- 1  $S^{k\rho_8-2}$
- 2  $\mathbb{Z}/8 \otimes_{\mathbb{Z}/4} S^\sigma$
- 3  $\mathbb{Z}/8 \otimes_{\mathbb{Z}/2} S^{\sigma-1}$
- 4  $S^k$

# Computing with Slices

## Key Fact

For spectra like  $MU$ , slices can be computed from equivariant simple chain complexes.

These algebraically describe the fixed points of the acceptable spheres.



## Cellular Chains for $S^{p_4-1}$

Gives the chain complex

$$\mathbb{Z}^4 \rightarrow \mathbb{Z}^4 \rightarrow \mathbb{Z}^2 \rightarrow \mathbb{Z} = C_{\bullet}.$$

Maps determined by

$$H_*(C_{\bullet}) = H_*(S^3).$$



# Gaps

## Theorem

*For any non-trivial subgroup  $H$  of  $\mathbb{Z}/8$  and for any slice sphere  $\mathbb{Z}/8 \otimes_H S^{\rho_H}$ ,*

$$H_{-2}(C_*^{\mathbb{Z}/8}) = 0$$

The proof is an easy direct computation:

- 1 If  $k \geq 0$ , then we are looking at something connected.
- 2 If  $k \leq 0$ , then we look at the associated cochain algebra.
- 3 In the relevant degrees, the complex is  $\mathbb{Z} \rightarrow \mathbb{Z}^2$  by  $1 \mapsto (1, 1)$ .

# Gap Theorem

## Theorem

$$\pi_{-2}(R) = 0.$$

## Proof.

- Slices of  $MU \otimes MU \otimes MU \otimes MU$  are all of the form

$$H\mathbb{Z} \otimes (\mathbb{Z}/8 \otimes_H S^{k\rho_H}).$$

- Class we are inverting is carried by an  $S^{k\rho_8}$ .
- Inversion is a colimit and first steps show  $\pi_{-2} = 0$ . □

## Take Home Message

Happy  $A_5$  Birthday,  
Bob and Ron!