

THE STRUCTURE OF THE COHOMOLOGY OF MORAVA STABILIZER ALGEBRA $S(3)$

ATSUSHI YAMAGUCHI

(Received April 26, 1991)

(Revised June 28, 1991)

Introduction.

Let X be a space and let p be a prime number. The E_2 -term of the Adams-Novikov spectral sequence associated with BP -theory at p converging to the p -localized homotopy group of X is given by $\text{Ext}_{BP_*BP}(BP_*, BP_*(X))$ ([1], [4]). This motivates to study $\text{Ext}_{BP_*BP}(BP_*, M)$ for a BP_*BP -comodule M . If X is a finite complex, $BP_*(X)$ is a finitely presented BP_* -module ([3]). Recall ([1], [14]) that $BP_* = \mathbf{Z}_{(p)}[v_1, v_2, \dots]$, $\deg v_n = 2(p^n - 1)$, and I_n denotes an invariant prime ideal $(p, v_1, v_2, \dots, v_{n-1})$ of BP_* . Landweber proved the following theorem.

Theorem ([6]). *Let M be a BP_*BP -comodule which is finitely presented as a BP_* -module. Then, M has a finite filtration by BP_*BP -subcomodules $0 = M_0 \subset M_1 \subset \dots \subset M_k = M$ such that for $1 \leq i \leq k$, M_i/M_{i-1} is isomorphic to BP_*/I_{n_i} for some $n_i \geq 0$ as a BP_*BP -comodule up to shifting degrees.*

By virtue of the above and a spectral sequence $E_2^{s,t} = \text{Ext}_{BP_*BP}^{s+t}(BP_*, M_i/M_{i-1}) \Rightarrow \text{Ext}_{BP_*BP}^{s+t}(BP_*, M)$ (See section 2) for M as above, we can relate $\text{Ext}_{BP_*BP}(BP_*, BP_*/I_n)$ ($n=0, 1, 2, \dots$) with $\text{Ext}_{BP_*BP}(BP_*, M)$. Hence it is necessary to know $\text{Ext}_{BP_*BP}(BP_*, BP_*/I_n)$ before we study the general case.

For small n , $\text{Ext}_{BP_*BP}(BP_*, BP/I_n)$ also has a geometric significance since there is a spectrum $V(n)$ whose BP -homology is isomorphic to BP_*/I_{n+1} , generalizing the Moore spectrum, if $p > 2n$ and $n=0, 1, 2, 3$ ([2], [15]). Hence $\text{Ext}_{BP_*BP}(BP_*, BP_*/I_{n+1})$ is the E_2 -term of the Adams-Novikov spectral sequence converging to the homotopy group of $V(n)$. We note that $V(n)$ is a ring spectrum if $p > 2n+2$ ([15]).

Since multiplication by v_n on BP_*/I_n is a BP_*BP -comodule homomorphism, $v_n^{-1}BP_*/I_n$ is a BP_*BP -comodule and $\text{Ext}_{BP_*BP}(BP_*, BP_*/I_n)$ is a module over $F_p[v_n]$ if $n > 0$. In fact, $\text{Ext}_{BP_*BP}(BP_*, BP_*/I_n)$ is a graded commutative algebra and $\text{Ext}_{BP_*BP}^0(BP_*, BP_*/I_n)$ is isomorphic to $F_p[v_n]$ if $n > 0$ ([5], [11]). Thus $v_n^{-1}\text{Ext}_{BP_*BP}(BP_*, BP/I_n)$ makes sense and it is obviously isomorphic to $\text{Ext}_{BP_*BP}(BP_*, v_n^{-1}BP_*/I_n)$.

We put $K(n)_* = \mathbf{F}_p[v_n, v_n^{-1}]$ and $\Sigma(n) = K(n)_* \otimes_{BP_*} BP_*BP \otimes_{BP_*} K(n)_*$, then $\Sigma(n)$ is a Hopf algebra over $K(n)_*$. It is shown in [9] that $\text{Ext}_{BP_*BP}(BP_*, v_n^{-1}BP_*/I_n)$ is isomorphic to $\text{Ext}_{\Sigma(n)}(K(n)_*, K(n)_*)$. Regarding \mathbf{F}_p as a $K(n)_*$ -algebra by $\rho: K(n)_* \rightarrow \mathbf{F}_p, \rho(v_n) = 1$, we put $S(n) = \Sigma(n) \otimes_{K(n)_*} \mathbf{F}_p$. $S(n)$ is a $\mathbf{Z}/2(p^n - 1)$ -graded Hopf algebra over \mathbf{F}_p and the dual Hopf algebra of $S(n)$ is called Morava stabilizer algebra. A functor from the category of graded $\Sigma(n)$ -comodules to the category of $\mathbf{Z}/2(p^n - 1)$ -graded $S(n)$ -comodules which assigns M to $M \otimes_{K(n)_*} \mathbf{F}_p$ is an equivalence of these categories. Thus $\text{Ext}_{\Sigma(n)}(K(n)_*, K(n)_*) \otimes_{K(n)_*} \mathbf{F}_p$ is isomorphic to $\text{Ext}_{S(n)}(\mathbf{F}_p, \mathbf{F}_p)$ and $\text{Ext}_{\Sigma(n)}(K(n)_*, K(n)_*)$ can be recovered from $\text{Ext}_{S(n)}(\mathbf{F}_p, \mathbf{F}_p)$. Therefore, the v_n -torsion free part of $\text{Ext}_{BP_*BP}(BP_*, BP_*/I_n)$ can be detected by the cohomology of Morava stabilizer algebra $\text{Ext}_{S(n)}(\mathbf{F}_p, \mathbf{F}_p)$ by the preceding argument.

Moreover, the chromatic spectral sequence $E_1^{s,t}(n) \Rightarrow \text{Ext}_{BP_*BP}^{s,t}(BP_*, BP_*/I_n)$ is constructed in [10], having the following properties; $E_1^{0,t}(n)$ is isomorphic to $\text{Ext}_{\Sigma(n)}^t(K(n)_*, K(n)_*)$ and the edge homomorphism $\text{Ext}_{BP_*BP}^t(BP_*, BP_*/I_n) \rightarrow E_1^{0,t}(n)$ can be identified with the localization map away from v_n . There is a Bockstein long exact sequence $\cdots \rightarrow E_1^{s-1,t}(n+1) \rightarrow E_1^{s,t}(n) \xrightarrow{v_n \times} E_1^{s,t}(n) \rightarrow E_1^{s-1,t+1}(n) \rightarrow \cdots$. Hence $\text{Ext}_{\Sigma(m)}(K(m)_*, K(m)_*)$ for $m = n, n+1, \dots$ relate with $\text{Ext}_{BP_*BP}(BP_*, BP_*/I_n)$ through the Bockstein spectral sequences and the chromatic spectral sequence.

The cohomology of Morava stabilizer algebra $\text{Ext}_{S(n)}(\mathbf{F}_p, \mathbf{F}_p)$ is calculated for $n = 1, 2$ in [13] and the Poincare series of $\text{Ext}_{S(3)}(\mathbf{F}_p, \mathbf{F}_p)$ is also given for $p \geq 5$. In this paper we determine the algebra structure of $\text{Ext}_{S(3)}(\mathbf{F}_p, \mathbf{F}_p)$ for $p \geq 5$. By the above explanation, our result is a part of the initial input for the chromatic spectral sequences and it also gives the v_3 -localization of the E_2 -term of the Adams-Novikov spectral sequence converging to $\pi_*(V(2))$. Since $v_3 \in E_2^{0,2(p^3-1)}$ is known to be a permanent cycle and $V(2)$ is a ring spectrum if $p \geq 7$, $\pi_*(V(2))$ is a module over $\mathbf{F}_p[v_3]$. Thus our result is expected to give some information on the v_3 -torsion free part of $\pi_*(V(2))$.

For the calculation, we apply the method of May and Ravenel ([13]), namely, define a certain filtration on $S(3)$ such that the dual of the associated graded Hopf algebra is primitively generated. By the theorem of Milnor-Moore [8], $(E_0S(3))^*$ is isomorphic to the universal enveloping algebra of a restricted Lie algebra $L(3) = P(E_0(S(3)))^*$. Let $L^u(3)$ denote the unrestricted Lie algebra obtained by forgetting the restriction of $L(3)$. We use the following spectral sequences which we review in section 2; $E_2^{s,t} = \text{Ext}_{E^0S(3)}^{s,t}(\mathbf{F}_p, \mathbf{F}_p)_t \Rightarrow \text{Ext}_{S^+(3)}^{s,t}(\mathbf{F}_p, \mathbf{F}_p)$ and $E_2^{s,t} = \text{Ext}_{U(L^u(3))}^s(\mathbf{F}_p, \mathbf{F}_p) \otimes P(s^2\pi L(3)^*)^t \Rightarrow \text{Ext}_{V(L(3))}^{s,t}(\mathbf{F}_p, \mathbf{F}_p) = \text{Ext}_{E^0S(3)}^{s,t}(\mathbf{F}_p, \mathbf{F}_p)$. The unrestricted Lie algebra $L^u(3)$ turns out to be a product of a nine-dimensional Lie algebra $M(3)$ and an abelian Lie algebra $I(3)$, and the edge homomorphism of the latter spectral sequence gives an isomorphism $\text{Ext}_{E^0S(3)}^t(\mathbf{F}_p, \mathbf{F}_p) \rightarrow E_\infty^{0,t} = E_3^{0,t} = \text{Ext}_{U(M(3))}^t(\mathbf{F}_p, \mathbf{F}_p)$. After we calculate the cohomology of $M(3)$, by show-

ing that the former spectral sequence collapses and the extension is trivial, we prove that the cohomology of $S(3)$ is isomorphic to that of $M(3)$.

In section 1, we review how to construct an economical resolution for the universal enveloping algebra of a restricted Lie algebra according to [4], [7]. We also summarize a part of Ravenel's work [13] needed for our calculation. In section 2, we set up two kinds of spectral sequences we mentioned above. Sections 3 and 4 are devoted to calculate the cohomology a certain nine-dimensional Lie algebra $M(3)$, applying a sort of Cartan-Eilenberg spectral sequence. In section 5, we show that the cohomology of $S(3)$ is isomorphic to that of $M(3)$.

1. Recollections

First, we recall from [4] and [7] how to construct economical resolutions.

NOTATIONS. For a graded vector space V and an integer l , we denote by $s^l V$ a bigraded vector space given by $(s^l V)_{i,j} = 0$ if $i \neq l$, $(s^l V)_{i,j} = V_j$ and πV denotes a graded vector space given by $(\pi V)_i = 0$ if p does not divide i , $(\pi V)_{p,j} = V_j$. We denote by x and \bar{x} the elements of $(sV)_{1,j}$ and $(s^2\pi V)_{2,pj}$ corresponding to an element x of V_j . We also denote by $E(V)$, $P(V)$ and $\Gamma(V)$ the exterior algebra, the polynomial algebra and the divided polynomial algebra generated by V , respectively. For an element x of V , let $\langle x \rangle$ and $\gamma_i(x)$ be typical generators of $E(V)$ and $\Gamma(V)$, respectively. For a bigraded vector space W and $l \in \mathbf{Z}$, we put $W_l = \sum_j W_{l,j}$.

Let K be a field of characteristic $p \neq 0$, and let $L = \sum_{i \geq 0} L_i$ be a graded restricted Lie algebra over K with restriction ξ . We denote by L^u the unrestricted Lie algebra obtained from L by forgetting the restriction of L . We put $L^+ = \sum_{i \geq 0} L_{2i}$, $L^- = \sum_{i \geq 0} L_{2i+1}$ if $p > 2$, and $L^+ = L$, $L^- = 0$ if $p = 2$. Let us denote by $U(L^u)$ and $V(L)$ the universal enveloping algebras of an unrestricted Lie algebra L^u and a restricted Lie algebra L , respectively. J.P. May ([7], see also [4]) constructed a $U(L^u)$ -free resolution $Y(L^u)$ of K as follows; $Y(L^u) = U(L^u) \otimes E(sL^+) \otimes \Gamma(sL^-)$ as a left $U(L^u)$ -module. Give $Y(L^u)$ a K -algebra structure such that the canonical inclusions of $U(L^u)$, $E(sL^+)$ and $\Gamma(sL^-)$ into each factor of $Y(L^u)$ are monomorphism of K -algebras and that the following relations hold for $x, x_i \in L^-$, $y, y_i \in L^+ (i=1, 2)$; $\langle \bar{y}_1 \rangle y_2 = y_2 \langle \bar{y}_1 \rangle + \langle \overline{[y_1, y_2]} \rangle$, $\langle \bar{y} \rangle x = -x \langle \bar{y} \rangle + \gamma_1(\overline{[y, x]})$, $\gamma_t(\bar{x})y = y\gamma_t(\bar{x}) + \gamma_t(\overline{[x, y]})\gamma_{t-1}(\bar{x})$, $\gamma_t(\bar{x}_1)x_2 = x_2\gamma_t(\bar{x}_1) + \langle \overline{[x_1, x_2]} \rangle \gamma_{t-1}(\bar{x}_1)$, $\gamma_t(\bar{x}) \langle \bar{y} \rangle = \langle \bar{y} \rangle \gamma_t(\bar{x})$. We note that $Y(L^u)$ is a bigraded algebra and, for $w \in L$, the bidegrees of $w \in U(L^u)$, $\langle \bar{w} \rangle \in E(sL^+)$ and $\gamma_t(\bar{w}) \in \Gamma(sL^-)$ are $(0, \text{deg } w)$, $(1, \text{deg } w)$ and $(t, t \text{ deg } w)$ respectively. The differential d of $Y(L^u)$ is given by $du = \varepsilon(u)$ for $u \in U(L^u)$, $d\langle \bar{y} \rangle = y$ for $y \in L^+$, $d\gamma_t(\bar{x}) = x\gamma_{t-1}(\bar{x}) + \frac{1}{2}\langle \overline{[x, x]} \rangle \gamma_{t-2}(\bar{x})$ for $x \in L^-$ satisfying the Leibniz formula $d(xy) = (dx)y + (-1)^{|x|}x dy$, where $\varepsilon: U(L^u) \rightarrow K$ is the augmentation and $|x|$ is the total degree of $x \in Y(L^u)$. We also define a coproduct $\varphi: Y(L^u) \rightarrow Y(L^u) \otimes Y(L^u)$ by $\varphi(z) = 1 \otimes z + z \otimes 1$ for $z \in L$, $\varphi(\langle \bar{y} \rangle) =$

$1 \otimes \langle \bar{y} \rangle + \langle \bar{y} \rangle \otimes 1$ for $y \in L^+$, $\varphi(\gamma_i(\bar{x})) = \sum_{i+j=t} \gamma_i(\bar{x}) \otimes \gamma_j(\bar{x})$ for $x \in L^-$ so that $Y(L^u)$ has a structure of differential Hopf algebra.

Put $W(L) = V(L) \otimes E(sL^+) \otimes \Gamma(sL^-)$ and let $q: Y(L^u) \rightarrow W(L)$ be the canonical projection induced by $U(L^u) \rightarrow V(L) = U(L^u) / (y^p - \xi(y) \mid y \in L)$, then $W(L)$ has a unique structure of differential Hopf algebra over K such that q is a morphism of differential Hopf algebras. The following is obvious.

Lemma 1.1. *q induces an isomorphism $\text{Hom}_{\mathcal{V}(L)}^*(W(L), K) \rightarrow \text{Hom}_{\mathcal{V}(L)}^*(Y(L^u), K)$ of chain complexes. Hence the cohomology of $\{\text{Hom}_{\mathcal{V}(L)}^*(W(L), K), d^*\}$ is isomorphic to $\text{Ext}_{U(L^u)}(K, K)$.*

We choose a K -basis $\{y_\alpha \mid \alpha \in \Lambda\}$ of L^+ . For each index $\alpha \in \Lambda$, let \mathbf{Z}_α^+ be a copy of a monoid of non-negative integers. For an element $R = (t_\alpha)_{\alpha \in \Lambda} \in \bigoplus_{\alpha \in \Lambda} \mathbf{Z}_\alpha^+$, we set $\gamma(R) = \prod_{\alpha \in \Lambda} \gamma_{t_\alpha}(\bar{y}_\alpha)$. Then $\{\gamma(R) \mid R \in \bigoplus_{\alpha \in \Lambda} \mathbf{Z}_\alpha^+\}$ forms a K -basis of $\Gamma(s^2\pi L^+)$. The bidegree of $\gamma(R)$ is $(2|R|, p \sum_{\alpha \in \Lambda} t_\alpha \deg y_\alpha)$, where we put $|R| = \sum_{\alpha \in \Lambda} t_\alpha$ for $R = (t_\alpha)_{\alpha \in \Lambda}$.

Lemma 1.2 ([4], [7]). 1) *There exists a twisting cochain $\theta = (\theta_{2l}, \theta_{2l}: \Gamma(s^2\pi L^+)_{2l} \rightarrow W(L)_{2l-1}$ satisfying $\theta_2(\gamma_1(\bar{y})) = y^{p-1} \langle \bar{y} \rangle - \langle \bar{\xi}(y) \rangle$ for $y \in L^+$.*
 2) *There exists a twisting diagonal cochain $\lambda = (\lambda_{2l}, \lambda_{2l}: \Gamma(s^2\pi L^+)_{2l} \rightarrow (W(L) \otimes W(L))_{2l}$ satisfying $\lambda_0(1) = 1 \otimes 1$, $\lambda_2(\gamma_1(\bar{y})) = \sum_{i=1}^{p-1} (-1)^i y^{i-1} \langle \bar{y} \rangle \otimes y^{p-1-i} \langle \bar{y} \rangle$.*

Consider a left $V(L)$ -module $X(L) = W(L) \otimes \Gamma(s^2\pi L^+)$ and define a differential d_θ and a coproduct D by $d_\theta(w \otimes \gamma(R)) = dw \otimes \gamma(R) + (-1)^{|w|} \sum_{S+T=R} w \cdot \theta(\gamma(S)) \otimes \gamma(T)$, $D(w \otimes \gamma(R)) = \phi(w) \cdot \sum_{S+T=R} \lambda(\gamma(S)) \cdot \nabla \gamma(T)$, where ϕ is the coproduct of $W(L)$ and ∇ is the standard coproduct of $\Gamma(s^2\pi L^+)$.

Theorem 1.3 ([4], [7]). *The complex $\{X(L), d_\theta\}$ is a $V(L)$ -free resolution of K . It is also a differential coalgebra with coproduct D .*

We set $\bar{Y}(L) = E(sL^+) \otimes \Gamma(sL^-)$ and define a filtration on $X(L)$ by $F_m X(L) = V(L) \otimes F_m \bar{X}(L)$ where $F_m \bar{X}(L) = \sum_{i \leq m} \bar{Y}(L)_{m-i} \otimes \Gamma(s^2\pi L^+)_i$. We state the following obvious fact for later use.

Proposition 1.4. 1) $0 = F_{-1} X(L)_i \subset W(L)_i = F_0 X(L)_i \subset \dots \subset F_i X(L)_i = X(L)_i$, $F_{2m} X(L) = F_{2m+1} X(L)$.
 2) *The inclusion $F_{m-1} X(L) \hookrightarrow F_m X(L)$ is a split monomorphism of $V(L)$ -modules.*
 3) $\{X(L), d_\theta, D\}$ is a filtered differential coalgebra and for each $w \otimes \gamma(R) \in F_{2|R|} X(L)$, $d_\theta(w \otimes \gamma(R)) \equiv dw \otimes \gamma(R)$ modulo $F_{2|R|-2} X(L)$, $D(w \otimes \gamma(R)) \equiv \phi(w) \cdot \nabla \gamma(R)$ modulo $F_{2|R|-2} X(L) \otimes X(L)$.

Next we recall some facts on Morava stabilizer algebra from [12].

Let (BP_*, BP_*BP) be the Hopf algebroid associated with BP -theory at a fixed prime p (See [14], for example). Put $K(n)_* = F_p[v_n, v_n^{-1}]$ and regard this as

a BP_* -algebra by $\pi: BP_* \rightarrow K(n)_*$, $\pi(v_i) = 0$ if $i \neq n$, $\pi(v_n) = v_n$. We set $\Sigma(n) = K(n)_* \otimes_{BP_*} BP_* BP \otimes_{BP_*} K(n)_*$, then it is known that $\Sigma(n)$ is isomorphic to $K(n)_*[t_1, t_2, \dots]/(v_n t_i^{p^n} - v_n^{p^i} t_i)$ ($\deg t_i = 2(p^i - 1)$) as a $K(n)_*$ -algebra. Note that Hopf algebroid $(K(n)_*, \Sigma(n))$ is in fact a Hopf algebra. Regarding F_p as a $K(n)_*$ -algebra by $\rho: K(n)_* \rightarrow F_p$, $\rho(v_n) = 1$, put $S(n) = \Sigma(n) \otimes_{K(n)_*} F_p$. $S(n)$ is a $\mathbb{Z}/2(p^n - 1)$ -graded Hopf algebra over F_p which is isomorphic to $F_p[t_1, t_2, \dots]/(t_i^{p^n} - t_i)$ as an F_p -algebra.

Theorem 1.5 ([12]). *Define integers d_i for $i \in \mathbb{Z}$ recursively by $d_i = 0$ if $i \leq 0$, $d_i = \max\{i, pd_{i-n}\}$ if $i > 0$. Then, there is a unique increasing Hopf algebra filtration on $S(n)$ with $t_i^{p^i} \in F_{d_i} S(n) - F_{d_i-1} S(n)$.*

Instead of considering the above filtration, we consider a new filtration $\{\widehat{F}_i S(n)\}$ defined by $\widehat{F}_{2i} S(n) = \widehat{F}_{2i+1} S(n) = F_i S(n)$ so that the associated $(\mathbb{Z} \times \mathbb{Z}/2(p^n - 1))$ -graded Hopf algebra $E^0 S(n)$ becomes graded commutative. Let $t_{i,j}$ denote the element of $E_{2d_i}^0 S(n)_{2p^j(p^i-1)}$ corresponding to $t_i^{p^j}$, where $j \in \mathbb{Z}/n$, then $E^0 S(n)$ is isomorphic to $F_p[t_{i,j} | i \geq 1, j \in \mathbb{Z}/n]/(t_{i,j}^{p^i})$ as an algebra. Consider the $(\mathbb{Z} \times \mathbb{Z}/2(p^n - 1))$ -graded dual $(E^0 S(n))^*$ of $E^0 S(n)$, and let $x_{i,j} \in (E^0 S(n))^{2d_i, 2p^j(p^i-1)}$ be the dual of $t_{i,j}$ with respect to the monomial basis. We set $L(n) = P(E^0 S(n))^*$, then $\{x_{i,j} | i \geq 1, j \in \mathbb{Z}/n\}$ spans $L(n)$. We note that this $L(n)$ is different from the one in [12], [13], [14], which coincides with an unrestricted Lie algebra $M(n)$ defined in the next section if $n < p - 1$. Since the p -th power map on $E^0 S(n)$ is trivial, it follows from [8] that $(E^0 S(n))^*$ is the universal enveloping algebra of the restricted Lie algebra $L(n)$. The bracket and the restriction are given by the following.

Theorem 1.6 ([12]). $[x_{i,j}, x_{k,l}] = \delta_{i+j}^l x_{i+k,j} - \delta_{k+l}^i x_{i+k,l}$ if $i+k \leq pn/(p-1)$, otherwise the bracket is trivial, where $\delta_i^s = 1$ if $s \equiv i \pmod n$, $\delta_i^s = 0$ otherwise. $\xi(x_{i,j}) = 0$ if $i \leq n/(p-1)$, $\xi(x_{i,j}) = -x_{i+n,j+1}$ otherwise.

REMARK 1.7. The p -th power map η on $S(n)$ is an automorphism of order n . Since η preserves the filtration, it induces an automorphism of $E^0 S(n)$ which we also denote by η . We note that η maps $t_{i,j}$ to $t_{i,j+1}$ and that η induces an automorphism η^\sharp of $L(n)$.

2. Spectral Sequences

Let R be a graded commutative ring. For a filtered R -modules M and N , we filter $M \otimes_R N$ by $F_s(M \otimes_R N) = \text{Im}(\sum_{i+j=s} F_i M \otimes_R F_j N \rightarrow M \otimes_R N)$ as usual. Then there is a natural epimorphism $\sum_{i+j=s} E_i^0 M \otimes_R E_j^0 N \rightarrow E_s^0(M \otimes_R N)$, where we put $E_i^0 M = F_i M / F_{i-1} M$. Note that this epimorphism is an isomorphism if $F_i M$ is flat over R for any i .

Let C be a filtered R -coalgebra which is flat over R , and let M be a filtered

left C -comodule. By the above remark, E^0C is an R -coalgebra and E^0M is a left E^0C -comodule. We give a decreasing filtration on the cobar complex $\Omega^*(C; M)$ ([9], [10], [14]) by $F^s\Omega^m(C; M) = \text{Im}(\sum_{i_0+\dots+i_m=m-s} F_{i_1}C \otimes_R F_{i_2}C \otimes_R \dots \otimes_R F_{i_m}C \otimes_R F_{i_0}M \rightarrow \Omega^m(C; M))$. Then the differential d of $\Omega^*(C; M)$ maps $F^s\Omega^m(C; M)$ into $F^{s+1}\Omega^{m+1}(C; M)$ and $E_0^{s,t} = F^s\Omega^{s+t}(C; M)/F^{s+1}\Omega^{s+t}(C; M)$ is isomorphic to $\Omega^{s+t}(E^0C; E^0M)_t = \sum_{i_0+\dots+i_{s+t}=t} E_{i_1}^0C \otimes_R E_{i_2}^0C \otimes_R \dots \otimes_R E_{i_{s+t}}^0C \otimes_R E_{i_0}^0M$. We rather call t of $E_0^{s,t}$ the filtration degree below. Put $D_1^{s,t} = H^{s+t}(F^s\Omega^*(C; M))$. $E_1^{s,t} = H^{s+t}(E_0^{s,*})$ and let $i_*: D_1^{s+1,t-1} \rightarrow D_1^{s,t}$ and $j_*: D_1^{s,t} \rightarrow E_1^{s,t}$ be the maps induced by inclusion $i: F^{s+1}\Omega^*(C; M) \hookrightarrow F^s\Omega^*(C; M)$ and projection $j: F^s\Omega^*(C; M) \rightarrow E_0^{s,*}$ respectively. $\partial: E_1^{s,t} \rightarrow D_1^{s+1,t}$ denotes the boundary homomorphism associated with a short exact sequence of complexes $0 \rightarrow F^{s+1}\Omega^*(C; M) \rightarrow F^s\Omega^*(C; M) \rightarrow E_0^{s,*} \rightarrow 0$. Consider the spectral sequence associated with exact couple $\langle D_1^{s,t}, E_1^{s,t}, i_*, j_*, \partial \rangle$. Then $E_1^{s,t} = E_0^{s,t}$ and the E_2 -term is given by $E_2^{s,t} = H^{s+t}(\Omega^*(E^0C; E^0M))_t = \text{Ext}_{E^0C}^{s+t}(R, E^0, M)_t$. Filter $H^*(\Omega^*(C; M)) = \text{Ext}_{\mathbb{Z}}^*(R, M)$ by putting $F^{s,t} = \text{Im}(H^{s+t}(F^s\Omega^*(C; M)) \rightarrow H^{s+t}(\Omega^*(C; M)))$. We assume that $C = \cup_s F_s C$ and $M = \cup_s F_s M$ hold and that $F_s C = F_s M = 0$ for sufficiently small s . Then the above spectral sequence converges to $\text{Ext}_{\mathbb{Z}}^*(R, M)$.

Applying the above spectral sequence to the case $R = M = F_p$, $C = S(n)$, we have a spectral sequence

$$(2.1) \quad E_2^{s,t} = \text{Ext}_{E^0S(n)}^{s+t}(F_p, F_p)_t \Rightarrow \text{Ext}_{S(n)}^{s+t}(F_p, F_p).$$

Let A be a graded algebra (not necessarily commutative) over a commutative ring. Let X be a filtered A -complex with differential $d: X_i \rightarrow X_{i-1}$. Put $E_{s,t}^0 = F_s X_{s+t}/F_{s-1} X_{s+t}$. Let M be a graded A -module. Consider a complex $\{C^*, d^*\}$ given by $C^i = \text{Hom}_A^i(X, M)$. Filter C^* by $F^s C^* = \text{Ker}(\text{Hom}_A^*(X, M) \rightarrow \text{Hom}_A^*(F_{s-1} X, M))$. We assume that the inclusions $F_s X/F_{s-1} X \hookrightarrow X/F_{s-1} X$ are split monomorphism of A -modules for any s (This holds if the inclusions $F_{s-1} X \hookrightarrow F_s X$ split for any s). Then we have short exact sequences of complexes $0 \rightarrow F^{s+1} C^* \xrightarrow{i} F^s C^* \xrightarrow{j} E_0^{s,*} \rightarrow 0$, where we set $E_0^{s,t} = \text{Hom}_A^{s+t}(E_{s,*}^0, M)$. Let $\Delta: H^*(E_0^{s,*}) \rightarrow H^{*+1}(F^{s+1} C^*)$ be the boundary homomorphism. Putting $D_1^{s,t} = H^{s+t}(F^s C^*)$, $E_1^{s,t} = H^{s+t}(E_0^{s,*})$, we consider a spectral sequence associated with an exact couple $\langle D_1^{s,t}, E_1^{s,t}, i_*, j_*, \Delta \rangle$. We define a filtration on $H^*(C^*)$ by $F^{s,t} = \text{Im}(H^{s+t}(F^s C^*) \rightarrow H^{s+t}(C^*))$. Suppose that, for each integer m , there exist integers $a(m)$ and $b(m)$ such that $F_s X_m = X_m$ if $s > a(m)$, $F_s X_m = 0$ if $s < b(m)$. Then the spectral sequence converges to $H^*(C^*)$.

By 1) and 2) of (1, 4), we can apply the above spectral sequence to the case $A = V(L)$, $X = X(L)$, $M = K$, and obtain a spectral sequence converging to $H^*(\text{Hom}_{V(L)}(X(L), K)) = \text{Ext}_{\mathbb{Z}(L)}^*(K, K)$. We note that the coproduct D of $X(L)$ makes this spectral sequence multiplicative. Identifying $E_{s,*}^0$ with $W(L) \otimes \Gamma(s^2 \pi L^+)_s$, it follows from (1, 4), 3) that d_θ and D induce $d \otimes 1: E_{s,*}^0 \rightarrow E_{s,*}^0$ and

$\phi \otimes \nabla: E_{s,*}^0 \rightarrow \sum_{i+j=s} E_{i,*}^0 \otimes E_{j,*}^0$. Therefore the E_1 -term is isomorphic to $P(s^2\pi(L^+)^*) \otimes \text{Ext}_{U(L^+)}^*(K, K)$ as an algebra by (1, 1), where $(L^+)^*$ denotes the graded K -dual of L^+ . Since $E_1^{s,t} = 0$ if s is odd or $s < 0$ or $t < 0$, $E_2^{s,t} = E_1^{s,t}$ holds and we have the edge homomorphism. Thus we have shown

Theorem 2.2 ([7]). *There is a multiplicative spectral sequence $E_2^{s,t} = P(s^2\pi(L^+)^*)^s \otimes \text{Ext}_{U(L^+)}^t(K, K) \Rightarrow \text{Ext}_{V(L)}^{s,t}(K, K)$, whose edge homomorphism $\text{Ext}_{V(L)}^t(K, K) = F^{0,t} \rightarrow E_\infty^{0,t} \hookrightarrow E_2^{0,t} = \text{Ext}_{U(L^+)}^t(K, K)$ is induced by the composite $Y(L^+) \xrightarrow{q} W(L) \hookrightarrow X(L)$.*

In particular, in the case $K = \mathbf{F}_p$, $L = L(n)$, let $M(n)$ and $I(n)$ be subspaces of $L^u(n)$ spanned by $\{x_{i,j} \mid i \leq pn/(p-1), j \in \mathbf{Z}/n\}$ and $\{x_{i,j} \mid i > pn/(p-1), j \in \mathbf{Z}/n\}$ respectively. It follows from (1, 6) that $M(n)$ is a Lie subalgebra of $L^u(n)$ and $I(n)$ is an ideal of $L^u(n)$. Obviously, $I(n)$ is an abelian Lie algebra and $L^u(n)$ is isomorphic to $M(n) \times I(n)$ as a Lie algebra. Therefore $U(L^u(n))$ is isomorphic to $U(M(n)) \otimes P(I(n))$. This implies that $\text{Ext}_{U(L^u(n))}(\mathbf{F}_p, \mathbf{F}_p)$ is isomorphic to $\text{Ext}_{U(M(n))}(\mathbf{F}_p, \mathbf{F}_p) \otimes E(\langle \bar{t}_{i,j} \rangle \mid i > pn/(p-1), j \in \mathbf{Z}/n)$ where $\text{deg} \langle \bar{t}_{i,j} \rangle = (1, 2d_i, 2p^j(p^i-1))$. Hence the E_2 -term of the spectral sequence is isomorphic to $P(\bar{t}_{i,j} \mid i \geq 1, j \in \mathbf{Z}/n) \otimes \text{Ext}_{U(M(n))}(\mathbf{F}_p, \mathbf{F}_p) \otimes E(\langle \bar{t}_{i,j} \rangle \mid i > pn/(p-1), j \in \mathbf{Z}/n)$. By (1, 2) and (1, 6), we have the following fact on the differential d_2 .

Lemma 2.3. ([13]). *$d_2 \langle \bar{t}_{i,j} \rangle = -\bar{t}_{i-n, j-1}$ for $i > pn/(p-1)$. Thus if $n < p-1$, $E_\infty^{s,t} = E_3^{s,t} = 0$ unless $s=0$, and the edge homomorphism maps $\text{Ext}_{V(L(n))}^t(\mathbf{F}_p, \mathbf{F}_p)$ bijectively onto $E_\infty^{0,t} = E_3^{0,t} = \text{Ext}_{U(M(n))}^t(\mathbf{F}_p, \mathbf{F}_p) \subset E_2^{0,t}$.*

3. Auxiliary Calculation

Let L be a graded unrestricted Lie algebra over a field K of finite type such that $L^- = 0$, and let $\{x_\lambda \mid \lambda \in \Lambda\}$ be a totally ordered basis of L . L^* denotes the graded dual of L . Take the dual basis $\{x_\lambda^* \mid \lambda \in \Lambda\}$ of $\{x_\lambda \mid \lambda \in \Lambda\}$. Define $\delta: E(sL^*)_i \rightarrow E(sL^*)_{i+1}$ by $\delta(\langle \overline{x_\lambda^*} \rangle) = -\sum_{\mu < \nu} \langle x_\mu^*, [x_\mu, x_\nu] \rangle \langle \overline{x_\mu^*} \rangle \langle \overline{x_\nu^*} \rangle$ satisfying the Leibniz formula, where $\langle, \rangle: L^* \otimes L \rightarrow K$ is the canonical pairing. It is straightforward to verify the following.

Lemma 3.1. *$\{E(sL^*), \delta\}$ is a differential algebra isomorphic to $\{\text{Hom}_{U(L)}^*(Y(L), K), d^*\}$. Hence $H^*(E(sL^*); \delta)$ is isomorphic to $\text{Ext}_{U(L)}^*(K, K)$.*

Now we concentrate on the computation of $\text{Ext}_{U(M(3))}^*(\mathbf{F}_p, \mathbf{F}_p)$ for $p \geq 5$. $M(3)$ is spanned by $\{x_{i,j} \mid i=1, 2, 3, j \in \mathbf{Z}/3\}$ over \mathbf{F}_p . Then $E(sM(3)^*) = E(t_{i,j} \mid i=1, 2, 3, j \in \mathbf{Z}/3)$ where we put $t_{i,j} = \langle \overline{x_{i,j}^*} \rangle$. It follows from (1.6) and (3.1) that δ is given by $\delta(t_{i,j}) = 0$, $\delta(t_{2,j}) = -t_{1,j}t_{1,j+1}$, $\delta(t_{3,j}) = t_{1,j-1}t_{2,j} - t_{1,j}t_{2,j+1}$ for $j \in \mathbf{Z}/3$. Let A be an ideal of $M(3)$ spanned by $\{x_{3,0}, x_{3,1}, x_{3,2}\}$ and we regard $\{E(s(M(3)/A)^*), \delta\}$ as a subcomplex of $\{E(sM(3)^*), \delta\}$. We remark that $M(3)$

and $M(3)/A$ are denoted by $L(3, 3)$ and $L(3, 2)$ respectively in [13], [14]. We can manage to compute the cohomology of $\{E(s(M(3)/A)^*), \delta\}$ directly by hand and the structure of $\text{Ext}_{U(M(3)/A)}(\mathbf{F}_p, \mathbf{F}_p)$ is described below.

For a cocycle z of $E(s(M(3)/A)^*)$, we denote by $[z]$ the cohomology class represented by z .

Lemma 3.2. 1) $\text{Ext}_{U(M(3)/A)}^*(\mathbf{F}_p, \mathbf{F}_p)$ is generated by the following seventeen elements as an algebra;

$$h_j = [t_{1,j}], g_j = [t_{1,j}t_{2,j}], g'_j = [t_{1,j+1}t_{2,j}], f_i = [t_{1,i-1}t_{2,i} - t_{1,i}t_{2,i+1}], d_j = [t_{1,j}t_{2,j-1}t_{2,j}], \\ e_j = [t_{1,j}t_{2,j}t_{2,j+1} + t_{1,j+1}t_{2,j-1}t_{2,j}], \text{ for } i, j \in \mathbf{Z}/3, i \neq 2.$$

2) $\text{Ext}_{U(M(3)/A)}^s(\mathbf{F}_p, \mathbf{F}_p) = 0$ for $s > 6$. A basis of $\text{Ext}_{U(M(3)/A)}^s(\mathbf{F}_p, \mathbf{F}_p)$ ($0 \leq s \leq 6$) is given as follows;

$$s=0; 1.$$

$$s=1; h_0, h_1, h_2.$$

$$s=2; g_0, g_1, g_2, g'_0, g'_1, g'_2, f_0, f_1.$$

$$s=3; h_1g_0, h_2g_1, h_0g_2, h_2g_0, h_0g_1, h_1g_2, d_0, d_1, d_2, e_0, e_1, e_2.$$

$$s=4; g_0g_1, g_1g_2, g_2g_0, g'_2g'_0, g'_0g'_1, g'_1g'_2, g_0g'_1, g_1g'_2.$$

$$s=5; g_1d_0, g_2d_1, g_0d_2.$$

$$s=6; g_0g_1g_2.$$

The operator η^* of $L(3)$ induces an algebra automorphism of $\text{Ext}_{U(M(3)/A)}^*(\mathbf{F}_p, \mathbf{F}_p)$ of order three, which we denote by η_* . Obviously, we have $\eta_*^i h_0 = h_i$, $\eta_*^i g_0 = g_i$, $\eta_*^i g'_0 = g'_i$, $\eta_*^i f_0 = f_i$, $\eta_*^i d_0 = d_i$, and $\eta_*^i e_0 = e_i$ for $i \in \mathbf{Z}/3$ where we put $f_2 = -f_0 - f_1$.

Lemma 3.3. Relations of $\text{Ext}_{U(M(3)/A)}^*(\mathbf{F}_p, \mathbf{F}_p)$ are given by the following and the relations obtained by applying η_* ($i=1, 2$) to them;

$$h_0h_i = 0, f_0d_i = 0, d_0d_i = d_0e_i = e_0e_i = 0 \text{ for } i \in \mathbf{Z}/3, h_0g_0 = 0, h_0g'_0 = -h_1g_0 \\ h_0g'_1 = -h_1g_2, h_0g'_2 = 0, h_0f_0 = h_0f_1 = -h_2g_0, g_0^2 = g_0'^2 = g_0g'_0 = g_0g'_2 = g_0f_0 = 0 \\ g_0f_1 = g'_2g'_0, g'_0f_0 = -g'_0f_1 = g_0g_1, g_2g'_0 = -g_0g'_1 - g_1g'_2, f_0^2 = 2g_0g'_1, f_0f_1 = 2g_2g'_0, \\ h_0d_0 = 0, h_0d_1 = -g_0g_1, h_0d_2 = g'_1g'_2, h_0e_0 = -g'_2g'_0, h_0e_1 = -g_0g'_1 + g_1g'_2, h_0e_2 = g_2g_0, \\ g_0d_0 = g_0d_1 = g_0e_0 = g_0e_2 = 0, g_0d_2 = -g_2d_0, g_0e_1 = -g_1d_0, g'_0d_0 = g'_0d_1 = g'_0e_0 = g'_0e_1 = 0, \\ g'_0e_2 = g_1d_0, f_0e_0 = g_1d_0, f_0e_1 = g_2d_1, f_0e_2 = -2g_0d_2, g'_0g'_1g'_2 = g_0g_1g_2.$$

4. Main Calculation

Let D be a subcomplex of $E(sM(3)^*)$ generated by $\{t_{i,j} \mid i=1, 2, 3, j \in \mathbf{Z}/3, j \neq 2 \text{ if } i=3\}$. We put $\zeta_3 = t_{3,0} + t_{3,1} + t_{3,2}$.

Lemma 4.1. $\delta(\zeta_3) = 0$ and $\{E(sM(3)^*), \delta\}$ is isomorphic to $\{D \otimes E(\zeta_3), \delta \otimes 1\}$. Therefore $\text{Ext}_{U(M(3))}^*(\mathbf{F}_p, \mathbf{F}_p)$ is isomorphic to $H^*(D) \otimes E(\zeta_3)$ as an algebra.

We filter $E(sM(3)^*)$ and D by $F^s E(sM(3)^*)^m = \sum_{i \leq m-s} E(s(M(3)/A)^*)^{m-i} \otimes$

$E(sA^*)^i$, $F^s D^m = D^m \cap F^s E(sM(3)^*)^m$. Then $F^s E(sM(3)^*)^m = F^s D^m + F^{s-1} D^{m-1} \zeta_3$ and $0 = F^{m+1} D^m \subset F^m D^m = E(s(M(3)/A)^*)^m \subset F^{m-1} D^m \subset F^{m-2} D^m = D^m$ hold. Also note that $\delta(F^s E(sM(3)^*)^{s+t}) \subset F^{s+1} E(sM(3)^*)^{s+t+1}$. Thus we have spectral sequences $E_1^{s,t} \Rightarrow \text{Ext}_{\tilde{U}(M(3))}^{s,t}(\mathbf{F}_p, \mathbf{F}_p)$ and $\tilde{E}_1^{s,t} \Rightarrow H^*(D)$ associated with these filtrations. By the above lemma, the former spectral sequence is isomorphic to $\{\tilde{E}_r^{s,t} \otimes E(\zeta_3), \tilde{d}_r \otimes 1\}$. Hence it suffices to compute the latter. The \tilde{E}_1 -term is given by $\tilde{E}_1^{s,t} = H^{s+t}(F^s D/F^{s+1} D) = E(s(M(3)/A)^*)^s \otimes E(t_{3,0}, t_{3,1})^t$ and \tilde{d}_1 coincides with $\delta \otimes 1$. Therefore $\tilde{E}_2^{s,t} = \text{Ext}_{\tilde{U}(M(3)/A)}^*(\mathbf{F}_p, \mathbf{F}_p) \otimes E(t_{3,0}, t_{3,1})^t$ and \tilde{d}_2 is given by $\tilde{d}_2(t_{3,j}) = f_j (j=0, 1)$. By computing the \tilde{E}_3 -term, we find that $\tilde{E}_3 = \tilde{E}_\infty$ for dimensional reasons. It is not difficult (but very tedious) to solve the extension problem and we can determine the structure of $\text{Ext}_{\tilde{U}(M(3))}^*(\mathbf{F}_p, \mathbf{F}_p)$ as given below.

Theorem 4.2. 1) $\text{Ext}_{\tilde{U}(M(3))}^*(\mathbf{F}_p, \mathbf{F}_p)$ is generated by the following twenty-six elements ;

$h_j = [t_{1,j}]$, $\zeta_3 = [t_{3,0} + t_{3,1} + t_{3,2}]$, $g_j = [t_{1,j} t_{2,j}]$, $g'_j = [t_{1,j+1} t_{2,j}]$, $a_0 = [t_{1,0} t_{3,0} - t_{1,0} t_{3,1} + t_{2,2} t_{2,0}]$, $a_1 = [t_{1,1} t_{3,0} + 2t_{1,1} t_{3,1} + t_{2,0} t_{2,1}]$, $a_2 = [-2t_{1,2} t_{3,0} - t_{1,2} t_{3,1} + t_{2,1} t_{2,2}]$, $b_0 = [t_{1,0} t_{2,0} t_{3,0}]$, $b_1 = [t_{1,1} t_{2,1} t_{3,1}]$, $b_2 = [t_{1,2} t_{2,2} (-t_{3,0} - t_{3,1})]$, $b'_0 = [t_{1,1} t_{2,0} (-t_{3,0} - t_{3,1})]$, $b'_1 = [t_{1,2} t_{2,1} t_{3,0}]$, $b'_2 = [t_{1,0} t_{2,2} t_{3,1}]$, $c = [t_{1,2} t_{2,0} t_{3,0} - t_{1,0} t_{2,1} t_{3,0} + t_{1,2} t_{2,0} t_{3,1} - t_{1,1} t_{2,2} t_{3,1} + t_{2,0} t_{2,1} t_{2,2}]$, $u_0 = [t_{1,0} t_{2,2} t_{2,0} t_{3,0}]$, $u_1 = [t_{1,1} t_{2,0} t_{2,1} t_{3,1}]$, $u_2 = [t_{1,2} t_{2,1} t_{2,2} (-t_{3,0} - t_{3,1})]$, $w_j = [t_{1,j} t_{2,j-1} t_{2,j} t_{3,0} t_{3,1}]$ for $j \in \mathbf{Z}/3$.

$H^*(D)$ is generated by the above elements except for ζ_3 .

2) $H^s(D) = 0$ for $s > 8$. A basis of $H^s(D)$ ($0 \leq s \leq 8$) is given as follows ;

- $s=0$; 1.
- $s=1$; h_0, h_1, h_2 .
- $s=2$; $g_0, g_1, g_2, g'_0, g'_1, g'_2, a_0, a_1, a_1$.
- $s=3$; $h_1 g_0, h_2 g_1, h_0 g_2, h_0 a_0, h_1 a_1, h_2 a_2, h_0 a_1, h_1 a_2, h_2 a_0, b_0, b_1, b_2, b'_0, b'_1, b'_2, c$.
- $s=4$; $h_1 b_0, h_2 b_1, h_0 b_2, h_2 b_0, h_0 b_1, h_1 b_2, h_0 b'_0, h_1 b'_1, h_2 b'_2, a_0^2, a_1^2, a_2^2, a_0 a_1, a_1 a_2, a_2 a_0, u_0, u_1, u_2$.
- $s=5$; $h_0 a_1 a_2, h_1 a_0 a_1, h_2 a_1 a_2, h_0 a_2 a_0, h_0 a_0 a_1, h_1 a_1 a_2, h_2 a_2 a_0, a_1 b_0, a_2 b_1, a_0 b_2, a_2 b_0, a_0 b_1, a_1 b_2, w_0, w_1, w_2$.
- $s=6$; $h_1 a_2 b_0, h_2 a_0 b_1, h_0 a_1 b_2, h_0 w_1, h_1 w_2, h_2 w_0, h_1 w_0, h_2 w_1, h_0 w_2$.
- $s=7$; $g_1 w_0, g_2 w_0, g_0 w_2$.
- $s=8$; $h_2 g_1 w_0$.

REMARKS. 1) By (4, 1) the basis given above is an $E(\zeta_3)$ -basis of $\text{Ext}_{\tilde{U}(M(3))}^*(\mathbf{F}_p, \mathbf{F}_p)$.

2) An element x of $\text{Ext}_{U(M(3))}(\mathbf{F}_p, \mathbf{F}_p)$ is said to be of filtration degree t if the image of x by the isomorphism $\text{Ext}_{U(M(3))}(\mathbf{F}_p, \mathbf{F}_p) \cong \text{Ext}_{V(L(3))}(\mathbf{F}_p, \mathbf{F}_p) = \text{Ext}_{E^0 S(3)}(\mathbf{F}_p, \mathbf{F}_p)$ of (2, 3) belongs to $E_2^{*,t}$ in the spectral sequence (2, 1). We denote by $f\text{-deg } x$ the filtration degree of x . Then it is easy to see $f\text{-deg } h_j = 2$, $f\text{-deg } \zeta_3 = f\text{-deg } g_j = f\text{-deg } g'_j = 6$, $f\text{-deg } a_j = 8$, $f\text{-deg } b_j = f\text{-deg } b'_j = f\text{-deg } c = 12$, $f\text{-deg } u_j = 16$, $f\text{-deg } w_j = 22$ for $j \in \mathbf{Z}/3$.

The internal degree of an element of $\text{Ext}_{U(M(3))}(\mathbf{F}_p, \mathbf{F}_p)$ is the degree coming

from the grading of $S(3)$ which takes value in $\mathbf{Z}/2(p^3-1)$. We denote by $i\text{-deg } x$ the internal degree of x . Put $q=2(p-1)$. Noting that $p^2q \equiv (-p-1)q$ modulo $2(p^3-1)$, the internal degrees of the generator are given as follows; $i\text{-deg } h_j = i\text{-deg } a_j = p^j q$, $i\text{-deg } \zeta_3 = i\text{-deg } c = 0$, $i\text{-deg } g_j = i\text{-deg } b_j = p^j(p+2)q$, $i\text{-deg } g'_j = i\text{-deg } b'_j = p^j(2p+1)q$, $i\text{-deg } u_j = i\text{-deg } w_j = 2p^j q$ for $j \in \mathbf{Z}/3$.

These two kinds of degrees play an important role in the next section.

Let η_* denote the operator on $\text{Ext}_{U(M(3))}(\mathbf{F}_p, \mathbf{F}_p)$ induced by $\eta^*: L(3) \rightarrow L(3)$. By the definition of the generators in (4, 2), it is easy to verify the following

Proposition 4.3. $h_j = \eta_*^j h_0$, $g_j = \eta_*^j g_0$, $g'_j = \eta_*^j g'_0$ for $j=1, 2$, $\zeta_3 = \eta_* \zeta_3$, $a_1 = \eta_* a_0 + h_1 \zeta_3$, $a_2 = \eta_*^2 a_0 - h_2 \zeta_3$, $b_1 = \eta_* b_0$, $b_2 = \eta_*^2 b_0 - g_2 \zeta_3$, $b'_1 = \eta_* b'_0 + g'_1 \zeta_3$, $b'_2 = \eta_*^2 b'_0 + g'_2 \zeta_3$, $c = \eta_* c$, $u_1 = \eta_* u_0$, $u_2 = \eta_*^2 u_0 - h_2 a_2 \zeta_3$, $w_1 = \eta_* w_0 - u_1 \zeta_3$, $w_2 = \eta_*^2 w_0 + u_2 \zeta_3 - a_2^2 \zeta_3 / 2$.

Since $\eta_* \zeta_3 = \zeta_3$, η_* induces an automorphism $\tilde{\eta}_*$ of $H^*(D) = \text{Ext}_{\mathbb{F}(M(3))}^*(\mathbf{F}_p, \mathbf{F}_p) / (\zeta_3)$ which maps x_j to x_{j+1} for $x = h, g, g', a, b, b', u, w$ and $j \in \mathbf{Z}/3$.

Theorem 4.4. *Relations of $H^*(D)$ are given by the following and the relations obtained by applying $\tilde{\eta}_*$ ($j=1, 2$) to them;*

$$\begin{aligned} h_0 h_j &= 0 \text{ for } j \in \mathbf{Z}/3; \\ h_0 g_0 &= h_0 g_1 = h_0 g'_1 = h_0 g'_2 = 0, h_0 g'_0 = -h_1 g_0, h_0 a_2 = h_2 a_0; \\ g_0 g_j &= g_0 g'_j = g'_0 g'_j = 0 \text{ for } j \in \mathbf{Z}/3, h_0 b_0 = h_0 b'_2 = 0, h_0 b'_1 = -h_1 b_2, h_0 c = -3h_2 b_0, a_0 g_0 = \\ a_0 g'_2 &= 0, a_0 g_1 = -3h_0 b_1, a_0 g_2 = 2h_2 b'_2 - h_0 b_2, a_0 g'_0 = h_0 b'_0 - h_1 b_0, a_0 g'_1 = -3h_1 b_2; \\ h_0 a_1 a_2 &= h_1 a_2 a_0 = h_2 a_0 a_1, h_0 a_0^2 = h_0 u_0 = h_0 u_2 = 0, h_0 a_1^2 = h_1 a_0 a_1, h_0 u_1 = h_1 a_0 a_1 / 2, g_0 b_0 = g_0 b'_0 = \\ g_0 b'_2 &= 0, g_0 b_1 = -h_1 a_0 a_1 / 2, g_0 b_2 = h_0 a_2 a_0 / 2, g_0 b'_1 = h_0 a_1 a_2 / 6, g_0 c = -h_0 a_0 a_1 / 2, g'_0 b_0 = g'_0 b'_1 = \\ g'_0 b'_2 &= 0, g'_0 b_2 = h_0 a_1 a_2 / 6, g'_0 c = h_2 a_1 a_2 / 2, g'_0 b'_1 = h_1 a_1 a_2 / 2, g'_0 b'_2 = -h_0 a_0 a_1 / 2, a_0 b_0 = a_0 b'_2 = \\ 0, a_0 b'_0 &= -a_1 b_0, a_0 b'_1 = -a_1 b_2, a_0 c = -3a_2 b_0; \\ h_0 a_0 b_2 &= 0, h_0 a_0 b_1 = h_1 a_2 b_0, g_0 u_0 = g_0 u_1 = g'_0 u_0 = g'_0 u_1 = 0, g_0 u_2 = -h_0 a_1 b_2 / 2, g'_0 u_2 = -h_2 a_0 b_1, \\ h_0 w_0 &= 0, a_0 a_1 a_2 = a_0^3 = a_0 u_0 = 0, a_0 a_1^2 = -6h_0 w_1, a_0 a_2^2 = 6h_0 w_2, a_0 u_1 = -h_0 w_1, a_0 u_2 = \\ 2h_0 w_2, b_0 b'_j &= 0 \text{ for } j \in \mathbf{Z}/3, b_0 b_1 = -h_0 w_1, b_2 b_0 = -h_2 w_0, b_0 c = h_1 w_0, b'_0 b'_1 = h_2 w_1, b'_2 b'_0 = \\ h_1 w_0, b'_0 c &= -h_1 w_2; \\ a_0 w_j &= b'_0 u_j = 0 \text{ for } j \in \mathbf{Z}/3, g_0 w_0 = g_0 w_1 = g'_0 w_0 = g'_0 w_1 = 0, g'_0 w_2 = -g_2 w_1, b_0 a_1^2 = b_0 u_0 = \\ b_0 u_1 &= c u_0 = 0, b_0 a_2^2 = 2g_0 w_2, b_0 a_1 a_2 = 2g_1 w_0, b_0 u_2 = g_0 w_2; \\ h_2 g_1 w_0 &= h_0 g_2 w_1 = h_1 g_0 w_2, c w_0 = 0, b_0 w_j = b'_0 w_j = u_0 u_j = 0 \text{ for } j \in \mathbf{Z}/3 \end{aligned}$$

This completes a description of the structure of $\text{Ext}_{\mathbb{F}^0 S(3)}^*(\mathbf{F}_p, \mathbf{F}_p)$ by virtue of (2.3) and (4,1).

5. The Algebra Structure of the Cohomology of $S(3)$

We consider the spectral sequence (2.1) for $n=3$, $p \geq 5$. This spectral sequence is $(\mathbf{Z} \times \mathbf{Z} \times \mathbf{Z} / 2(p^3-1))$ -graded and we denote $E_r^{s,t,u}$ and $F^{s,t,u}$ the subspaces of $E_r^{s,t}$ and $F^{s,t}$ spanned by elements of internal degree u . From the calculation of the previous section, we have the following table of the E_2 -term, where the numbers in the parentheses in the table indicate the filtration degree

t of the elements.

$u \setminus s+t$	0	1	2	3	4	5
0	1(0)	$\zeta_3(6)$	—	$c(12)$	$c\zeta_3(18)$	$h_0 a_1 a_2(18)$
$p^j q$	—	$h_j(2)$	$a_j, h_j \zeta_3(8)$	$a_j \zeta_3(14)$	$h_{j+2} b_j(14)$	$a_{j+2} b_j, h_{j+2} b_j \zeta_3(20)$
$p^j(p+1)q$	—	—	—	$h_j a_{j+1}(10)$	$a_j a_{j+1}, h_j a_{j+1} \zeta_3(16)$	$a_j a_{j+1} \zeta_3(22)$
$p^j(p+2)q$	—	—	$g_j(6)$	$b_j, g_j \zeta_3(12)$	$b_j \zeta_3(18)$	$h_j a_j a_{j+1}(18)$
$p^j(2p+1)q$	—	—	$g'_j(6)$	$b'_j, g'_j \zeta_3(12)$	$b'_j \zeta_3(18)$	$h_{j+1} a_j a_{j+1}(18)$
$2p^j(p+1)q$	—	—	—	$h_{j+1} g_j(8)$	$h_{j+1} b_j, h_j b'_j, h_{j+1} g_j \zeta_3(14)$	$a_{j+1} b_j, h_{j+1} b_j \zeta_3(20)$
$2p^j q$	—	—	—	$h_j a_j(10)$	$a_j^2, u_j, h_j a_j \zeta_3(16)$	$w_j, a_j^2 \zeta_3, u_j \zeta_3(22)$

$u \setminus s+t$	6	7	8	9
0	$h_0 a_1 a_2 \zeta_3(24)$	—	$h_2 g_1 w_0(30)$	$h_2 g_1 w_0 \zeta_3(36)$
$p^j q$	$a_{j+2} b_j \zeta_3(26)$	—	—	—
$p^j(p+1)q$	$h_{j+1} a_{j+2} b_j(22)$	$g_{j+1} w_j, h_{j+1} a_{j+2} b_j \zeta_3(28)$	$g_{j+1} w_j \zeta_3(34)$	—
$p^j(p+2)q$	$h_{j+1} w_j, h_j a_j a_{j+1} \zeta_3(24)$	$h_{j+1} w_j \zeta_3(30)$	—	—
$p^j(2p+1)q$	$h_j w_{j+1}, h_{j+1} a_j a_{j+1} \zeta_3(24)$	$h_j w_{j+1} \zeta_3(30)$	—	—
$2p^j(p+1)q$	$a_{j+1} b_j \zeta_3(26)$	—	—	—
$2p^j q$	$w_j \zeta_3(28)$	—	—	—

The following facts are immediately verified from the table.

Lemma 5.1. *If $E_2^{m-t,t} \neq 0, E_2^{m+1-s,s} = 0$ holds for $s < t$. Therefore the spectral sequence of $(2, 1)$ collapses, that is, $E_2^{s,t} = E_\infty^{s,t}$.*

Lemma 5.2. *If $\sum_{s+t=m} E_2^{s,t,u} \neq 0$ for given $m \in \mathbf{Z}$ and $u \in \mathbf{Z}/2(p^3-1)$, then $E_2^{m-t,t,u} = 0$ for all but only one t . Hence if $\text{Ext}_{S(3)}^m(\mathbf{F}_p, \mathbf{F}_p) \neq 0$, there is a unique $t = \tau(m, u)$ such that $F^{m-t+1,t-1,u} = 0$ and $F^{m-t,t,u} = \text{Ext}_{S(3)}^m(\mathbf{F}_p, \mathbf{F}_p)$.*

Thus there are unique elements $h_j, \zeta_3, g_j, g'_j, a_j, b_j, b'_j, c, u_j, w_j (j \in \mathbf{Z}/3)$ of $\text{Ext}_{S(3)}(\mathbf{F}_p, \mathbf{F}_p)$ corresponding to the elements of the E_2 -term denoted by the same symbols. Let \bar{B} be a set of monomials of the above elements which corresponds to the $E(\zeta_3)$ -basis of the E_2 -term given in the previous section. We put $B = \bar{B} \cup$

$\{x'_3 | x \in \bar{B}\}$, then B is a basis of $\text{Ext}_{S(3)}(\mathbf{F}_p, \mathbf{F}_p)$. For $x \in \text{Ext}_{S(3)}^m(\mathbf{F}_p, \mathbf{F}_i)$, we denote by \tilde{x} the element of $E_2^{m-t, t, u}$ corresponding to x where $t = \tau(m, u)$. For $x \in B \cap \text{Ext}_{S(3)}^m(\mathbf{F}_p, \mathbf{F}_p)$, $y \in B \cap \text{Ext}_{S(3)}^{l, v}(\mathbf{F}_p, \mathbf{F}_p)$, suppose that $\tilde{x}\tilde{y} = \sum_i \nu_i z_i$ holds for $\nu_i \in \mathbf{F}_p, z_i \in B$ in $E_2^{m+l-t-t', t+t', u+v}$ where $t = \tau(m, u), t' = \tau(l, v)$, in other words, $xy = \sum_i \nu_i z_i$ holds modulo $F^{m+l-t-t'+1, t+t'-1, u+v}$. If $\tilde{x}\tilde{y} \neq 0$, (5.2) implies that $F^{m+l-t-t'+1, t+t'-1, u+v} = 0$. Hence xy exactly equals to $\sum_i \nu_i z_i$ in this case. In the case $\tilde{x}\tilde{y} = 0$, we can verify $xy = 0$ case by case. In fact, it suffices to deal with the case $F^{m+l-t-t', t+t'} \neq 0$. Then we only have to check the cases $(x, y) = (a_j, b_j), (a_j, b'_{j-1}), (a_0 a_1, a_2), (a_j^2, a_j), (h_j, w_j), (a_j, u_j), (b_i, b'_j), (a_i, w_j), (b_i, w_j), (b'_i, w_j), (c, w_j), (u_i, u_j)$ for $i, j \in \mathbb{Z}/3$. In any of these cases, since $\text{Ext}_{S(3)}^{m+l, u+v}(\mathbf{F}_p, \mathbf{F}_p) = 0$, the assertion follows. Similarly, for $x \in B, \eta_* \tilde{x} = \sum_i \mu_i \tilde{y}_i (\mu_i \in \mathbf{F}_p, y_i \in B)$ implies $\eta_* x = \sum_i \mu_i y_i$ where the latter η_* is the operation of $\text{Ext}_{S(3)}(\mathbf{F}_p, \mathbf{F}_p)$ induced by the p -th power map of $S(3)$. Thus we have shown

Theorem 5.3. *$\text{Ext}_{S(3)}(\mathbf{F}_p, \mathbf{F}_p)$ is isomorphic to $\text{Ext}_{U(M(3))}(\mathbf{F}_p, \mathbf{F}_p)$ as an algebra over \mathbf{F}_p and the isomorphism commutes with the operations induced by the p -th power map of $S(3)$.*

References

- [1] J.F. Adams: Stable Homotopy and Generalized Homology, University of Chicago Press, Chicago, 1974.
- [2] J.F. Adams: Localisation and Completion, Mathematics Lecture Notes, University of Chicago, 1975.
- [3] P.E. Conner and L. Smith: *On the complex bordism of finite complexes*, Publ. Math. IHES no. 37 (1970), 117-212.
- [4] A. Iwai and N. Shimada: *A remark on resolutions for Hopf algebras*, Publ. RIMS Kyoto Univ. 1 (1966), 187-198.
- [5] P.S. Landweber: *Annihilator ideals and primitive elements in complex cobordism*, Ill. J. Math. 17 (1973), 273-284.
- [6] P.S. Landweber: *Associated prime ideals and Hopf algebras*, J. Pure Appl. Algebra 3 (1973), 43-58.
- [7] J.P. May: *The cohomology of restricted Lie algebras and of Hopf algebras*, J. Algebra 12 (1969), 533-568.
- [8] J.W. Milnor and J.C. Moore: *On the structure of Hopf algebras*, Ann. of Math. 81 (1965), 211-264.
- [9] Miller, H.R. and D.C. Ravenel: *Morava stabilizer algebras and the localization of Novikov's E_2 -term*, Duke Math. J. 44 (1977), 433-447.
- [10] H.R. Miller and D.C. Ravenel, and W.S. Wilson: *Periodic phenomena in the Adams-Novikov spectral sequence*, Ann. of Math. 106 (1977), 469-516.
- [11] J. Morava: *Structure theorem for cobordism comodules*, Preprint.
- [12] D.C. Ravenel: *The structure of Morava stabilizer algebras*, Invent. Math. 37 (1976), 106-120.
- [13] D.C. Ravenel: *The cohomology of the Morava stabilizer algebra*, Math. Zeit.

- 152 (1977), 287–297.
- [14] D.C. Ravenel: *Complex Cobordism and Stable Homotopy Groups of Spheres*, Academic Press, 1986.
- [15] H. Toda: *On spectra realizing exterior parts of the Steenrod algebra*, *Topology* **10** (1971), 53–65.

Department of Mathematics
University of Osaka Prefecture
Sakai, Osaka 591
Japan

