

## On an extension of the notion of Reedy category

Clemens Berger · Ieke Moerdijk

Received: 30 September 2009 / Accepted: 26 July 2010 / Published online: 9 September 2010  
© The Author(s) 2010. This article is published with open access at Springerlink.com

**Abstract** We extend the classical notion of a Reedy category so as to allow non-trivial automorphisms. Our extension includes many important examples occurring in topology such as Segal’s category  $\Gamma$ , or the total category of a crossed simplicial group such as Connes’ cyclic category  $\Lambda$ . For any generalized Reedy category  $\mathbb{R}$  and any cofibrantly generated model category  $\mathcal{E}$ , the functor category  $\mathcal{E}^{\mathbb{R}}$  is shown to carry a canonical model structure of Reedy type.

**Keywords** Generalized Reedy category · Quillen model category · Crossed group · Dendroidal set

**Mathematics Subject Classification (2000)** Primary 18G55 · 55U35;  
Secondary 18G30 · 20N99

### 0 Introduction

A Reedy category is a category  $\mathbb{R}$  equipped with a structure which makes it possible to prove that, for any Quillen [32] model category  $\mathcal{E}$ , the functor category  $\mathcal{E}^{\mathbb{R}}$  inherits a model structure, in which the cofibrations, weak equivalences and fibrations can all three be described explicitly in terms of those in  $\mathcal{E}$ . Prime examples of such Reedy categories are the simplex category  $\Delta$  and its dual  $\Delta^{\text{op}}$ ; the corresponding model structure on cosimplicial spaces goes back to Bousfield–Kan [9], while the model structure on simplicial objects in a model category  $\mathcal{E}$  is described in an unpublished manuscript by Reedy [34]. The general result for an

---

C. Berger (✉)

Lab. J.-A. Dieudonné, Université de Nice-Sophia Antipolis, Parc Valrose, 06108 Nice, France  
e-mail: cberger@math.unice.fr

I. Moerdijk

Mathematisch Instituut, Postbus 80.010, 3508 TA Utrecht, The Netherlands  
e-mail: moerdijk@math.uu.nl

arbitrary model category  $\mathcal{E}$  and Reedy category  $\mathbb{R}$  is by now a standard and important tool in homotopy theory, well explained in several textbooks, see e.g. [20, 22, 23].

As is well known, Reedy categories are skeletal, and moreover do not permit non-trivial automorphisms. There are, however, important cases in which it is possible to establish a Reedy-like model structure on the functor category  $\mathcal{E}^{\mathbb{R}}$  even though  $\mathbb{R}$  does have non-trivial automorphisms. One example is the strict model structure on  $\Gamma$ -spaces (space-valued presheaves on Segal's [36] category  $\Gamma$ ) established by Bousfield–Friedlander [8]. Another example is the case of cyclic spaces (space-valued presheaves on Connes' category  $\Lambda$ , see [14]). This paper grew out of a third example, namely the category of dendroidal spaces [30], which carries a Reedy-like model structure, although a dendroidal space is by definition a presheaf on a category  $\Omega$  of trees containing many automorphisms. Using the results of this paper, it has been shown that a localization of our Reedy-like model structure on dendroidal spaces is equivalent to a model structure on coloured topological operads, cf. [12, 13].

In this paper, we introduce the notion of a *generalized Reedy category*, and prove that for any such category  $\mathbb{R}$  and any  $\mathbb{R}$ -projective (e.g. cofibrantly generated) Quillen model category  $\mathcal{E}$ , the functor category  $\mathcal{E}^{\mathbb{R}}$  inherits a model structure, in which the cofibrations, weak equivalences and fibrations can again be described explicitly in terms of those in  $\mathcal{E}$ . Any classical Reedy category is a generalized Reedy category in our sense; in fact, a generalized Reedy category is equivalent to a classical one if and only if it has no non-trivial automorphisms. Segal's category  $\Gamma$  (as well as its dual) and the cyclic category  $\Lambda$  of Connes are examples of generalized Reedy categories, as is any (finite) group or groupoid. The cyclic category is an example of the total category associated to a crossed simplicial group [18, 26]; we will show that the total category of any *crossed group* on a classical Reedy category is a generalized Reedy category. This method yields many interesting examples of generalized Reedy categories with non-trivial automorphisms. In particular, the category  $\Omega$  mentioned above is of this type. Other examples of generalized Reedy categories relevant in homotopy theory are the *orbit category* of a finite or compact Lie group, and the total category associated to a *complex of groups*, see e.g. [21].

The results of this paper lead to interesting applications and further questions. We already mentioned the comparison between dendroidal spaces and coloured topological operads, which is analogous to the comparison between complete Segal spaces (a localization of the Reedy model structure on simplicial spaces) and topologically enriched categories—see [6, 11–13, 25, 27, 35]. We expect the Reedy model structure on spaces over a complex of groups to be useful in describing the derived category of the corresponding orbifold. A precise comparison would refine the weak homotopy equivalence between (the classifying spaces of) the complex of groups and the proper étale groupoid of the corresponding orbifold, cf. [29]. Another topic to be explored further is the relation between various models for cyclic homology (see e.g. [33]) and the Reedy model structure on cyclic spaces given by applying our main theorem to Connes' category  $\Lambda$ . In this context, we note that it is known [10] that a localization of this model structure is Quillen equivalent to the model structure [15] on cyclic sets.

In a recent paper, Angeltveit [1] studies Reedy categories enriched in a monoidal model category, and obtains examples of such from non-symmetric operads. We expect that a similar enrichment is possible for our generalized Reedy categories, so that Angeltveit's construction can be applied to symmetric operads as well. It would also be of interest to study the functoriality of the model structure on  $\mathcal{E}^{\mathbb{R}}$  as a functor of  $\mathbb{R}$ , and extend the results of Barwick [3] to our context.

To conclude this introduction, we describe the contents of the different sections of this paper. In Sect. 1, we present our notion of generalized Reedy category, state the main theorem

on the existence of a model structure (Theorem 1.6), and list some of the main examples. In Sect. 2, we explain a general method for constructing generalized Reedy categories out of classical ones by means of crossed groups. Sections 3 and 4 contain some technical preliminaries for the proof of the main theorem which will be given in Sect. 5. In Sect. 6, we give a brief introduction into skeleta and coskeleta for functor categories of the form  $\mathcal{E}^{\mathbb{R}}$ . We then discuss a special class of dualizable generalized Reedy categories  $\mathbb{R}$  for which the skeleta of set-valued presheaves on  $\mathbb{R}$  have a simple, explicit description. In Sect. 7, we obtain a refinement of the main theorem (Theorem 7.6) giving sufficient conditions on  $\mathbb{R}$  and  $\mathcal{E}$  for the Reedy model structure on  $\mathcal{E}^{\mathbb{R}^{\text{op}}}$  to be monoidal.

### 1 Generalized Reedy categories

Recall that a subcategory  $\mathbb{S}$  of  $\mathbb{R}$  is called *wide* if  $\mathbb{S}$  has the same objects as  $\mathbb{R}$ . An example of a wide subcategory of  $\mathbb{R}$  is the *maximal subgroupoid*  $\text{Iso}(\mathbb{R})$  of  $\mathbb{R}$ .

**Definition 1.1** A *generalized Reedy structure* on a small category  $\mathbb{R}$  consists of wide subcategories  $\mathbb{R}^+$ ,  $\mathbb{R}^-$ , and a degree-function  $d : \text{Ob}(\mathbb{R}) \rightarrow \mathbb{N}$  satisfying the following four axioms:

- (i) non-invertible morphisms in  $\mathbb{R}^+$  (resp.  $\mathbb{R}^-$ ) raise (resp. lower) the degree; isomorphisms in  $\mathbb{R}$  preserve the degree;
- (ii)  $\mathbb{R}^+ \cap \mathbb{R}^- = \text{Iso}(\mathbb{R})$ ;
- (iii) every morphism  $f$  of  $\mathbb{R}$  factors as  $f = gh$  with  $g \in \mathbb{R}^+$  and  $h \in \mathbb{R}^-$ , and this factorization is unique up to isomorphism;
- (iv) If  $\theta f = f$  for  $\theta \in \text{Iso}(\mathbb{R})$  and  $f \in \mathbb{R}^-$ , then  $\theta$  is an identity.

A generalized Reedy structure is *dualizable* if in addition the following axiom holds:

- (iv)' If  $f\theta = f$  for  $\theta \in \text{Iso}(\mathbb{R})$  and  $f \in \mathbb{R}^+$ , then  $\theta$  is an identity.

A (dualizable) generalized Reedy category is a small category equipped with a (dualizable) generalized Reedy structure.

A morphism of generalized Reedy categories  $\mathbb{R} \rightarrow \mathbb{S}$  is a functor which takes  $\mathbb{R}^+$  (resp.  $\mathbb{R}^-$ ) to  $\mathbb{S}^+$  (resp.  $\mathbb{S}^-$ ) and which preserves the degree.

*Remark 1.2* The inclusion from left to right in axiom (ii) follows from axiom (i). Axiom (iv) says that automorphisms in  $\mathbb{R}$  consider morphisms of  $\mathbb{R}^-$  as epimorphisms. This last axiom implies that the isomorphism in (iii) is unique. The axioms (i)-(ii) are self-dual while axiom (iv) is dual to axiom (iv)'. A generalized Reedy category  $\mathbb{R}$  is thus dualizable if and only if  $\mathbb{R}^{\text{op}}$  is also a generalized Reedy category. Most of the examples that we are aware of are dualizable. The asymmetry in the definition is related to the asymmetry of the projective model structure on objects with a group action, which enters in Theorem 1.6; cf. the proof of Lemma 5.5.

*Remark 1.3* If  $\mathbb{R}$  is a generalized Reedy category, an equivalence of categories  $\mathbb{R}' \xrightarrow{\sim} \mathbb{R}$  induces a generalized Reedy structure on  $\mathbb{R}'$ . In this sense, the existence of a generalized Reedy structure is invariant under equivalence of categories.

*Remark 1.4* Recall that in the literature (cf. [20, 22, 23, 34]) a category  $\mathbb{R}$ , equipped with  $\mathbb{R}^+$ ,  $\mathbb{R}^-$  and  $d$  as above, is called a *Reedy category* if it satisfies the following two axioms:

- (i) non-identity morphisms in  $\mathbb{R}^+$  (resp. in  $\mathbb{R}^-$ ) raise (resp. lower) degree;
- (ii) every morphism in  $\mathbb{R}$  factors uniquely as a morphism in  $\mathbb{R}^-$  followed by one in  $\mathbb{R}^+$ .

Any such Reedy category is a dualizable generalized Reedy category in our sense. To emphasize the distinction with generalized Reedy categories we will refer to the classical ones as *strict Reedy categories*. The notion of a strict Reedy category is not invariant under equivalence of categories. In fact, one checks that in a strict Reedy category every isomorphism is an identity.<sup>1</sup> A generalized Reedy category is equivalent to a strict one if and only if it has no non-trivial automorphisms, and is itself strict if and only if it is moreover skeletal.

*Remark 1.5* As for strict Reedy categories, all the results concerning a fixed generalized Reedy category  $\mathbb{R}$  go through if the degree-function takes values in an arbitrary well-ordered set. (However, with these more general degree-functions, the notion of a morphism of Reedy categories is more subtle to define).

For a generalized Reedy category  $\mathbb{R}$ , we introduce the following notions, which are classical in the case of a strict Reedy category. For each object  $r$  of  $\mathbb{R}$ , the category  $\mathbb{R}^+(r)$  has as objects the *non-invertible* morphisms in  $\mathbb{R}^+$  with codomain  $r$ , and as morphisms from  $u : s \rightarrow r$  to  $u' : s' \rightarrow r$  all  $w : s \rightarrow s'$  such that  $u = u'w$ . Observe that axiom (iii) implies that  $w \in \mathbb{R}^+$ ; moreover, the automorphism group  $\text{Aut}(r)$  acts on the category  $\mathbb{R}^+(r)$  by composition. For each functor  $X : \mathbb{R} \rightarrow \mathcal{E}$  and each object  $r$  of  $\mathbb{R}$ , the *r-th latching object*  $L_r(X)$  of  $X$  is defined to be

$$L_r(X) = \lim_{s \rightarrow r} X_s$$

where the colimit is taken over the category  $\mathbb{R}^+(r)$ . We will always assume  $\mathcal{E}$  to be sufficiently cocomplete for this colimit to exist (in many examples this colimit is finite). Note that  $\text{Aut}(r)$  acts on  $L_r(X)$ .

Dually, for each object  $r$  of  $\mathbb{R}$ , the category  $\mathbb{R}^-(r)$  has as objects the *non-invertible* morphisms in  $\mathbb{R}^-$  with domain  $r$ , and as morphisms from  $u : r \rightarrow s$  to  $u' : r \rightarrow s'$  all  $w : s \rightarrow s'$  such that  $u' = wu$ . Observe that axiom (iii) implies that  $w \in \mathbb{R}^-$ ; moreover, the automorphism group  $\text{Aut}(r)$  acts on the category  $\mathbb{R}^-(r)$  by precomposition. For each object  $X$  of  $\mathcal{E}^{\mathbb{R}}$  and each object  $r$  of  $\mathbb{R}$ , the *r-th matching object*  $M_r(X)$  of  $X$  is defined to be

$$M_r(X) = \lim_{r \rightarrow s} X_s$$

where the limit is taken over the category  $\mathbb{R}^-(r)$ . We will always assume  $\mathcal{E}$  to be sufficiently complete for this limit to exist (in many examples this limit is finite). Note that  $\text{Aut}(r)$  acts on  $M_r(X)$ .

Each object  $X$  of the functor category  $\mathcal{E}^{\mathbb{R}}$  defines for any object  $r$  of  $\mathbb{R}$  natural  $\text{Aut}(r)$ -equivariant maps  $L_r(X) \rightarrow X_r \rightarrow M_r(X)$ . For a map  $f : X \rightarrow Y$  in  $\mathcal{E}^{\mathbb{R}}$  these give rise to *relative latching*, resp. *matching* maps

$$X_r \cup_{L_r(X)} L_r(Y) \longrightarrow Y_r, \quad \text{resp.} \quad X_r \longrightarrow M_r(X) \times_{M_r(Y)} Y_r.$$

Recall that for any group (or groupoid)  $\Gamma$  and any *cofibrantly generated* model category  $\mathcal{E}$ , the category  $\mathcal{E}^\Gamma$  of objects of  $\mathcal{E}$  with right  $\Gamma$ -action carries a projective model structure, in which weak equivalences and fibrations are defined by forgetting the  $\Gamma$ -action. In general,

<sup>1</sup> Indeed, for an isomorphism  $f$ , let  $f = gh$  and  $hf^{-1} = g'h'$  be the unique factorizations. Then  $id = ghf^{-1} = (gg')h'$ , so  $h' = id$  and  $gg' = id$ , whence  $g = id$  and  $g' = id$  since  $g, g' \in \mathbb{R}^+$ . Thus  $f = h \in \mathbb{R}^-$ . The same argument applied to  $f^{-1}$  shows that  $f$  preserves the degree, hence  $f = id$ .

a Quillen model category  $\mathcal{E}$  will be called  $\mathbb{R}$ -projective, if for each object  $r$  of  $\mathbb{R}$ , the category  $\mathcal{E}^{\text{Aut}(r)}$  admits a projective model structure. For  $\mathbb{R}$ -projective model categories  $\mathcal{E}$ , we introduce the following notions:

A map  $f : X \rightarrow Y$  in  $\mathcal{E}^{\mathbb{R}}$  is called a

- *Reedy cofibration* if for each  $r$ , the relative latching map  $X_r \cup_{L_r(X)} L_r(Y) \rightarrow Y_r$  is a cofibration in  $\mathcal{E}^{\text{Aut}(r)}$ ;
- *Reedy weak equivalence* if for each  $r$ , the induced map  $f_r : X_r \rightarrow Y_r$  is a weak equivalence in  $\mathcal{E}^{\text{Aut}(r)}$ ;
- *Reedy fibration* if for each  $r$ , the relative matching map  $X_r \rightarrow M_r(X) \times_{M_r(Y)} Y_r$  is a fibration in  $\mathcal{E}^{\text{Aut}(r)}$ .

Observe that the automorphism group  $\text{Aut}(r)$  really enters only in the definition of a Reedy cofibration, by definition of the model structure on  $\mathcal{E}^{\text{Aut}(r)}$  just described.

**Theorem 1.6** *Let  $\mathbb{R}$  be a generalized Reedy category and let  $\mathcal{E}$  be an  $\mathbb{R}$ -projective Quillen model category in which the relevant limits and colimits exist (for instance,  $\mathcal{E}$  can be any cofibrantly generated model category). With the above classes of Reedy cofibrations, Reedy weak equivalences and Reedy fibrations, the functor category  $\mathcal{E}^{\mathbb{R}}$  is a Quillen model category.*

The proof will be supplied in Sect. 5. Notice that if  $\mathbb{R} = \mathbb{R}^+$  then the constant functor  $\mathcal{E} \rightarrow \mathcal{E}^{\mathbb{R}}$  sends weak equivalences and fibrations in  $\mathcal{E}$  to Reedy weak equivalences and Reedy fibrations in  $\mathcal{E}^{\mathbb{R}}$ . Thus, we obtain the following corollary which is well known for strict Reedy categories, cf. the two books [22,23].

**Corollary 1.7** *Let  $\mathcal{E}$  and  $\mathbb{R}$  be as in Theorem 1.6. If  $\mathbb{R} = \mathbb{R}^+$  then  $\varinjlim : \mathcal{E}^{\mathbb{R}} \rightarrow \mathcal{E}$  is a left Quillen functor.*

*Remark 1.8* A Quillen model category  $\mathcal{E}$  could be called  $\mathbb{R}$ -injective, if for each object  $r$  of  $\mathbb{R}$ , the category  $\mathcal{E}^{\text{Aut}(r)}$  admits an injective model structure, i.e. one for which the forgetful functor to  $\mathcal{E}$  detects weak equivalences and cofibrations. The proof of Theorem 1.6 will show that if  $\mathbb{R}^{\text{op}}$  is a generalized Reedy category and  $\mathcal{E}$  is  $\mathbb{R}$ -injective then  $\mathcal{E}^{\mathbb{R}}$  admits a model structure in which the three classes of maps are defined in a similar manner, but with reference to the injective model structure on each  $\mathcal{E}^{\text{Aut}(r)}$ . In this case, a dual of Corollary 1.7 holds, stating that if  $\mathbb{R} = \mathbb{R}^-$  then  $\varprojlim : \mathcal{E}^{\mathbb{R}} \rightarrow \mathcal{E}$  is a right Quillen functor.

*Examples 1.9* We give a list of first examples which, among other things, show that generalized Reedy categories occur naturally in several different contexts in homotopy theory. More examples are provided in Sect.2.

- (a) For completeness, we mention (again) that any strict Reedy category is a dualizable generalized Reedy category (cf. Remark 1.4). This applies in particular to standard examples of Reedy categories such as the simplex category  $\Delta$  and its dual, as well as to  $(\mathbb{N}, <)$ ,  $\leftarrow \cdot \rightarrow$ ,  $\cdot \dashrightarrow$  (relevant for homotopy colimits of sequences, for homotopy pushouts and for homotopy coequalizers). Other examples are Joyal’s category of finite disks and its dual  $\Theta$  (cf. [5,24]).
- (b) Segal’s category  $\Gamma$  (cf. [36]) is a dualizable generalized Reedy category. In fact,  $\Gamma^{\text{op}}$  is equivalent to the category  $\text{Fin}_*$  of finite pointed sets, and one can take  $\text{Fin}_*^+$  to consist of monomorphisms and  $\text{Fin}_*^-$  of epimorphisms, while the degree-function is given by cardinality. If  $\mathcal{E}$  is the category of simplicial sets, the Reedy model structure on  $\mathcal{E}^{\Gamma^{\text{op}}}$  given by Theorem 1.6 was discussed in Bousfield–Friedlander [8] and referred to as the strict model structure on  $\Gamma$ -spaces. The simplicial circle  $\Delta[1]/\partial\Delta[1]$ , when viewed as a functor  $\Delta \rightarrow \Gamma$  (cf. [5,36]), is a morphism of generalized Reedy categories.

- (c) The category  $\text{Fin}$  of finite sets carries a dualizable generalized Reedy structure, analogous to the pointed case. A skeleton of  $\text{Fin}$  is often denoted by  $\Delta_{\text{sym}}$ , and  $\mathcal{E}^{\text{op}}_{\Delta_{\text{sym}}}$  is referred to as the category of *symmetric simplicial objects* in  $\mathcal{E}$ , cf. [2, 10]. The inclusion  $\Delta \hookrightarrow \Delta_{\text{sym}}$  is a morphism of generalized Reedy categories.
- (d) Any group(oid) is a generalized Reedy category.
- (e) *Orbit categories.* The orbit category  $\mathcal{O}(G)$  of a *finite group*  $G$  has the subgroups of  $G$  as objects, and the  $G$ -equivariant maps  $G/H \rightarrow G/K$  as morphisms. This orbit category is a generalized Reedy category with  $\mathcal{O}(G) = \mathcal{O}(G)^-$  and  $d(H) = \text{card}(G/H)$  (the index of  $H$  in  $G$ ). There is also a generalized Reedy structure on  $\mathcal{O}(G)$  with  $\mathcal{O}(G) = \mathcal{O}(G)^+$  and  $d(H) = \text{card}(H)$ . If  $G$  is not finite, the first structure still makes sense for subgroups of finite index, the second one for finite subgroups. The orbit category  $\mathcal{O}(G)$  of a *compact Lie group*  $G$  is the category with closed subgroups of  $G$  as objects and  $G$ -homotopy classes of  $G$ -maps  $G/H \rightarrow G/K$  as morphisms from  $H$  to  $K$ . This is again a generalized Reedy category with  $\mathcal{O}(G) = \mathcal{O}(G)^+$ . The degree of an object  $H$  now takes values in  $\mathbb{N} \times \mathbb{N}$  with the lexicographical ordering, and is defined by  $d(H) = (\text{dim}(H), \text{card}(\pi_0 H))$ . Notice that this generalized Reedy structure does not admit a dual structure in which  $\mathcal{O}(G) = \mathcal{O}(G)^-$  like in the case of finite groups, because there may be infinite increasing sequences of closed subgroups, e.g. the subgroups  $\mathbb{Z}/p^n\mathbb{Z}$  of the circle  $S^1$ . This contradicts the existence of a degree-function for which the morphisms are degree-lowering.
- (f) *Complexes of groups.* Let  $X$  be a simplicial complex. Recall that a complex of groups  $G$  over  $X$  assigns to each simplex  $\sigma \in X$  a group  $G_\sigma$ , to each inclusion  $\sigma \subseteq \tau$  an *injective* group homomorphism  $\phi_{\sigma,\tau} : G_\tau \rightarrow G_\sigma$ , and to each sequence  $\rho \subseteq \sigma \subseteq \tau$  a specific element  $g = g_{\rho,\sigma,\tau} \in G_\rho$  such that the triangle

$$\begin{array}{ccc} G_\tau & \rightarrow & G_\sigma \\ & \searrow & \downarrow \\ & & G_\rho \end{array}$$

commutes up to conjugation by  $g$ , i.e. for each  $x \in G_\tau$ :

$$g\phi_{\rho,\tau}(x)g^{-1} = \phi_{\rho,\sigma}(\phi_{\sigma,\tau}(x)).$$

Moreover, for  $\pi \subseteq \rho \subseteq \sigma \subseteq \tau$ , the following coherence condition should be satisfied:

$$\phi_{\pi,\rho}(g_{\rho,\sigma,\tau})g_{\pi,\rho,\tau} = g_{\pi,\rho,\sigma}g_{\pi,\sigma,\tau}.$$

Such complexes of groups can be used to model orbifold structures on a triangulated space  $|X|$ , see [21, 29]. To each complex of groups  $G$  over  $X$  is associated a category  $\Delta_X(G)$  whose objects are the simplices  $\sigma \in X$ ; if  $\sigma \subseteq \tau$  then morphisms  $y : \sigma \rightarrow \tau$  in  $\Delta_X(G)$  are given by elements  $y \in G_\sigma$ . Composition of  $y : \sigma \rightarrow \tau$  and  $x : \rho \rightarrow \sigma$  is defined to be  $\phi_{\rho,\sigma}(y)x : \rho \rightarrow \sigma$ . The coherence condition implies that this composition is associative. The category  $\Delta_X(G)$  is a generalized Reedy category in which the degree of  $\sigma$  is the dimension of the simplex, and for which  $\Delta_X(G) = \Delta_X(G)^+$ .

The class of generalized Reedy categories is closed under arbitrary coproducts and under finite products. A more subtle closure property is the following:

**Proposition 1.10** *Let  $\mathbb{S} \rightarrow \mathbb{R}$  be a fibered category over  $\mathbb{R}$ . Suppose that the base  $\mathbb{R}$  and each of the fibers  $\mathbb{S}_r$  are equipped with generalized Reedy structures. Assume furthermore that for each morphism  $\alpha : r \rightarrow s$  in the base  $\mathbb{R}$ ,*

- (i) the base change  $\alpha^* : \mathbb{S}_s \rightarrow \mathbb{S}_r$  preserves the degree;
- (ii) if  $\alpha$  belongs to  $\mathbb{R}^+$  then  $\alpha^*$  takes  $\mathbb{S}_s^+$  to  $\mathbb{S}_r^+$ ;
- (iii) if  $\alpha$  belongs to  $\mathbb{R}^-$  then  $\alpha^*$  has a left adjoint  $\alpha_!$  which takes  $\mathbb{S}_r^-$  to  $\mathbb{S}_s^-$ .

Then  $\mathbb{S}$  can be equipped with a generalized Reedy structure such that the fiber inclusions  $\mathbb{S}_r \hookrightarrow \mathbb{S}$  and the projection  $\mathbb{S} \rightarrow \mathbb{R}$  preserve the factorization systems.

*Proof* Consider a morphism  $f : x \rightarrow y$  in  $\mathbb{S}$  over  $\alpha : r \rightarrow s$  in  $\mathbb{R}$ . Say  $f \in \mathbb{S}^+$  if  $\alpha \in \mathbb{R}^+$  and the unique morphism  $x \rightarrow \alpha^*(y)$  in  $\mathbb{S}_r$  determined by a cartesian lift  $\alpha^*(y) \rightarrow y$  of  $\alpha$  lies in  $\mathbb{S}_r^+$ . Say  $f \in \mathbb{S}^-$  if  $\alpha \in \mathbb{R}^-$  and the unique morphism  $\alpha_!(x) \rightarrow y$  in  $\mathbb{S}_y$  determined by a cocartesian lift  $x \rightarrow \alpha_!(x)$  of  $\alpha$  lies in  $\mathbb{S}_s^-$ . For  $x \in \mathbb{S}_r$ , define the degree by  $d_{\mathbb{S}}(x) = d_{\mathbb{R}}(r) + d_{\mathbb{S}_r}(x)$ . With these definitions, it is straightforward to verify that  $\mathbb{S}$  is a generalized Reedy category.

Example 1.9(f) is a special instance of Proposition 1.10.

**Corollary 1.11** *Let  $\mathbb{R}$  be a generalized Reedy category for which  $\mathbb{R} = \mathbb{R}^+$ , and let  $\Phi : \mathbb{R}^{\text{op}} \rightarrow \text{Cat}$  be a diagram of Reedy categories and morphisms of Reedy categories. Then the Grothendieck construction  $\mathbb{S} = \int_{\mathbb{R}} \Phi$  is again a generalized Reedy category.*

## 2 Crossed groups

In this section, we introduce the notion of a *crossed group*  $G$  on a category  $\mathbb{R}$ , and discuss the construction of the associated *total category*  $\mathbb{R}G$ . We will show that for any strict Reedy category  $\mathbb{R}$  and crossed  $\mathbb{R}$ -group  $G$ , the total category  $\mathbb{R}G$  is a generalized Reedy category, which is no longer strict unless  $G$  is trivial. Many of our examples of generalized Reedy categories are instances of this construction.

Crossed groups on the simplex category have been studied in the literature under the name *skew-simplicial groups* (see Krasauskas [26]), resp. *crossed simplicial groups* (see Fiedorowicz–Loday [18]). Recently, Batanin–Markl [4, 2.2] considered *crossed cosimplicial groups* which are crossed groups on the dual of the simplex category. Feigin–Tsygan already spelled out the axioms of a crossed group in [17, A4.1–4]. Cisinski considers the more general concept of a *thickening* in [10, 8.5.8].

**Definition 2.1** For any small category  $\mathbb{R}$ , a *crossed  $\mathbb{R}$ -group*  $G$  is a set-valued presheaf on  $\mathbb{R}$ , together with, for each object  $r$  of  $\mathbb{R}$ ,

- (i) a group structure on  $G_r$ ,
- (ii) left  $G_r$ -actions on the hom-sets  $\text{Hom}_{\mathbb{R}}(s, r)$  with codomain  $r$ ,

such that the following identities hold for all  $g, h \in G_r$ ,  $\alpha : s \rightarrow r$ ,  $\beta : t \rightarrow s$ ,

$$g_*(\alpha \circ \beta) = g_*(\alpha) \circ \alpha^*(g)_*(\beta), \tag{1}$$

$$g_*(1_r) = 1_r, \tag{2}$$

$$\alpha^*(g \cdot h) = h_*(\alpha)^*(g) \cdot \alpha^*(h), \tag{3}$$

$$\alpha^*(e_r) = e_s, \tag{4}$$

where the presheaf action of  $\alpha : s \rightarrow r$  is denoted by  $\alpha^* : G_r \rightarrow G_s$  and the group action of  $g \in G_r$  is denoted by  $g_* : \text{Hom}_{\mathbb{R}}(s, r) \rightarrow \text{Hom}_{\mathbb{R}}(s, r)$ . Moreover, for each object  $r$ , the identity of  $r$  (resp. neutral element of  $G_r$ ) is denoted by  $1_r$  (resp.  $e_r$ ).

*Remark 2.2* In what follows we shall make no difference in notation between composition in  $\mathbb{R}$  and composition in  $G_r$ , especially since both structures will agree in the total category  $\mathbb{R}G$ . In addition to the four identities spelled out in Definition 2.1, the following four identities also hold in any  $\mathbb{R}$ -crossed group  $G$  (by the axioms for a presheaf, resp. group action):

$$(\alpha\beta)^*(g) = \beta^*\alpha^*(g), \tag{5}$$

$$1_r^*(g) = g, \tag{6}$$

$$(gh)_*(\alpha) = g_*h_*(\alpha), \tag{7}$$

$$(e_r)_*(\alpha) = \alpha. \tag{8}$$

**Definition 2.3** For any small category  $\mathbb{R}$  and crossed  $\mathbb{R}$ -group  $G$ , the *total category*  $\mathbb{R}G$  is the category with the same objects as  $\mathbb{R}$ , and with morphisms  $r \rightarrow s$  the pairs  $(\alpha, g)$  where  $\alpha : r \rightarrow s$  belongs to  $\mathbb{R}$ , and  $g \in G_r$ . Composition of  $(\alpha, g) : s \rightarrow t$  and  $(\beta, h) : r \rightarrow s$  is defined as

$$(\alpha, g) \circ (\beta, h) = (\alpha \cdot g_*(\beta), \beta^*(g) \cdot h).$$

One easily checks that this composition is associative and has a two-sided unit  $(1_r, e_r)$  for each object  $r$  of  $\mathbb{R}G$ .

*Remark 2.4* In the special case where  $G$  is a constant presheaf (i.e.  $G = G_r$  for a fixed group  $G$  and  $\alpha^*(g) = g$  for all  $g$  and all  $\alpha$ ), the total category  $\mathbb{R}G$  reduces to the familiar Grothendieck construction for a diagram of categories on  $G$ .

In the special case where the left action of  $G$  on  $\mathbb{R}$  is trivial (i.e.  $g_*(\alpha) = \alpha$  for all  $g$  and all  $\alpha$ ), the crossed group is actually a presheaf of groups, and the total category  $\mathbb{R}G$  again reduces to a Grothendieck construction, this time for a diagram of groups on  $\mathbb{R}^{\text{op}}$ .

In these cases, Proposition 2.5 below becomes a special instance of Corollary 1.11.

Returning to the general case of a crossed  $\mathbb{R}$ -group  $G$ , notice that we always have a canonical embedding  $\mathbb{R} \hookrightarrow \mathbb{R}G$  which sends  $\alpha : r \rightarrow s$  to  $(\alpha, e_r) : r \rightarrow s$ , and identifies  $\mathbb{R}$  with a wide subcategory of  $\mathbb{R}G$ . Elements  $g \in G_r$  of the crossed group may be identified with *special automorphisms*  $(1_r, g)$  in the total category  $\mathbb{R}G$ , and every morphism  $(\alpha, g)$  in  $\mathbb{R}G$  factors uniquely as a special automorphism  $(1_r, g)$  followed by a morphism  $(\alpha, e_r)$  in  $\mathbb{R}$ . This *unique factorization property* is characteristic for total categories of crossed groups as asserted by:

**Proposition 2.5** *Let  $\mathbb{R} \subseteq \mathbb{S}$  be a wide subcategory and assume that there exist subgroups  $G_s \subseteq \text{Aut}_{\mathbb{S}}(s)$  of special automorphisms such that each morphism in  $\mathbb{S}$  factors uniquely as a special automorphism followed by a morphism in  $\mathbb{R}$ . Then the groups  $G_s$  define a crossed  $\mathbb{R}$ -group, and  $\mathbb{S}$  is isomorphic to  $\mathbb{R}G$  (under  $\mathbb{R}$ ).*

*Proof* For any morphism  $\alpha : r \rightarrow s$  of  $\mathbb{R}$  and special automorphism  $g \in G_s$ , the presheaf action of  $\mathbb{R}$  as well as the group action of  $G$  are defined by factoring the composite  $g\alpha : r \rightarrow s$  uniquely as in the hypothesis of the proposition, as

$$\begin{array}{ccc} r & \xrightarrow{\alpha} & s \\ \alpha^*(g) \downarrow & & \downarrow g \\ r & \xrightarrow{g_*(\alpha)} & s. \end{array}$$

With this explicit description, the proof of the identities of Definition 2.1 and of the isomorphism  $\mathbb{S} \cong \mathbb{R}G$  is a matter of (lengthy but) straightforward verification.



*Remark 2.6* Fiedorowicz–Loday [18] take Proposition 2.5 for  $\mathbb{R} = \Delta$  as the definition of a crossed simplicial group (with contravariant instead of covariant group action), and state Definition 2.1 of a crossed  $\Delta$ -group as a proposition.

*Example 2.7* The most prominent example of a crossed group is the *simplicial circle*  $C = \Delta[1]/\partial\Delta[1]$  whose total category  $\Delta C$  is isomorphic to the *cyclic category*  $\Lambda$  of Connes [14]. It is convenient to embed  $C$  in a larger crossed  $\Delta$ -group  $\Sigma$ , formed by the permutation groups  $\Sigma_{[n]}$  of the sets  $[n] = \{0, 1, \dots, n\}$ . The crossed  $\Delta$ -group structure of  $\Sigma$  is defined as follows: given  $\alpha : [m] \rightarrow [n]$  in  $\Delta$  and  $g : [n] \rightarrow [n]$  in  $\Sigma_{[n]}$ , the map  $\alpha^*(g) : [m] \rightarrow [m]$  is the unique permutation which is *order-preserving on the fibers* of  $\alpha$ , and for which  $g_*(\alpha) = g \circ \alpha \circ \alpha^*(g)^{-1} : [m] \rightarrow [n]$  is order-preserving:

$$\begin{array}{ccc} [m] & \xrightarrow{\alpha} & [n] \\ \alpha^*(g) \downarrow & & \downarrow g \\ [m] & \xrightarrow{g_*(\alpha)} & [n]. \end{array}$$

Let  $C_{[n]} \subset \Sigma_{[n]}$  be the subgroup generated by the cycle  $0 \mapsto 1 \mapsto \dots \mapsto n \mapsto 0$ . One checks that if  $g \in C_{[n]}$  then  $\alpha^*(g) \in C_{[m]}$  for each  $\alpha : [m] \rightarrow [n]$  in  $\Delta$ , so that  $C$  inherits a crossed  $\Delta$ -group structure. The total category  $\Delta C$  is then isomorphic to the cyclic category  $\Lambda$  of Connes [14], and embeds in the total category  $\Delta\Sigma$ . The latter has been described in detail by Feigin–Tsygan [17, A10] and plays an important role in the general classification of crossed  $\Delta$ -groups, see [18, 26].

*Example 2.8* One of the examples of a generalized Reedy category which motivated this paper is the category  $\Omega$  of trees introduced by Moerdijk–Weiss in [30]. The objects of this category are finite trees with a distinguished output edge and a set of distinguished input edges, as common in the context of operads. Any such tree  $T$  freely generates a symmetric coloured operad  $\Omega(T)$  whose colour-set is the set  $E(T)$  of edges of  $T$ ; the morphisms  $T \rightarrow T'$  in  $\Omega$  are the maps of symmetric coloured operads  $\Omega(T) \rightarrow \Omega(T')$ . For a more precise description, we refer to [30]. Here, it is enough to observe that any such morphism  $T \rightarrow T'$  induces a map  $E(T) \rightarrow E(T')$  in a functorial way, and that this induced map completely determines the morphism. The category  $\Omega$  carries a natural dualizable generalized Reedy structure, for which the degree is given by the number of vertices in the tree, while a morphism belongs to  $\Omega^+$  (resp.  $\Omega^-$ ) when it induces an injection (resp. surjection) between the sets of edges.

For such a tree  $T$ , one can consider the set of planar structures  $p$  on  $T$ . Since every tree in  $\Omega$  carries at least one planar structure, the category  $\Omega$  is equivalent to the category  $\Omega'$  whose objects are planar trees  $(T, p)$ , and whose morphisms  $(T, p) \rightarrow (T', p')$  are the morphisms  $T \rightarrow T'$  in  $\Omega$ . For every such morphism, one can pull back the planar structure  $p'$  on  $T'$  to one on  $T$ , and call the morphism planar if this pulled back structure coincides with  $p$ . The planar morphisms form a wide subcategory of  $\Omega'$ , denoted  $\Omega_{planar}$ ; in this latter category, every automorphism is trivial, and  $\Omega_{planar}$  is equivalent to a strict Reedy category. Every morphism in  $\Omega'$  factors uniquely as an automorphism followed by a planar map. This shows by Proposition 2.5 that the category  $\Omega$  is equivalent to the total category of a crossed group on  $\Omega_{planar}$ .

The embedding  $i : \Delta \hookrightarrow \Omega$  (cf. [30]) is a morphism of generalized Reedy categories, and Theorem 1.6 gives a Reedy model structure on *dendroidal spaces*, which is compatible with the Reedy model structure on simplicial spaces. At the end of Sect. 7 (cf. Example 7.7(iii)), we will show that the model structure on dendroidal spaces is *monoidal* (in the sense of

Hovey [23]) with respect to the Boardman-Vogt tensor product on dendroidal spaces (cf. [30, Appendix]).

Consider a crossed  $\mathbb{R}$ -group  $G$ , and suppose that  $\mathbb{R}$  carries a generalized Reedy structure. We will say that the crossed  $\mathbb{R}$ -group is *compatible with the generalized Reedy structure* if the following two conditions hold:

- (i) the  $G$ -action respects  $\mathbb{R}^+$  and  $\mathbb{R}^-$  (i.e. if  $\alpha : r \rightarrow s$  belongs to  $\mathbb{R}^\pm$  and  $g \in G_s$  then  $g_*(\alpha) : r \rightarrow s$  belongs to  $\mathbb{R}^\pm$ );
- (ii) if  $\alpha : r \rightarrow s$  belongs to  $\mathbb{R}^-$  and  $g \in G_s$  is such that  $\alpha^*(g) = e_r$  and  $g_*(\alpha) = \alpha$ , then  $g = e_s$ .

*Remark 2.9* Observe that condition (i) is in particular satisfied if any morphism in  $\mathbb{R}$ , which in  $\mathbb{R}G$  is isomorphic to a morphism in  $\mathbb{R}^\pm$ , already belongs to  $\mathbb{R}^\pm$ . Condition (ii) is equivalent to the condition that  $\mathbb{R}^-$  fulfills axiom (iv) of Definition 1.1 with respect to special automorphisms of  $\mathbb{R}G$ , cf. the proof of Proposition 2.5.

Because in the simplex category  $\Delta$  the morphisms of  $\Delta^+$  (resp. of  $\Delta^-$ ) are the monomorphisms (resp. split epimorphisms) of  $\Delta$ , any crossed  $\Delta$ -group is compatible with the Reedy structure of  $\Delta$ . The same property holds for crossed groups on  $\Omega_{planar}$ , cf. Example 2.8, and in general for crossed groups on strict EZ-categories, cf. Definition 6.7.

**Proposition 2.10** *Let  $\mathbb{R}$  be a strict Reedy category, and let  $G$  be a compatible crossed  $\mathbb{R}$ -group. Then there is a unique dualizable generalized Reedy structure on  $\mathbb{R}G$  for which the embedding  $\mathbb{R} \hookrightarrow \mathbb{R}G$  is a morphism of generalized Reedy categories.*

*Proof* Necessarily,  $(\mathbb{R}G)^\pm$  consists of those morphisms  $(\alpha, g)$  for which  $\alpha \in \mathbb{R}^\pm$ . Because of compatibility condition (i),  $(\mathbb{R}G)^\pm$  is closed under composition. It is now straightforward to verify that this defines a generalized Reedy structure on  $\mathbb{R}G$ . In particular, axiom (iv) follows from compatibility condition (ii) and the fact that all automorphisms of  $\mathbb{R}G$  are special since  $\mathbb{R}$  is a strict Reedy category. The dual axiom (iv)' holds automatically.

### 3 Kan extensions and the projection formula

In this section we recall some basic facts about Kan extensions for diagram categories. Let  $\phi : \mathbb{D} \rightarrow \mathbb{C}$  be a functor between small categories, and write  $\phi^* : \mathcal{E}^{\mathbb{C}} \rightarrow \mathcal{E}^{\mathbb{D}}$  for precomposition with  $\phi$ . The left and right adjoints of  $\phi^*$  are usually called *left* and *right Kan extension* along  $\phi$ .

If  $\mathcal{E}$  is sufficiently cocomplete, the left Kan extension  $\phi_! : \mathcal{E}^{\mathbb{D}} \rightarrow \mathcal{E}^{\mathbb{C}}$  can be computed pointwise by

$$\phi_!(X)_c = \lim_{\phi/c} X \circ \pi_c$$

where  $\phi/c$  is the comma category with objects  $(d, u : \phi(d) \rightarrow c)$  and morphisms  $(d, u) \rightarrow (d', u')$  given by  $f : d \rightarrow d'$  in  $\mathbb{D}$  such that  $u' \circ \phi(d) = u$ . The functor  $\pi_c : \phi/c \rightarrow \mathbb{D}$  is defined by  $(d, u) \mapsto d$ . We will often informally write

$$\phi_!(X)_c = \lim_{\phi(d) \rightarrow c} X_d.$$

The formula for left Kan extension simplifies if the functor  $\phi : \mathbb{D} \rightarrow \mathbb{C}$  is cofibered. Recall (cf. [7]) that for a given functor  $\phi : \mathbb{D} \rightarrow \mathbb{C}$ , a morphism  $f : d \rightarrow d'$  in  $\mathbb{D}$  is *cocartesian* if

for any  $g : d \rightarrow d''$  such that  $\phi(g) = h\phi(f)$ , there is a unique  $k : d' \rightarrow d''$  such that  $g = kf$  and  $\phi(k) = h$ . The functor  $\phi$  is called *cofibered*, if morphisms in  $\mathbb{C}$  have cocartesian lifts in  $\mathbb{D}$ . If  $\phi$  is cofibered, then for any object  $c$  of  $\mathbb{C}$ , the embedding of the fiber  $\phi^{-1}(c)$  into the comma category  $\phi/c$  (given on objects by  $d \mapsto (d, 1_{\phi(c)})$ ) has a left adjoint, so  $\phi^{-1}(c)$  is cofinal in  $\phi/c$ , and hence

$$\phi_!(X)_c = \lim_{\rightarrow \phi^{-1}(c)} X$$

is the colimit over the fiber. This implies that for any commutative diagram of categories

$$\begin{array}{ccc} \mathbb{D}' & \xrightarrow{\beta} & \mathbb{D} \\ \psi \downarrow & & \downarrow \phi \\ \mathbb{C}' & \xrightarrow{\alpha} & \mathbb{C} \end{array}$$

the natural transformation of functors (induced by adjunction)

$$\psi_! \beta^* \longrightarrow \alpha^* \phi_!$$

is an *isomorphism* whenever the diagram is a pullback with  $\phi$  (and hence  $\psi$ ) cofibered. This is often called the *projection formula*, and will be applied below in the special case where  $\mathbb{D} = \int_{\mathbb{C}} F$  is the Grothendieck construction of a covariant diagram  $F : \mathbb{C} \rightarrow \text{Cat}$ .

Dually, if  $\mathcal{E}$  is sufficiently complete, the right Kan extension  $\phi_* : \mathcal{E}^{\mathbb{D}} \rightarrow \mathcal{E}^{\mathbb{C}}$  can be computed pointwise by

$$\phi_*(X)_c = \lim_{\leftarrow c \rightarrow \phi(d)} X_d$$

and this formula simplifies for fibered functors  $\phi : \mathbb{D} \rightarrow \mathbb{C}$ . Recall that a functor  $\phi$  is called *fibered*, if morphisms in  $\mathbb{C}$  have *cartesian* lifts in  $\mathbb{D}$ . If  $\phi$  is fibered, then for any object  $c$  of  $\mathbb{C}$ , the embedding of the fiber  $\phi^{-1}(c)$  into the comma category  $c/\phi$  (given on objects by  $d \mapsto (1_{\phi(c)}, d)$ ) has a right adjoint, so  $\phi^{-1}(c)$  is final in  $c/\phi$ , and hence

$$\phi_*(X)_c = \lim_{\leftarrow \phi^{-1}(c)} X$$

is the limit over the fiber. This implies that for any commutative diagram of categories

$$\begin{array}{ccc} \mathbb{D}' & \xrightarrow{\beta} & \mathbb{D} \\ \psi \downarrow & & \downarrow \phi \\ \mathbb{C}' & \xrightarrow{\alpha} & \mathbb{C} \end{array}$$

the natural transformation of functors

$$\alpha^* \phi_* \longrightarrow \psi_* \beta^*$$

is an *isomorphism* whenever the diagram is a pullback with  $\phi$  (and hence  $\psi$ ) fibered. This *dual projection formula* will be applied below in the special case where  $\mathbb{D} = \int_{\mathbb{C}} F$  is the Grothendieck construction of a contravariant diagram  $F : \mathbb{C}^{\text{op}} \rightarrow \text{Cat}$ .

### 4 Latching and matching objects

In this section we give an alternative, more global definition of latching and matching objects. Throughout, we consider a fixed generalized Reedy category  $\mathbb{R}$  with wide subcategories  $\mathbb{R}^\pm$  and degree-function  $d$  as in Definition 1.1, and assume that  $\mathcal{E}$  is a sufficiently bicomplete category.

#### 4.1 The groupoids of objects of fixed degree

For each natural number  $n$ , the full subcategory of  $\mathbb{R}$  of objects of degree  $\leq n$  will be denoted  $\mathbb{R}_{\leq n}$ , the full subgroupoid of  $\text{Iso}(\mathbb{R})$  spanned by the objects of degree  $n$  will be denoted  $\mathbb{G}_n(\mathbb{R})$ , and the discrete category of objects of  $\mathbb{R}$  of degree  $n$  will be denoted  $\mathbb{R}_n$ .

#### 4.2 Overcategories

For each natural number  $n$ , the category  $\mathbb{R}^+(n)$  has as objects the non-invertible morphisms  $u : s \rightarrow r$  in  $\mathbb{R}^+$  such that  $d(r) = n$ , and as morphisms from  $u$  to  $u'$  the commutative squares

$$\begin{array}{ccc} s & \xrightarrow{f} & s' \\ u \downarrow & & \downarrow u' \\ r & \xrightarrow{g} & r' \end{array}$$

such that  $f \in \mathbb{R}^+$  and  $g \in \mathbb{G}_n(\mathbb{R})$ .

The wide subcategory  $\mathbb{R}^+(n)$  of  $\mathbb{R}^+(n)$  contains those morphisms for which  $g$  is an identity. The category  $\mathbb{R}^+(r)$  of Sect.1 may thus be identified with the full subcategory of  $\mathbb{R}^+(n)$  spanned by the objects with codomain  $r$ . Notice that

$$\mathbb{R}^+(n) = \coprod_{d(r)=n} \mathbb{R}^+(r).$$

The categories introduced so far assemble into the following commutative diagram:

$$\begin{array}{ccccc} \mathbb{R} & \xleftarrow{d_n} & \mathbb{R}^+(n) & \xrightarrow{c_n} & \mathbb{G}_n(\mathbb{R}) & \xrightarrow{j_n} & \mathbb{R} \\ & & \uparrow k_n & & \uparrow i_n & & \\ & & \mathbb{R}^+(n) & \xrightarrow{b_n} & \mathbb{R}_n & & \end{array}$$

where  $d_n$  denotes the domain-functor,  $b_n$  and  $c_n$  denote codomain-functors, and  $i_n, j_n$  and  $k_n$  are inclusion-functors. Note that  $c_n$  is cofibered, i.e.

$$\mathbb{R}^+(n) \cong \int_{\mathbb{G}_n(\mathbb{R})} \mathbb{R}^+(-),$$

and that the square is a pullback. In particular, the projection formula yields

$$i_n^*(c_n)! \cong (b_n)!k_n^*.$$

#### 4.3 Latching objects

The definition of the latching object  $L_n(X)$  for an object  $X$  of  $\mathcal{E}^{\mathbb{R}}$  now takes the following form:

$$L_n(X) = (c_n)!d_n^*(X) \in \mathcal{E}^{\mathbb{G}_n(\mathbb{R})}.$$

We write  $X_n = j_n^*(X) = X_{|\mathbb{G}_n(\mathbb{R})}$ , so that we get in each degree  $n$  a *latching map*

$$L_n(X) \longrightarrow X_n.$$

Note that, since  $c_n$  is cofibered, we have more concretely:

$$L_n(X)_r = \lim_{\substack{\longrightarrow \\ s \rightarrow r}} X_s,$$

where the colimit is taken over the category  $\mathbb{R}^+(r)$  as in Sect. 1. Accordingly, we will often simplify notation and write  $L_r(X)$  for  $L_n(X)_r$ .

Observe that a morphism  $\phi : \mathbb{S} \rightarrow \mathbb{R}$  of generalized Reedy categories induces for  $k \in \mathbb{N}$  a commutative diagram of functors

$$\begin{array}{ccccc} \mathbb{S} & \xleftarrow{\bar{d}_k} & \mathbb{S}^+((k)) & \xrightarrow{\bar{c}_k} & \mathbb{G}_k(\mathbb{S}) \\ \phi \downarrow & & \phi_k^+ \downarrow & & \downarrow \phi_k \\ \mathbb{R} & \xleftarrow{d_k} & \mathbb{R}^+((k)) & \xrightarrow{c_k} & \mathbb{G}_k(\mathbb{R}) \end{array}$$

and hence a natural transformation

$$(\bar{c}_k)! \phi_k^* \longrightarrow \phi_k^*(c_k)!$$

and in particular for each object  $X$  of  $\mathcal{E}^{\mathbb{R}}$  a natural map

$$(\bar{c}_k)! \bar{d}_k^* \phi^* X = (\bar{c}_k)! \phi_k^* d_k^* X \longrightarrow \phi_k^*(c_k)! d_k^* X,$$

in other words a natural comparison map

$$L_k(\phi^*(X)) \longrightarrow \phi_k^*(L_k(X)).$$

**Lemma 4.4** *Let  $\phi : \mathbb{S} \rightarrow \mathbb{R}$  be a morphism of generalized Reedy categories. Suppose that the induced square*

$$\begin{array}{ccc} \mathbb{S}^+((k)) & \xrightarrow{\bar{c}_k} & \mathbb{G}_k(\mathbb{S}) \\ \phi_k^+ \downarrow & & \downarrow \phi_k \\ \mathbb{R}^+((k)) & \xrightarrow{c_k} & \mathbb{G}_k(\mathbb{R}) \end{array}$$

is a pullback. Then, for each object  $X$  of  $\mathcal{E}^{\mathbb{R}}$ , the natural comparison map of latching objects  $L_k(\phi^*(X)) \rightarrow \phi_k^*(L_k(X))$  is an isomorphism.

The pullback hypothesis holds in particular in the following two cases:

- (i)  $\mathbb{S} = \mathbb{R}^+(n)$  and  $\phi = d_n k_n : \mathbb{R}^+(n) \rightarrow \mathbb{R}$  is the domain functor;
- (ii)  $\mathbb{S} = \mathbb{R}_{\leq n}$  and  $\phi : \mathbb{R}_{\leq n} \rightarrow \mathbb{R}$  is the canonical embedding.

*Proof* The natural comparison map is an isomorphism because  $\bar{c}_k$  and  $c_k$  are cofibered functors, see Sect. 3 and end of Sect. 4.2.

If  $\mathbb{S} = \mathbb{R}^+(n)$  then  $\mathbb{S}^+((k))$  has as objects the composable pairs  $t \rightarrow s \rightarrow r$  of non-invertible, degree-raising maps such that  $d(r) = n$  and  $d(s) = k$ , and as morphisms those transformations of diagrams which are the identity on the last object, an isomorphism on the intermediate object, and degree-raising on the first object; this category coincides with the fiber product of  $\phi_k : \mathbb{G}_k(\mathbb{S}) \rightarrow \mathbb{G}_k(\mathbb{R})$  and  $c_k : \mathbb{R}^+((k)) \rightarrow \mathbb{G}_k(\mathbb{R})$ .

If  $\mathbb{S} = \mathbb{R}_{\leq n}$  the pullback hypothesis follows from the fact that an object of  $\mathbb{R}^+((k))$  belongs to  $\mathbb{S}^+((k))$  if and only if its codomain is of degree  $\leq n$ .

### 4.5 Undercategories

The category  $\mathbb{R}^-(n)$  has as objects the non-invertible morphisms  $u : r \rightarrow s$  in  $\mathbb{R}^-$  such that  $d(r) = n$ , and as morphisms from  $u$  to  $u'$  the commutative squares

$$\begin{array}{ccc} r & \xrightarrow{g} & r' \\ u \downarrow & & \downarrow u' \\ s & \xrightarrow{f} & s' \end{array}$$

such that  $f \in \mathbb{R}^-$  and  $g \in \mathbb{G}_n(\mathbb{R})$ .

The wide subcategory  $\mathbb{R}^-(n)$  of  $\mathbb{R}^-(n)$  contains those morphisms for which  $g$  is an identity. The category  $\mathbb{R}^-(r)$  of Sect. 1 may then be identified with the full subcategory of  $\mathbb{R}^-(n)$  spanned by the objects with domain  $r$ . Notice that

$$\mathbb{R}^-(n) = \coprod_{d(r)=n} \mathbb{R}^-(r).$$

The categories introduced so far assemble into the following commutative diagram:

$$\begin{array}{ccccc} \mathbb{R} & \xleftarrow{\gamma_n} & \mathbb{R}^-(n) & \xrightarrow{\delta_n} & \mathbb{G}_n(\mathbb{R}) & \xrightarrow{j_n} & \mathbb{R} \\ & & \kappa_n \uparrow & & \uparrow i_n & & \\ & & \mathbb{R}^-(n) & \xrightarrow{\beta_n} & \mathbb{R}_n & & \end{array}$$

where  $\gamma_n$  denotes the codomain-functor,  $\beta_n$  and  $\delta_n$  denote domain-functors, and  $i_n, j_n$  and  $\kappa_n$  are inclusion-functors.

Note that  $\delta_n$  is fibered, i.e.

$$\mathbb{R}^-(n) \cong \int_{\mathbb{G}_n(\mathbb{R})} \mathbb{R}^-(\_)$$

and that the square is a pullback. In particular, the dual projection formula yields

$$i_n^*(\delta_n)_* \cong (\beta_n)_* \kappa_n^*.$$

### 4.6 Matching objects

The definition of the *matching object*  $M_n(X)$  of an object  $X$  of  $\mathcal{E}^{\mathbb{R}}$  now takes the following form:

$$M_n(X) = (\delta_n)_* \gamma_n^*(X) \in \mathcal{E}^{\mathbb{G}_n(\mathbb{R})}.$$

We write  $X_n = j_n^*(X) = X|_{\mathbb{G}_n(\mathbb{R})}$ , so that we get in each degree  $n$  a *matching map*

$$X_n \longrightarrow M_n(X).$$

Note that, since  $\delta_n$  is fibered, we have more concretely:

$$M_n(X)_r = \lim_{\substack{\longleftarrow \\ r \rightarrow s}} X_s,$$

where the limit is taken over the category  $\mathbb{R}^-(r)$  as in Sect. 1. Accordingly, we will often simplify notation and write  $M_r(X)$  for  $M_n(X)_r$ .

**Lemma 4.7** *Let  $\phi : \mathbb{S} \rightarrow \mathbb{R}$  be a morphism of generalized Reedy categories. Suppose that the induced square*

$$\begin{CD} \mathbb{S}^-(k) @>>> \mathbb{G}_k(\mathbb{S}) \\ @V\phi_k^-VV @VV\phi_k V \\ \mathbb{R}^-(k) @>>> \mathbb{G}_k(\mathbb{R}) \end{CD}$$

*is a pullback. Then, for each object  $X$  of  $\mathcal{E}^{\mathbb{R}}$ , the natural comparison map of matching objects  $\phi_k^*(M_k(X)) \rightarrow M_k(\phi^*(X))$  is an isomorphism.*

*The pullback hypothesis holds in particular in the following two cases:*

- (i)  $\mathbb{S} = \mathbb{R}^-(n)$  and  $\phi = \gamma_n \kappa_n : \mathbb{R}^-(n) \rightarrow \mathbb{R}$  is the codomain functor;
- (ii)  $\mathbb{S} = \mathbb{R}_{\leq n}$  and  $\phi : \mathbb{R}_{\leq n} \rightarrow \mathbb{R}$  is the canonical embedding.

*Proof* Dual to the proof of Lemma 4.4.

### 5 The Reedy model structure

In this section we prove Theorem 1.6. In particular, we assume throughout that  $\mathbb{R}$  is a generalized Reedy category, and  $\mathcal{E}$  is an  $\mathbb{R}$ -projective Quillen model category. Recall that  $\mathbb{R}$ -projectivity of  $\mathcal{E}$  simply means that for each object  $r$  of  $\mathbb{R}$ , the category  $\mathcal{E}^{\text{Aut}(r)}$  admits a projective model structure. We can reformulate the definition of the classes of maps in Sect. 1 as follows:

**Lemma 5.1** *A map  $X \rightarrow Y$  in  $\mathcal{E}^{\mathbb{R}}$  is a Reedy cofibration (resp. a Reedy weak equivalence, resp. a Reedy fibration) if and only if, for each natural number  $n$ , the map  $X_n \cup_{L_n(X)} L_n(Y) \rightarrow Y_n$  (resp.  $X_n \rightarrow Y_n$ , resp.  $X_n \rightarrow M_n(X) \times_{M_n(Y)} Y_n$ ) is a cofibration (resp. a weak equivalence, resp. a fibration) in  $\mathcal{E}^{\mathbb{G}_n(\mathbb{R})}$ .*

*Proof* This just follows from the equivalence of categories

$$\mathcal{E}^{\mathbb{G}_n(\mathbb{R})} \xrightarrow{\sim} \prod_r \mathcal{E}^{\text{Aut}(r)}$$

where  $r$  runs through a set of representatives for the connected components of the groupoid  $\mathbb{G}_n(\mathbb{R})$ .

A Reedy (co)fibration which is also a Reedy weak equivalence will be referred to as a *trivial Reedy (co)fibration*. The following lemmas are preparatory for the proof of Theorem 1.6.

**Lemma 5.2** *Let  $f : A \rightarrow B$  be a trivial Reedy cofibration; suppose that, for each  $n$ , the induced map  $L_n(f) : L_n(A) \rightarrow L_n(B)$  is a pointwise trivial cofibration (i.e.  $L_n(f)_r$  is a trivial cofibration in  $\mathcal{E}$  for each object  $r$  of  $\mathbb{R}$ ). Then  $f : A \rightarrow B$  has the left lifting property with respect to Reedy fibrations.*

*Proof* Consider a commutative square in  $\mathcal{E}^{\mathbb{R}}$

$$\begin{CD} A @>\alpha>> Y \\ @VfVV @VVgV \\ B @>\beta>> X \end{CD}$$

where  $f$  is a trivial Reedy cofibration and  $g$  is a Reedy fibration, and furthermore  $L_n(f) : L_n(A) \rightarrow L_n(B)$  is a pointwise trivial cofibration for all  $n$ . We construct a diagonal filler  $\gamma : B \rightarrow Y$  by constructing inductively a filler  $\gamma_{\leq n} : B_{\leq n} \rightarrow Y_{\leq n}$  on the full subcategory  $\mathbb{R}_{\leq n}$  of objects of  $\mathbb{R}$  of degree  $\leq n$ . For  $n = 0$ , we get a diagonal filler  $\gamma_0 : B_0 \rightarrow Y_0$  in

$$\begin{array}{ccc} A_0 & \xrightarrow{\alpha_0} & Y_0 \\ f_0 \downarrow & & \downarrow g_0 \\ B_0 & \xrightarrow{\beta_0} & X_0 \end{array}$$

since  $\mathbb{R}_{\leq 0}$  is the groupoid  $\mathbb{G}_0(\mathbb{R})$ , and  $L_0(A) = 0, M_0(X) = 1$ , so that by hypothesis  $f_0$  is a trivial cofibration in  $\mathcal{E}^{\mathbb{G}_0(\mathbb{R})}$  and  $g_0$  is a fibration in  $\mathcal{E}^{\mathbb{G}_0(\mathbb{R})}$ .

Assume by induction that a filler  $\gamma_{\leq n-1} : B_{\leq n-1} \rightarrow Y_{\leq n-1}$  has been found for

$$\begin{array}{ccc} A_{\leq n-1} & \xrightarrow{\alpha_{\leq n-1}} & Y_{\leq n-1} \\ f_{\leq n-1} \downarrow & & \downarrow g_{\leq n-1} \\ B_{\leq n-1} & \xrightarrow{\beta_{\leq n-1}} & X_{\leq n-1}. \end{array}$$

This yields composite maps

$$L_n(B) \longrightarrow L_n(Y) \longrightarrow Y_n \text{ and } B_n \longrightarrow M_n(B) \longrightarrow M_n(Y)$$

as well as a commutative square

$$\begin{array}{ccc} A_n \cup_{L_n(A)} L_n(B) & \longrightarrow & Y_n \\ v_n \downarrow & & \downarrow w_n \\ B_n & \longrightarrow & X_n \times_{M_n(X)} M_n(Y). \end{array}$$

A  $\mathbb{G}_n(\mathbb{R})$ -equivariant filler is exactly what is needed in order to complete the inductive step. To see that such a filler exists, note that by hypothesis  $v_n$  is a cofibration and  $w_n$  is a fibration in  $\mathcal{E}^{\mathbb{G}_n(\mathbb{R})}$ . It is thus enough to check that  $v_n$  is a weak equivalence. For this, consider the following diagram in which the square is a pushout:

$$\begin{array}{ccccc} L_n(A)_r & \longrightarrow & A_r & \xrightarrow{f_r} & B_r \\ L_n(f)_r \downarrow & & \downarrow & \nearrow (v_n)_r & \\ L_n(B)_r & \longrightarrow & (A \cup_{L_n(A)} L_n(B))_r & & \end{array}$$

Since, by hypothesis,  $L_n(f)_r$  is a trivial cofibration in  $\mathcal{E}$ , and  $f_r$  is a weak equivalence,  $v_n$  is a weak equivalence as required.

**Lemma 5.3** *Let  $f : A \rightarrow B$  be a Reedy cofibration such that  $f_r : A_r \rightarrow B_r$  is a weak equivalence for all objects  $r$  of  $\mathbb{R}$  of degree  $< n$ . Then, the induced map  $L_n(f) : L_n(A) \rightarrow L_n(B)$  is a pointwise trivial cofibration.*

*Proof* For  $n = 0$ , there is nothing to prove; therefore, we can assume inductively that  $L_k(f) : L_k(A) \rightarrow L_k(B)$  is a pointwise trivial cofibration for  $k < n$ . We want to show that  $i_n^* L_n(f)$  is a trivial cofibration in  $\mathcal{E}^{\mathbb{R}^n}$ . To this end, we have to find a filler for any commutative square

$$\begin{array}{ccc} i_n^* L_n(A) & \longrightarrow & Y \\ i_n^* L_n(f) \downarrow & & \downarrow g \\ i_n^* L_n(B) & \longrightarrow & X \end{array}$$



in  $\mathcal{E}^{\mathbb{R}_n}$  in which  $g : Y \rightarrow X$  is a fibration. Since  $i_n^* L_n = i_n^*(c_n)!d_n^* = (b_n)!k_n^*d_n^*$ , a filler for the former square is the same as a filler for the following square in  $\mathcal{E}^{\mathbb{R}^+(n)}$ :

$$\begin{array}{ccc} k_n^*d_n^*(A) & \longrightarrow & b_n^*(Y) \\ k_n^*d_n^*(f) \downarrow & & \downarrow b_n^*(g) \\ k_n^*d_n^*(B) & \longrightarrow & b_n^*(X). \end{array}$$

In order to finish the proof, we shall apply Lemma 5.2 to this square. The category  $\mathbb{S} = \mathbb{R}^+(n)$  is a generalized Reedy category for which  $\mathbb{S} = \mathbb{S}^+$ . In particular, Reedy fibrations are the same as pointwise fibrations, so  $b_n^*(g)$  is a Reedy fibration. Moreover,  $k_n^*d_n^*(f)$  is a Reedy weak equivalence in  $\mathcal{E}^{\mathbb{S}}$ , since the objects of  $\mathbb{S}$  have degree  $< n$ . It remains to be shown that  $k_n^*d_n^*(f)$  is a Reedy cofibration whose induced maps on latching objects of degree  $< n$  are pointwise trivial cofibrations.

Write  $\phi = d_n k_n$ . By Lemma 4.4, the functor  $\phi_k^* : \mathcal{E}^{\mathbb{G}_k(\mathbb{R})} \rightarrow \mathcal{E}^{\mathbb{G}_k(\mathbb{S})}$  induces a canonical isomorphism  $L_k(\phi^*(A)) \cong \phi_k^*(L_k(A))$ . Therefore, the relative latching map  $\phi^*(A) \cup_{L_k(\phi^*(A))} L_k(\phi^*(B)) \rightarrow \phi^*(B)$  may be identified with  $\phi_k^*$  of the relative latching map  $A_k \cup_{L_k(A)} L_k(B) \rightarrow B_k$ . Observe that  $\phi_k : \mathbb{G}_k(\mathbb{S}) \rightarrow \mathbb{G}_k(\mathbb{R})$  is a faithful functor between groupoids, so  $\phi_k^*$  preserves cofibrations, thus  $k_n^*d_n^*(f)$  is a Reedy cofibration. Moreover,  $L_k(\phi^*(A)) \rightarrow L_k(\phi^*(B))$  is a pointwise trivial cofibration for  $k < n$ , since  $L_k(A) \rightarrow L_k(B)$  is so by induction hypothesis.

**Lemma 5.4** *Let  $g : Y \rightarrow X$  be a trivial Reedy fibration; suppose that for each  $n$ , the induced map  $M_n(g) : M_n(Y) \rightarrow M_n(X)$  is a (pointwise) trivial fibration. Then  $g : Y \rightarrow X$  has the right lifting property with respect to Reedy cofibrations.*

*Proof* Dual to the proof of Lemma 5.2.

**Lemma 5.5** *Let  $g : Y \rightarrow X$  be a Reedy fibration such that  $g_r : Y_r \rightarrow X_r$  is a weak equivalence for all objects  $r$  of  $\mathbb{R}$  of degree  $< n$ . Then, the induced map  $M_n(g) : M_n(Y) \rightarrow M_n(X)$  is a (pointwise) trivial fibration.*

*Proof* For  $n = 0$ , there is nothing to prove; therefore, we can assume inductively that  $M_k(g) : M_k(Y) \rightarrow M_k(X)$  is a trivial fibration for  $k < n$ . We want to show that  $i_n^* M_n(g)$  is a trivial fibration in  $\mathcal{E}^{\mathbb{R}_n}$ . To this end, we have to find a filler for any commutative square

$$\begin{array}{ccc} A & \longrightarrow & i_n^* M_n(Y) \\ f \downarrow & & \downarrow i_n^* M_n(g) \\ B & \longrightarrow & i_n^* M_n(X) \end{array}$$

in  $\mathcal{E}^{\mathbb{R}_n}$  in which  $f : A \rightarrow B$  is a cofibration. Since  $i_n^* M_n = i_n^*(\delta_n)_* \gamma_n^* = (\beta_n)_* \kappa_n^* \gamma_n^*$ , a filler for the former square is the same as a filler for the following square in  $\mathcal{E}^{\mathbb{R}^-(n)}$ :

$$\begin{array}{ccc} \beta_n^*(A) & \longrightarrow & \kappa_n^* \gamma_n^*(Y) \\ \beta_n^*(f) \downarrow & & \downarrow \kappa_n^* \gamma_n^*(g) \\ \beta_n^*(B) & \longrightarrow & \kappa_n^* \gamma_n^*(X). \end{array}$$

In order to finish the proof, we shall apply Lemma 5.4 to this square. The category  $\mathbb{S} = \mathbb{R}^-(n)$  is a generalized Reedy category for which  $\mathbb{S} = \mathbb{S}^-$ ; notice that  $\mathbb{S}$  has no non-trivial automorphisms in virtue of axiom (iv) of Definition 1.1; in other words,  $\mathbb{S}$  is equivalent to a strict Reedy category. Therefore, Reedy cofibrations are the same as pointwise cofibrations, so  $\beta_n^*(f)$  is a Reedy cofibration. Moreover,  $\kappa_n^* \gamma_n^*(f)$  is a Reedy weak equivalence in  $\mathcal{E}^{\mathbb{S}}$ , since

the objects of  $\mathbb{S}$  have degree  $< n$ . It remains to be shown that  $\kappa_n^* \gamma_n^*(f)$  is a Reedy fibration whose induced maps on matching objects of degree  $< n$  are trivial fibrations.

Write  $\phi = \gamma_n \kappa_n$ . By Lemma 4.7, the functor  $\phi_k^* : \mathcal{E}^{\mathbb{G}_k(\mathbb{R})} \rightarrow \mathcal{E}^{\mathbb{G}_k(\mathbb{S})}$  induces a canonical isomorphism  $\phi_k^*(M_k(Y)) \cong M_k(\phi^*(X))$ . Therefore, the relative matching map  $\phi^*(Y) \rightarrow \phi^*(X) \times_{M_k(\phi^*(X))} M_k(\phi^*(Y))$  may be identified with  $\phi_k^*$  of the relative matching map  $Y_k \rightarrow X_k \times_{M_k(X)} M_k(Y)$ . Observe that  $\phi_k^*$  preserves fibrations, thus  $\kappa_n^* \gamma_n^*(g)$  is a Reedy fibration. Moreover,  $M_k(\phi^*(Y)) \rightarrow M_k(\phi^*(X))$  is a trivial fibration for  $k < n$ , since  $M_k(Y) \rightarrow M_k(X)$  is so by induction hypothesis.

*Remark 5.6* The reader may observe that the proofs of Lemmas 5.3 and 5.5, although very similar, are not exactly dual to each other. In particular, while the fact that  $b_n^*(g)$  is a Reedy fibration does not use axiom (iv) of a generalized Reedy category, the proof that  $\beta_n^*(f)$  is a Reedy cofibration does. On the other hand, while the preservation under  $\phi_k^*$  of fibrations follows from our use of projective model structures, the preservation of cofibrations relies on the faithfulness of  $\phi_k$ .

*Proof of Theorem 1.6* Limits and colimits in  $\mathcal{E}^{\mathbb{R}}$  are constructed pointwise. The class of Reedy weak equivalences has the two-out-of-three property. Moreover, all three classes are closed under retract. It remains to be shown that the lifting and factorization axioms of a Quillen model category hold.

For the lifting axiom, observe that by Lemma 5.3, trivial Reedy cofibrations fulfill the hypothesis of Lemma 5.2, and therefore have the left lifting property with respect to Reedy fibrations. Dually, Lemmas 5.5 and 5.4 imply that trivial Reedy fibrations have the right lifting property with respect to Reedy cofibrations.

We now come to the factorization axiom. Given a map  $f : X \rightarrow Y$  in  $\mathcal{E}^{\mathbb{R}}$ , we shall construct inductively a factorization  $X \rightarrow A \rightarrow Y$  of  $f$  into a trivial Reedy cofibration followed by a Reedy fibration. To do so, we use that  $\mathcal{E}$  is  $\mathbb{R}$ -projective by hypothesis so that for each  $n$ , the category  $\mathcal{E}^{\mathbb{G}_n(\mathbb{R})}$  admits a projective model structure; cf. the proof of Lemma 5.1.

For  $n = 0$ , factor  $f_0$  in  $\mathcal{E}^{\mathbb{G}_0(\mathbb{R})}$  as  $X_0 \rightarrow A_0 \rightarrow Y_0$  into a trivial cofibration followed by a fibration. Next, if  $X_{\leq n-1} \rightarrow A_{\leq n-1} \rightarrow Y_{\leq n-1}$  is a factorization of  $f_{\leq n-1}$  into trivial Reedy cofibration followed by Reedy fibration in  $\mathcal{E}^{\mathbb{R}_{\leq n-1}}$ , we obtain the following commutative diagram in  $\mathcal{E}^{\mathbb{G}_n(\mathbb{R})}$  :

$$\begin{array}{ccccc}
 L_n(X) & \rightarrow & L_n(A) & \rightarrow & L_n(Y) \\
 \downarrow & & & & \downarrow \\
 X_n & & & & Y_n \\
 \downarrow & & & & \downarrow \\
 M_n(X) & \rightarrow & M_n(A) & \rightarrow & M_n(Y).
 \end{array}$$

This diagram induces a map  $X_n \cup_{L_n(X)} L_n(A) \rightarrow M_n(A) \times_{M_n(Y)} Y_n$  which we factor as a trivial cofibration followed by a fibration in  $\mathcal{E}^{\mathbb{G}_n(\mathbb{R})}$ :

$$X_n \cup_{L_n(X)} L_n(A) \xrightarrow{\sim} A_n \rightarrow M_n(A) \times_{M_n(Y)} Y_n.$$

The object  $A_n$  of  $\mathcal{E}^{\mathbb{G}_n(\mathbb{R})}$  together with the maps  $L_n(A) \rightarrow A_n \rightarrow M_n(A)$  define an extension of  $A_{\leq n-1}$  to an object  $A_{\leq n}$  in  $\mathcal{E}^{\mathbb{R}_{\leq n}}$  together with a factorization of  $f_{\leq n} : X_{\leq n} \rightarrow Y_{\leq n}$  into a Reedy cofibration  $X_{\leq n} \rightarrow A_{\leq n}$  followed by a Reedy fibration  $A_{\leq n} \rightarrow Y_{\leq n}$ . The former map is a *trivial* Reedy cofibration, because the map  $X_n \rightarrow A_n$  decomposes into two maps  $X_n \rightarrow X_n \cup_{L_n(X)} L_n(A) \rightarrow A_n$ , the first one of which is a weak equivalence by Lemma 5.3, the second one by construction. This defines the required factorization of  $f_{\leq n}$  in  $\mathcal{E}^{\mathbb{R}_{\leq n}}$ .

The factorization of  $f$  into a Reedy cofibration followed by a trivial Reedy fibration is constructed in a dual manner using Lemma 5.5 instead of Lemma 5.3.

The proof of Theorem 1.6 uses implicitly that trivial Reedy (co)fibrations may be characterized in terms of relative matching (latching) maps. Since this is a pivotal property of the Reedy model structure, we state it explicitly:

**Proposition 5.7** *A map  $f : A \rightarrow B$  in  $\mathcal{E}^{\mathbb{R}}$  is a trivial Reedy cofibration if and only if, for each  $n$ , the relative latching map  $A_n \cup_{L_n(A)} L_n(B) \rightarrow B_n$  is a trivial cofibration in  $\mathcal{E}^{\mathbb{G}_n(\mathbb{R})}$ .*

*A map  $g : Y \rightarrow X$  in  $\mathcal{E}^{\mathbb{R}}$  is a trivial Reedy fibration if and only if, for each  $n$ , the relative matching map  $Y_n \rightarrow X_n \times_{M_n(X)} M_n(Y)$  is a trivial fibration in  $\mathcal{E}^{\mathbb{G}_n(\mathbb{R})}$ .*

*Proof* For each  $n$ , the induced map  $f_n : A_n \rightarrow B_n$  in  $\mathcal{E}^{\mathbb{G}_n(\mathbb{R})}$  factors as

$$A_n \xrightarrow{u_n} A_n \cup_{L_n(A)} L_n(B) \xrightarrow{v_n} B_n.$$

If  $f$  is a trivial Reedy cofibration then  $f_n$  is a weak equivalence, so that, by Lemma 5.3,  $u_n$  is a weak equivalence, and hence  $v_n$  is a trivial cofibration. Conversely, if each  $v_n$  is a trivial cofibration then an induction on  $n$ , based on Lemma 5.3, shows that  $u_n$  is a weak equivalence, and hence  $f$  is a trivial Reedy cofibration.

The dual proof for a trivial Reedy fibration  $g : Y \rightarrow X$  uses Lemma 5.5 instead of Lemma 5.3.

### 6 Skeleta and coskeleta

In this section we define the skeletal filtration and the coskeletal tower of any functor  $X : \mathbb{R} \rightarrow \mathcal{E}$  on a generalized Reedy category  $\mathbb{R}$ , and study their interaction with the Reedy model structure on  $\mathcal{E}^{\mathbb{R}}$  for a Quillen model category  $\mathcal{E}$ . We then introduce a special class of dualizable generalized Reedy categories for which the skeleta in  $\text{Sets}^{\mathbb{R}^{\text{op}}}$  are simple to describe.

Recall that for any object  $X$  of  $\mathcal{E}^{\mathbb{R}}$ , the restriction  $j_n^* X : \mathbb{G}_n(\mathbb{R}) \rightarrow \mathcal{E}$  along the inclusion  $j_n : \mathbb{G}_n(\mathbb{R}) \rightarrow \mathbb{R}$  is denoted  $X_n$ . We shall write  $t_n : \mathbb{R}_{\leq n} \hookrightarrow \mathbb{R}$  for the full embedding of the subcategory of objects of degree  $\leq n$ , cf. Sect. 4.1.

**Definition 6.1** The *n-skeleton functor* is the endofunctor  $sk_n = t_n!t_n^*$ . The *n-coskeleton functor* is the endofunctor  $cosk_n = t_n^*t_n^*$ .

Since  $t_n : \mathbb{R}_{\leq n} \hookrightarrow \mathbb{R}$  is a full embedding, the unit of the  $(t_n!, t_n^*)$ -adjunction (resp. the counit of the  $(t_n^*, t_n^*)$ -adjunction) is an isomorphism; in other words, the endofunctor  $sk_n$  (resp.  $cosk_n$ ) is an idempotent comonad (resp. monad) on  $\mathcal{E}^{\mathbb{R}}$ .

The counit of the  $(t_n!, t_n^*)$ -adjunction (resp. unit of the  $(t_n^*, t_n^*)$ -adjunction) provides for each object  $X$  of  $\mathcal{E}^{\mathbb{R}}$  a map  $sk_n(X) \rightarrow X$  (resp.  $X \rightarrow cosk_n(X)$ ) in  $\mathcal{E}^{\mathbb{R}}$ . Observe however that these maps need not be monic (resp. epic) for general  $X$ .

For consistency,  $sk_{-1}(X)$  (resp.  $cosk_{-1}(X)$ ) will denote an initial (resp. terminal) object of  $\mathcal{E}^{\mathbb{R}}$ .

**Lemma 6.2** *For each object  $X$  of  $\mathcal{E}^{\mathbb{R}}$ , the n-th latching object  $L_n(X)$  is canonically isomorphic to  $sk_{n-1}(X)_n$ , and the n-th matching object  $M_n(X)$  is canonically isomorphic to  $cosk_{n-1}(X)_n$ .*

*Under these isomorphisms, the n-th latching map  $L_n(X) \rightarrow X_n$  is induced by the counit  $sk_{n-1}(X) \rightarrow X$ , and the n-th matching map  $X_n \rightarrow M_n(X)$  is induced by the unit  $X \rightarrow cosk_{n-1}(X)$ .*

*Proof* This follows from the explicit formulas for the left and right Kan extensions  $t_{n!}$  and  $t_{n*}$  (cf. Sect.3), and from axiom (iii) of Definition 1.1. Indeed, the latter implies that for any object  $r$  of  $\mathbb{R}$ , the category  $\mathbb{R}^+(r)$  is cofinal in the comma category  $\mathbb{R}_{\leq n}/r$ , while the category  $\mathbb{R}^-(r)$  is final in the comma category  $r/\mathbb{R}_{\leq n}$ . Moreover, the latching map  $L_n(X) \rightarrow X_n$  of Sect. 4.3 factors canonically through the counit  $sk_{n-1}(X)_n \rightarrow X_n$ , while the matching map  $X_n \rightarrow M_n(X)$  of Sect. 4.6 factors canonically through the unit  $X_n \rightarrow cosk_{n-1}(X)_n$ .

**Lemma 6.3** *For any natural numbers  $m \leq n$ , there are canonical isomorphisms  $sk_n \circ sk_m \cong sk_m \cong sk_m \circ sk_n$  as well as  $cosk_n \circ cosk_m \cong cosk_m \cong cosk_m \circ cosk_n$ .*

*Proof* Write  $\alpha : \mathbb{R}_{\leq m} \rightarrow \mathbb{R}_{\leq n}$  for the full embedding. It induces a pair of adjoint functors  $(\alpha_!, \alpha^*)$ . Since by definition  $t_n \alpha = t_m$ , we get as required

$$\begin{aligned} sk_n sk_m &= t_n! t_n^* t_m! t_m^* \\ &= t_n! t_n^* t_n! \alpha_! \alpha^* t_n^* \\ &\cong t_n! \alpha_! \alpha^* t_n^* \\ &= t_m! t_m^* \\ &= sk_m, \end{aligned}$$

where the isomorphism is induced by the unit  $id \xrightarrow{\sim} t_n^* t_n!$  of the monad  $t_n$ . The other cases are equally straightforward.

Lemma 6.3 yields for each object  $X$  of  $\mathcal{E}^{\mathbb{R}}$  natural systems of maps

$$\begin{aligned} sk_0 X &\longrightarrow sk_1 X \longrightarrow sk_2 X \longrightarrow \dots, \\ cosk_0 X &\longleftarrow cosk_1 X \longleftarrow cosk_2 X \longleftarrow \dots. \end{aligned}$$

The colimit of the first sequence and the limit of the second sequence are both isomorphic to  $X$ . We now discuss for which objects  $X$  of  $\mathcal{E}^{\mathbb{R}}$ , this defines a skeletal filtration (resp. coskeletal tower), i.e. for which  $X$  these maps are cofibrations (resp. fibrations) in  $\mathcal{E}^{\mathbb{R}}$ . Recall that a functor between Quillen model categories is called a *left* (resp. *right*) *Quillen functor* if it preserves cofibrations and trivial cofibrations (resp. fibrations and trivial fibrations).

**Lemma 6.4** *Let  $\mathcal{E}$  be a Quillen model category and let  $\mathbb{R}$  be a generalized Reedy category. We endow  $\mathcal{E}^{\mathbb{R}}$  and  $\mathcal{E}^{\mathbb{R}_{\leq n}}$  with their Reedy model structures. Then,*

- (i) *the left Kan extension  $t_{n!}$  is a left Quillen functor;*
- (ii) *the right Kan extension  $t_{n*}$  is a right Quillen functor;*
- (iii) *the restriction functor  $t_n^*$  is simultaneously a left and right Quillen functor.*

*In particular,  $sk_n$  (resp.  $cosk_n$ ) is a left (resp. right) Quillen endofunctor of  $\mathcal{E}^{\mathbb{R}}$ .*

*Proof* By adjointness, (iii) is equivalent to the conjunction of (i) and (ii). Under the identification  $\mathbb{G}_k(\mathbb{R}_{\leq n}) \cong \mathbb{G}_k(\mathbb{R})$ , Lemma 4.4(ii) gives for each object  $X$  of  $\mathcal{E}^{\mathbb{R}}$  and each  $n$ , an isomorphism  $L_k(t_n^* X) \cong L_k(X)$ . It follows then from the definition of a Reedy cofibration and from the characterization 5.7 of trivial Reedy cofibrations that  $t_n^*$  preserves both classes, and is hence a left Quillen functor. The proof that  $t_n^*$  is also a right Quillen functor follows similarly using Lemma 4.7(ii).

**Proposition 6.5** *Let  $\mathcal{E}$  be a Quillen model category and let  $\mathbb{R}$  be a generalized Reedy category. For any  $m < n \leq \infty$ , and any object  $X$  of  $\mathcal{E}^{\mathbb{R}}$ ,*

- (i) if  $X$  is Reedy cofibrant, the canonical map  $sk_m(X) \rightarrow sk_n(X)$  is a Reedy cofibration between Reedy cofibrant objects;
- (ii) if  $X$  is Reedy fibrant, the canonical map  $cosk_n(X) \rightarrow cosk_m(X)$  is a Reedy fibration between Reedy fibrant objects.

*Proof* The proofs of (i) and (ii) are dual; we shall establish (i). By Lemma 6.3, we can stick to the case  $n = \infty$ , i.e. to the case  $sk_m(X) \rightarrow sk_\infty(X) = X$ . For this, consider the commutative square:

$$\begin{array}{ccc} L_k(sk_m(X)) & \rightarrow & L_k(X) \\ \downarrow & & \downarrow \\ sk_m(X)_k & \longrightarrow & X_k. \end{array}$$

For  $k \leq m$ , the horizontal maps are isomorphisms, thus the relative latching map  $sk_m(X)_k \cup_{L_k(sk_m(X))} L_k(X) \rightarrow X_k$  is an isomorphism too. For  $k > m$ , the left vertical map is an isomorphism by Lemmas 6.2 and 6.3, thus the relative latching map coincides with  $L_k(X) \rightarrow X_k$  which is a cofibration by hypothesis. Moreover, Lemma 6.4 shows that  $sk_m(X)$  is Reedy cofibrant.

We shall now introduce a special class of generalized Reedy categories  $\mathbb{R}$  for which the skeletal filtration in  $\text{Sets}^{\mathbb{R}^{\text{op}}}$  admits a particularly simple description, as in Corollary 6.10 below. In the particular case of the simplex category  $\Delta$ , this proposition was first observed by Eilenberg and Zilber (see [16, 19]), and therefore we have chosen to name these special categories *Eilenberg–Zilber categories*, or briefly *EZ-categories*. Their formal definition makes use of the notion of absolute pushout which we recall first:

*Remark 6.6* A pushout in a category  $\mathbb{R}$  is called *absolute* if it is preserved by the Yoneda embedding  $\mathbb{R} \hookrightarrow \widehat{\mathbb{R}} = \text{Sets}^{\mathbb{R}^{\text{op}}}$ . (It can be shown that this is equivalent to being preserved by any functor, but we will not use this fact.) We will be concerned with absolute pushouts of two split epimorphisms (i.e. epimorphisms having sections). Notice that the image of a map  $f : r \rightarrow s$  in  $\mathbb{R}$  under the Yoneda embedding is an epimorphism in  $\widehat{\mathbb{R}}$  if and only if  $f$  is a split epimorphism. Since pushouts of epimorphisms are epimorphisms, this shows that an absolute pushout of split epimorphisms yields in fact a commutative square of split epimorphisms.

**Definition 6.7** An *EZ-category* is a small category  $\mathbb{R}$ , equipped with a degree-function  $d : \text{Ob}(\mathbb{R}) \rightarrow \mathbb{N}$ , such that

- (i) monomorphisms preserve (resp. raise) the degree if and only if they are invertible (resp. non-invertible);
- (ii) every morphism factors as a split epimorphism followed by a monomorphism;
- (iii) any pair of split epimorphisms with common domain has an absolute pushout.

Observe that assuming axiom (ii), axiom (i) implies

- (i)' split epimorphisms preserve (resp. lower) the degree if and only if they are invertible (resp. non-invertible).

Any EZ-category is a *dualizable* generalized Reedy category where  $\mathbb{R}^+$  (resp.  $\mathbb{R}^-$ ) is defined to be the wide subcategory containing all monomorphisms (resp. split epimorphisms). Notice however that, although the dual of an EZ-category is a generalized Reedy category, it is in general *not* an EZ-category. We are mostly interested in *presheaves* on  $\mathbb{R}$ , so that the

reader should be aware of the fact that the roles of  $\mathbb{R}^\pm$  have to be reversed in the definitions of Sects. 4–6. Axiom (ii) expresses that the epi-mono factorization system of the presheaf category  $\widehat{\mathbb{R}}$  restricts (under the Yoneda-embedding) to  $\mathbb{R}$ , while axiom (iii) can be restated as follows: in  $\widehat{\mathbb{R}}$ , the pushout of any pair of representable epimorphisms with common domain is representable.

*Examples 6.8* As important examples of EZ-categories we mention the simplex category  $\Delta$ , Segal’s category  $\Gamma$ , the category  $\Delta_{sym}$  (cf. 1.9.a–c), as well as the total category  $\mathbb{R}G$  of a crossed group  $G$  on an EZ-category  $\mathbb{R}$  whose underlying Reedy category is strict (e.g. the category  $\Lambda$  for cyclic sets, resp. the category  $\Omega$  for dendroidal sets, cf. 2.7 and 2.8). To see that axiom (iii) of an EZ-category holds for  $\mathbb{R}G$ , one can either construct the pushout of split epimorphisms in  $\mathbb{R}G$  explicitly from such pushouts in  $\mathbb{R}$ , or use the following more abstract argument: the inclusion functor  $i : \mathbb{R} \hookrightarrow \mathbb{R}G$  induces adjoint functors  $i_! : \widehat{\mathbb{R}} \rightleftarrows \widehat{\mathbb{R}G} : i^*$ . Because  $i$  is bijective on objects, the restriction functor  $i^*$  is faithful. This implies that the category  $\widehat{\mathbb{R}G}$  is equivalent to the category of algebras for the monad  $i^*i_!$  on  $\widehat{\mathbb{R}}$ . Therefore, the restriction functor  $i^*$  creates absolute pushouts.

The presheaf  $\mathbb{R}(-, r)$  represented by an object  $r$  of  $\mathbb{R}$  will be denoted  $\mathbb{R}[r]$ . The split epimorphisms of an EZ-category will be called *degeneracy operators*; the monomorphisms will be called *face operators*. Recall that the Yoneda-lemma allows us to identify *elements* of a set-valued presheaf  $X$  on  $\mathbb{R}$  with *maps*  $x : \mathbb{R}[r] \rightarrow X$  in  $\widehat{\mathbb{R}}$ ; such a map (or element)  $x$  will be called *degenerate* if  $x$  factors through a non-invertible degeneracy  $\mathbb{R}[r] \rightarrow \mathbb{R}[s]$ , and *non-degenerate* otherwise.

**Proposition 6.9** *Let  $\mathbb{R}$  be an EZ-category and let  $X$  be a presheaf on  $\mathbb{R}$ . Then any element  $x : \mathbb{R}[r] \rightarrow X$  factors in an essentially unique way as a degeneracy  $\rho_x : \mathbb{R}[r] \rightarrow \mathbb{R}[s]$  followed by a non-degenerate element  $\sigma_x : \mathbb{R}[s] \rightarrow X$ .*

Any such decomposition will be referred to as a *standard decomposition* of  $x$ .

*Proof* The existence of a standard decomposition follows from the facts that the degree-function takes values in  $\mathbb{N}$ , and that non-invertible degeneracies lower the degree by 6.7(i)’. For the essential uniqueness, observe first that there can be at most one comparison map from a standard decomposition  $x = \sigma_x \rho_x$  to another  $x = \sigma'_x \rho'_x$ , since degeneracies are epic. It remains to be shown that such a comparison map always exists. Take the absolute pushout of  $\rho_x$  and  $\rho'_x$ , as provided by 6.7(iii):

$$\begin{array}{ccc} \mathbb{R}[r] & \xrightarrow{\rho_x} & \mathbb{R}[s] \\ \rho'_x \downarrow & & \downarrow \tau_x \\ \mathbb{R}[s'] & \xrightarrow{\tau'_x} & \mathbb{R}[t] \end{array}$$

There exists therefore a map  $\phi_x : \mathbb{R}[t] \rightarrow X$  such that  $\phi_x \tau_x = \sigma_x$  and  $\phi_x \tau'_x = \sigma'_x$ . Since  $\sigma_x$  and  $\sigma'_x$  are non-degenerate, the split epimorphisms  $\tau_x$  and  $\tau'_x$  must preserve the degree. It then follows from 6.7(i)’ that  $\tau_x$  and  $\tau'_x$  are invertible so that  $(\tau'_x)^{-1} \tau_x$  provides the required comparison map.

**Corollary 6.10** *Let  $\mathbb{R}$  be an EZ-category and let  $X$  be a presheaf on  $\mathbb{R}$ . Then the counit  $sk_n(X) \rightarrow X$  is monic and its image is the subobject  $X^{(n)}$  of those elements of  $X$  which factor through an element  $\mathbb{R}[s] \rightarrow X$  of degree  $s \leq n$ .*

*Proof* Notice that the counit  $sk_n(X) \rightarrow X$  factors through  $X^{(n)}$  since by definition, for each object  $r$  of  $\mathbb{R}$ , we have

$$sk_n(X)_r = t_n!t_n^*(X)_r = \lim_{r \rightarrow s, d(s) \leq n} X_s,$$

where the last equation follows from the pointwise formula for the left Kan extension, cf. Sect. 3. The induced map  $sk_n(X) \rightarrow X^{(n)}$  is pointwise surjective. To see that  $sk_n(X) \rightarrow X^{(n)}$  is pointwise injective, take two elements  $x, y$  in  $sk_n(X)$  giving rise to the same element  $z$  in  $X^{(n)}$ . Then, the essential uniqueness of the standard decomposition of  $z$  shows that  $x$  and  $y$  define the same element in  $sk_n(X)$ .

### 7 Monoidal Reedy model structures

From now on, we shall assume that  $\mathcal{E} = (\mathcal{E}, \otimes_{\mathcal{E}}, I_{\mathcal{E}}, \tau_{\mathcal{E}})$  is a *closed symmetric monoidal category*, see e.g. [7]. Observe that if arbitrary (small) coproducts of the unit object  $I_{\mathcal{E}}$  exist, there is a canonical functor  $I_{\mathcal{E}} : \text{Sets} \rightarrow \mathcal{E}$  given by  $S \mapsto I_{\mathcal{E}}[S] = \coprod_S I_{\mathcal{E}}$ . The symmetric monoidal structure will be called *solid* if these coproducts exist, and if moreover the resulting functor from the category of sets to  $\mathcal{E}$  is *faithful*. Objects and morphisms of  $\mathcal{E}$  which belong to the essential image of this functor will be called *discrete*. Likewise, the presheaf category  $\widehat{\mathbb{R}}$  maps to  $\mathcal{E}^{\text{Rop}}$ . Observe that both functors have right adjoints, and hence preserve colimits.

Recall that, according to Hovey [23], a *monoidal model category* is a category which is simultaneously a closed symmetric monoidal category and a Quillen model category such that *unit* and *pushout-product* axioms hold. For brevity, we shall say that a monoidal model category  $\mathcal{E}$  is *solid* if

- (i) the symmetric monoidal structure is *solid* in the sense mentioned above;
- (ii) the unit  $I_{\mathcal{E}}$  is *cofibrant*;
- (iii) for any discrete group  $G$ , and any map of  $G$ -sets  $f : A \rightarrow B$ , if  $I_{\mathcal{E}}[f]$  is a cofibration in  $\mathcal{E}^G$  then  $f$  is a *free  $G$ -extension*.<sup>2</sup>

Observe that condition (ii) makes the unit axiom redundant, and condition (iii) (applied to the trivial group) implies that discrete cofibrations in  $\mathcal{E}$  are monic. Condition (ii) implies that the converse of (iii) also holds. Examples of solid monoidal model categories include the category of compactly generated spaces, the category of simplicial sets (both equipped with Quillen’s model structure), and the category of differential graded  $R$ -modules with the projective model structure.

#### 7.1 Boundary inclusions and cofibrations

For each object  $r$  of an EZ-category  $\mathbb{R}$ , the *formal boundary*  $\partial\mathbb{R}[r]$  of  $\mathbb{R}[r]$  is defined to be the subobject of those elements of  $\mathbb{R}[r]$  which factor through a non-invertible face operator  $s \rightarrow r$ . By Corollary 6.10, we have  $\partial\mathbb{R}[r] = sk_{d(r)-1}\mathbb{R}[r]$ . Our main purpose here is to single out a class of maps in  $\widehat{\mathbb{R}}$  which induce *Reedy cofibrations* in  $\mathcal{E}^{\text{Rop}}$  for *any* solid monoidal model category  $\mathcal{E}$ . This class coincides with Cisinski’s class of *normal monomorphisms*, see [10, 8.1.30].

<sup>2</sup> A  $G$ -equivariant map of  $G$ -sets  $f : A \rightarrow B$  is a *free  $G$ -extension* iff  $f$  is monic and  $G$  acts freely on the complement  $B \setminus f(A)$ .

**Proposition 7.2** For a map  $\phi : X \rightarrow Y$  of set-valued presheaves on an EZ-category  $\mathbb{R}$ , the following three properties are equivalent:

- (i) for each object  $r$  of  $\mathbb{R}$ , the relative latching map  $X_r \cup_{L_r(X)} L_r(Y) \rightarrow Y_r$  is a free  $\text{Aut}(r)$ -extension;
- (ii)  $\phi$  is monic, and for each object  $r$  of  $\mathbb{R}$  and each non-degenerate element  $y \in Y_r \setminus \phi(X)_r$ , the isotropy group  $\{g \in \text{Aut}(r) \mid g^*(y) = y\}$  is trivial;
- (iii) for each  $n \geq 0$ , the relative  $n$ -skeleton  $sk_n(\phi) = X \cup_{sk_n(X)} sk_n(Y)$  is obtained from the relative  $(n - 1)$ -skeleton  $sk_{n-1}(\phi)$  by attaching a coproduct of representable presheaves of degree  $n$  along their formal boundary.

*Proof* (ii)  $\implies$  (i). By Lemmas 6.10 and 6.2, the latching object  $L_r(X)$  may be identified with the subobject of degenerate elements of  $X_r$ . Since  $\phi$  is monic, the induced map  $L_r(\phi) : L_r(X) \rightarrow L_r(Y)$  is monic; moreover,  $\phi$  takes non-degenerate elements of  $X$  to non-degenerate elements of  $Y$ , as follows from the left lifting property of split epimorphisms with respect to monomorphisms:

$$\begin{array}{ccc}
 \mathbb{R}[r] & \rightarrow & X \\
 \downarrow & \nearrow \text{dotted} & \downarrow \phi \\
 \mathbb{R}[s] & \rightarrow & Y
 \end{array}$$

In particular, the complement  $X_r \setminus L_r(X)$  is taken to the complement  $Y_r \setminus L_r(Y)$ , i.e. the relative latching map  $X_r \cup_{L_r(X)} L_r(Y) \rightarrow Y_r$  is monic, and the complement of its image may be identified with the set of non-degenerate elements of  $Y_r \setminus \phi(X)_r$ .

(i)  $\implies$  (iii). It follows from Lemmas 6.3 and 6.2 that the canonical map  $sk_{n-1}(\phi) \rightarrow sk_n(\phi)$ , evaluated at objects  $r$  of degree  $< n$ , is an isomorphism, while at objects  $r$  of degree  $n$ , it evaluates to  $X_r \cup_{L_r(X)} L_r(Y) \rightarrow Y_r$ . The latter is a free  $\text{Aut}(r)$ -extension by hypothesis. Since neither  $sk_{n-1}(\phi)$  nor  $sk_n(\phi)$  contain non-degenerate elements of degree  $> n$ , this shows that  $sk_n(\phi)$  is obtained from  $sk_{n-1}(\phi)$  by attaching, for each  $\text{Aut}(r)$ -orbit in  $sk_n(\phi)_r \setminus sk_{n-1}(\phi)_r$ , a distinct copy of  $\mathbb{R}[r]$ ; since the orbit is free, the complement  $\mathbb{R}[r] \setminus \partial \mathbb{R}[r]$  is freely attached.

(iii)  $\implies$  (ii). Since property (ii) is stable under pushout and sequential colimit, it suffices to show that the boundary inclusions  $\partial \mathbb{R}[r] \hookrightarrow \mathbb{R}[r]$  have property (ii). The only non-degenerate elements of  $\mathbb{R}[r] \setminus \partial \mathbb{R}[r]$  are the automorphisms of  $r$ ; the latter have trivial isotropy groups.

Cisinski shows the equivalence of 7.2(ii) and 7.2(iii) in a slightly different setting, cf. [10, 8.1.1, 8.1.29–35]. In the special cases  $\mathbb{R} = \Gamma$  and  $\mathbb{R} = \Omega$ , the skeletal filtration 7.2(iii) has been described by Lydakis [28] and Moerdijk–Weiss [31], respectively.

A map  $\phi : X \rightarrow Y$  in  $\widehat{\mathbb{R}}$ , fulfilling one of the equivalent conditions of Proposition 7.2, will be called a *cofibration*. Condition 7.2(i) readily implies

**Corollary 7.3** Let  $\mathbb{R}$  be an EZ-category and  $\mathcal{E}$  be a solid monoidal model category. A map of set-valued presheaves on  $\mathbb{R}$  is a cofibration if and only if the induced map in  $\mathcal{E}^{\mathbb{R}^{\text{op}}}$  is a Reedy cofibration.

**Definition 7.4** An EZ-category  $\mathbb{R}$  is called *quasi-monoidal* if the presheaf category  $\widehat{\mathbb{R}}$  carries a symmetric monoidal structure  $(\widehat{\mathbb{R}}, \square, I_\square, \tau_\square)$  such that

- (i) the bifunctor  $-\square- : \widehat{\mathbb{R}} \times \widehat{\mathbb{R}} \rightarrow \widehat{\mathbb{R}}$  preserves colimits in both variables;
- (ii) the unit  $I_\square$  is cofibrant;



(iii) for all objects  $r, s$  of  $\mathbb{R}$ , the boundary inclusions induce a pullback square

$$\begin{array}{ccc} \partial\mathbb{R}[r] \square \partial\mathbb{R}[s] & \rightarrow & \partial\mathbb{R}[r] \square \mathbb{R}[s] \\ \downarrow & & \downarrow \\ \mathbb{R}[r] \square \partial\mathbb{R}[s] & \longrightarrow & \mathbb{R}[r] \square \mathbb{R}[s] \end{array}$$

in  $\widehat{\mathbb{R}}$  consisting of cofibrations.

Since cofibrations are monic, the induced map

$$\mathbb{R}[r] \square \partial\mathbb{R}[s] \cup_{\partial\mathbb{R}[r] \square \partial\mathbb{R}[s]} \partial\mathbb{R}[r] \square \mathbb{R}[s] \rightarrow \mathbb{R}[r] \square \mathbb{R}[s]$$

is also monic, and hence, by 7.2(ii), a cofibration. It then follows from 7.2(iii) and 7.4(i) that the class of cofibrations in  $\widehat{\mathbb{R}}$  satisfies Hovey’s pushout-product axiom.

### 7.5 Convolution product

The monoidal structure  $\square$  on  $\widehat{\mathbb{R}}$  and the monoidal structure  $\otimes_{\mathcal{E}}$  on  $\mathcal{E}$  together induce a (closed symmetric) monoidal structure on  $\mathcal{E}^{\mathbb{R}^{\text{op}}}$ , which will be denoted  $\square_{\mathcal{E}}$ , and called the *convolution* of  $\square$  and  $\otimes_{\mathcal{E}}$ . The construction uses that  $\mathcal{E}$  is cocomplete and can be obtained as an enriched Kan extension, see [30, Appendix]. For objects  $X$  and  $Y$  of  $\mathcal{E}^{\mathbb{R}^{\text{op}}}$  the convolution product  $X \square_{\mathcal{E}} Y$  is explicitly given by the formula

$$(X \square_{\mathcal{E}} Y)_t = \varinjlim_{\mathbb{R}[r] \rightarrow \mathbb{R}[r] \square \mathbb{R}[s]} X_r \otimes_{\mathcal{E}} Y_s.$$

Notice that the functor  $\widehat{\mathbb{R}} \rightarrow \mathcal{E}^{\mathbb{R}^{\text{op}}}$  discussed at the beginning of this section is strongly symmetric monoidal because of 7.4(i). In particular, it preserves the unit.

**Theorem 7.6** *Let  $\mathbb{R}$  be a quasi-monoidal EZ-category and let  $\mathcal{E}$  be a cofibrantly generated, solid monoidal model category. Then the functor category  $\mathcal{E}^{\mathbb{R}^{\text{op}}}$ , equipped with the Reedy model structure of 1.6 and with the symmetric monoidal structure obtained by convolution, is a cofibrantly generated, solid monoidal model category.*

*Proof* We shall first show that  $\mathcal{E}^{\mathbb{R}^{\text{op}}}$  is cofibrantly generated. The generating (trivial) Reedy cofibrations of  $\mathcal{E}^{\mathbb{R}^{\text{op}}}$  are obtained by “twisting” the generating (trivial) cofibrations of  $\mathcal{E}$  against the boundary inclusions of  $\widehat{\mathbb{R}}$ . To be more precise, let  $f : A \rightarrow B$  be an arbitrary generating (trivial) cofibration of  $\mathcal{E}$  and let  $i_r : \partial\mathbb{R}[r] \rightarrow \mathbb{R}[r]$  be a boundary inclusion of  $\widehat{\mathbb{R}}$ . For brevity, for any object  $A$  of  $\mathcal{E}$  and any set  $S$ , the tensor  $A \otimes_{\mathcal{E}} (\coprod_S I_{\mathcal{E}})$  will be written  $A \otimes_{\mathcal{E}} S$ , and similarly for set-valued presheaves on  $\mathbb{R}$ . We thus obtain the following commutative square

$$\begin{array}{ccc} A \otimes_{\mathcal{E}} \partial\mathbb{R}[r] & \rightarrow & B \otimes_{\mathcal{E}} \partial\mathbb{R}[r] \\ \downarrow & & \downarrow \\ A \otimes_{\mathcal{E}} \mathbb{R}[r] & \longrightarrow & B \otimes_{\mathcal{E}} \mathbb{R}[r] \end{array}$$

in  $\mathcal{E}^{\mathbb{R}^{\text{op}}}$ . The induced comparison map

$$A \otimes_{\mathcal{E}} \mathbb{R}[r] \cup_{A \otimes_{\mathcal{E}} \partial\mathbb{R}[r]} B \otimes_{\mathcal{E}} \partial\mathbb{R}[r] \rightarrow B \otimes_{\mathcal{E}} \mathbb{R}[r]$$

is a *generating (trivial) Reedy cofibration* of  $\mathcal{E}^{\mathbb{R}^{\text{op}}}$ , and they are all of this form. Indeed, since the Reedy model structure on  $\mathcal{E}^{\mathbb{R}^{\text{op}}}$  is well defined by Theorem 1.6, the generating property just expresses that a map  $X \rightarrow Y$  is a trivial Reedy fibration (resp. Reedy fibration) *if and*

only if it has the right lifting property with respect to the generating Reedy cofibrations (resp. trivial Reedy cofibrations). This in turn follows from the fact that, by adjointness, one of the following two squares

$$\begin{array}{ccc}
 A \otimes_{\mathcal{E}} \mathbb{R}[r] \cup_{A \otimes_{\mathcal{E}} \partial \mathbb{R}[r]} B \otimes_{\mathcal{E}} \partial \mathbb{R}[r] & \rightarrow & X \\
 \downarrow & & \downarrow \\
 B \otimes_{\mathcal{E}} \mathbb{R}[r] & \longrightarrow & Y
 \end{array}
 \quad
 \begin{array}{ccc}
 A & \longrightarrow & X_r \\
 \downarrow & & \downarrow \\
 B & \rightarrow & Y_r \times_{M_r(Y)} M_r(X)
 \end{array}$$

has a diagonal filler if and only if the other has.

The fact that  $\mathcal{E}^{\mathbb{R}^{\text{op}}}$  is solid follows from the facts that  $\mathcal{E}$  is solid and that  $\widehat{\mathbb{R}} \rightarrow \mathcal{E}^{\mathbb{R}^{\text{op}}}$  is a strong monoidal functor preserving cofibrations, cf. Corollary 7.3.

So, it remains to check the pushout-product axiom. To this end, take two generating cofibrations  $f : A \rightarrow B$ ,  $g : C \rightarrow D$  in  $\mathcal{E}$  as well as two boundary inclusions  $i_r, i_s$  of  $\mathbb{R}$ , and consider the associated generating Reedy cofibrations

$$\begin{aligned}
 f/i_r &: A \otimes_{\mathcal{E}} \mathbb{R}[r] \cup_{A \otimes_{\mathcal{E}} \partial \mathbb{R}[r]} B \otimes_{\mathcal{E}} \partial \mathbb{R}[r] \rightarrow B \otimes_{\mathcal{E}} \mathbb{R}[r], \\
 g/i_s &: C \otimes_{\mathcal{E}} \mathbb{R}[s] \cup_{C \otimes_{\mathcal{E}} \partial \mathbb{R}[s]} D \otimes_{\mathcal{E}} \partial \mathbb{R}[s] \rightarrow D \otimes_{\mathcal{E}} \mathbb{R}[s].
 \end{aligned}$$

We shall denote them by  $f/i_r : A/i_r \rightarrow B/i_r$  and  $g/i_s : C/i_s \rightarrow D/i_s$ . We have to show that the pushout-product map

$$(A/i_r \square_{\mathcal{E}} D/i_s) \cup_{(A/i_r \square_{\mathcal{E}} C/i_s)} (B/i_r \square_{\mathcal{E}} C/i_s) \rightarrow (B/i_r \square_{\mathcal{E}} D/i_s) \tag{9}$$

is a Reedy cofibration which is trivial if  $f$  or  $g$  is trivial.

The operation  $(f, i_r) \mapsto f/i_r$  extends in an evident way to a bifunctor

$$-/- : \text{Arr}(\mathcal{E}) \times \text{Arr}(\widehat{\mathbb{R}}) \rightarrow \text{Arr}(\mathcal{E}^{\mathbb{R}^{\text{op}}}),$$

where  $\text{Arr}(\mathbb{C})$  denotes the category of arrows in  $\mathbb{C}$ .

It is now straightforward to verify that (9) is isomorphic to  $h/i_t$  where

$$\begin{aligned}
 h &: A \otimes_{\mathcal{E}} D \cup_{A \otimes_{\mathcal{E}} C} B \otimes_{\mathcal{E}} C \rightarrow B \otimes_{\mathcal{E}} D \quad \text{and} \\
 i_t &: \mathbb{R}[r] \square \partial \mathbb{R}[s] \cup_{\partial \mathbb{R}[r] \square \partial \mathbb{R}[s]} \partial \mathbb{R}[r] \square \mathbb{R}[s] \rightarrow \mathbb{R}[r] \square \mathbb{R}[s]
 \end{aligned}$$

are the canonical comparison maps. Since in  $\mathcal{E}$  and  $\widehat{\mathbb{R}}$  the pushout-product axiom holds, it remains to be shown that for a (trivial) cofibration  $h$  in  $\mathcal{E}$ , and cofibration  $i_t$  in  $\widehat{\mathbb{R}}$ , the map  $h/i_t$  is a (trivial) Reedy cofibration in  $\mathcal{E}^{\mathbb{R}^{\text{op}}}$ . By the adjointness argument given above, this holds whenever  $h$  is a generating (trivial) cofibration, and  $i_t$  a boundary inclusion; the general case reduces to this special case, since the operation  $(h, i_t) \mapsto h/i_t$  commutes with sequential colimits and retracts in each variable, and takes pushout squares in each variable to pushout squares in  $\mathcal{E}^{\mathbb{R}^{\text{op}}}$ .

*Examples 7.7* (a) The simplex category  $\Delta$  is a quasi-monoidal EZ-category for the cartesian product on  $\widehat{\Delta}$ . Therefore, the category of simplicial spaces is a monoidal model category for the cartesian product, where “space” means either compactly generated topological space or simplicial set. In the latter case, the Reedy cofibrations are precisely the monomorphisms, and the result is of course well-known. Notice that in general, even for strict EZ-categories  $\mathbb{R}$ , the exactness axiom 7.4(iii) may not be true for the cartesian product on  $\widehat{\mathbb{R}}$ .

(b) Segal’s [36] category  $\Gamma$  is a (quasi-)monoidal EZ-category for the smash product on  $\widehat{\Gamma}$ , as can be deduced from the work of Lydakis [28]. This means that the category of  $\Gamma$ -spaces, equipped with the strict model structure of Bousfield–Friedlander [8], is a monoidal model category.

- (c) The category  $\Omega$  for dendroidal sets (see 2.8) is a quasi-monoidal EZ-category for the Boardman-Vogt tensor product on  $\widehat{\Omega}$ , cf. [11, 30]. Therefore, the category of dendroidal spaces is a monoidal model category in such a way that the embedding  $i : \Delta \hookrightarrow \Omega$  induces a monoidal Quillen adjunction between simplicial spaces and dendroidal spaces. It is shown in [12] that, in complete analogy to Rezk's localization of simplicial spaces (the model structure for complete Segal spaces, cf. [6, 25, 27, 35]), there is a localization of the model category of dendroidal spaces which is Quillen equivalent to the model category of quasi-operads introduced in [11].

**Acknowledgements** The main results of this paper were presented at the CRM in Barcelona in February 2008, in the context of the Program on Homotopy Theory and Higher Categories. The actual writing of this paper was done while the second author was visiting the University of Nice in May 2008. He would like to express his gratitude to the CRM and the University of Nice for their hospitality and support. Both authors would like to thank the referee for his careful reading and useful suggestions.

**Open Access** This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

## References

1. Angeltveit, V.: Enriched Reedy categories. *Proc. Am. Math. Soc.* **136**, 2323–2332 (2008)
2. Antokoletz, E.R.: Nonabelian Algebraic Models for Classical Homotopy Types. PhD thesis, University of California at Berkeley (2008)
3. Barwick, C.: On Reedy Model Categories. arXiv:math/0708.2832
4. Batanin, M.A., Markl, M.: Crossed interval groups and operations on the Hochschild cohomology. arXiv:math/0803.2249
5. Berger, C.: Iterated wreath product of the simplex category and iterated loop spaces. *Adv. Math.* **213**, 230–270 (2007)
6. Bergner, J.E.: Three models for the homotopy theory of homotopy theories. *Topology* **46**, 397–436 (2007)
7. Bourceux, F.: Handbook of Categorical Algebra II. *Enc Math*, vol. 51. Cambridge University Press, Cambridge (1994)
8. Bousfield, A.K., Friedlander, E.M.: Homotopy theory of  $\Gamma$ -spaces, spectra, and bisimplicial sets. *Lecture Notes in Math*, vol. 658, pp. 80–130. Springer, Berlin (1978)
9. Bousfield, A.K., Kan, D.M.L.: Homotopy limits, completions, and localization. *Lecture Notes in Mathematics*, vol. 304. Springer, Berlin (1972)
10. Cisinski, D.-C.: Les préfaisceaux comme modèles des types d'homotopie. *Astérisque*, vol. 308. Soc. Math. France (2006)
11. Cisinski, D.-C., Moerdijk, I.: Dendroidal sets as a model for homotopy operads. arXiv:0809.3341
12. Cisinski, D.-C., Moerdijk, I.: Dendroidal Segal spaces and  $\infty$ -operads, 30 pp, preprint (2010)
13. Cisinski, D.-C., Moerdijk, I.: Dendroidal sets and simplicial operads (in preparation)
14. Connes, A.: Cyclic homology and functor  $Ext^n$ . *C.R. Acad. Sci. Paris* **296**, 953–958 (1983)
15. Dwyer, W.G., Hopkins, M.J., Kan, D.M.: The homotopy theory of cyclic sets. *Trans. Am. Math. Soc.* **291**, 281–289 (1985)
16. Eilenberg, S., Zilber, J.A.: Semi-simplicial complexes and singular homology. *Ann. Math.* **51**, 499–513 (1950)
17. Feigin, B.L., Tsygan B.L.: Additive  $K$ -theory. *Lecture Notes in Mathematics*, vol. 1289, pp. 67–209. Springer, Berlin (1987)
18. Fiedorowicz, Z., Loday, J.-L.: Crossed simplicial groups and their associated homology. *Trans. Am. Math. Soc.* **326**, 57–87 (1991)
19. Gabriel, P., Zisman, M.: Calculus of fractions and homotopy theory. In: *Ergebnisse der Mathematik und ihrer Grenzgebiete*, vol. 35. Springer, Berlin (1967)
20. Goerss P.F., Jardine R.J.: *Simplicial homotopy theory*. In: *Progress in Mathematics*, vol. 174, vi+510 pp. Birkhäuser, Basel (1999)
21. Haefliger, A.: Extension of complexes of groups. *Ann. Inst. Fourier* **42**, 275–311 (1992)

22. Hirschhorn, P.: *Model Categories and Their Localizations*. Math. Surveys Monogr., vol. 99. Am. Math. Soc. (2003)
23. Hovey, M.: *Model Categories*. Math. Surveys Monogr., vol. 63. Am. Math. Soc. (1999)
24. Joyal, A.: Disks, duality and  $\theta$ -categories, 6 pp, preprint (1997)
25. Joyal, A., Tierney, M.: Quasi-categories vs Segal spaces. *Contemp. Math.* **431**, 277–326 (2007)
26. Krasauskas, R.: Skew-simplicial groups. *Lithuanian Math. J.* **27**, 47–54 (1987)
27. Lurie, J.: Higher topos theory. In: *Annals of Mathematics Studies*, vol. 170, viii+925 pp. Princeton University Press, Princeton (2009)
28. Lydakis, M.: Smash products and  $\Gamma$ -spaces. *Math. Camb. Philos. Soc.* **126**, 311–328 (1999)
29. Moerdijk, I.: Orbifolds as groupoids: an introduction. *Contemp. Math.* **310**, 205–221 (2002)
30. Moerdijk, I., Weiss, I.: Dendroidal sets. *Alg. Geom. Top.* **7**, 1441–1470 (2007)
31. Moerdijk, I., Weiss, I.: On inner Kan complexes in the category of dendroidal sets. *Adv. Math.* **221**, 343–389 (2009)
32. Quillen, D.G.: *Homotopical Algebra*. Lecture Notes in Mathematics, vol. 43, iv+156 pp. Springer, Berlin (1967)
33. Quillen, D.G.: Bivariant cyclic cohomology and models for cyclic homology types. *J. Pure Appl. Algebra* **101**, 1–33 (1995)
34. Reedy, C.L.: Homotopy theories of model categories, 8 pp, preprint (1973)
35. Rezk, C.: A model for the homotopy theory of homotopy theory. *Trans. Am. Math. Soc.* **353**, 973–1007 (2003)
36. Segal, G.: Categories and cohomology theories. *Topology* **13**, 293–312 (1974)