

## MORAVA $E$ -HOMOLOGY OF BOUSFIELD-KUHN FUNCTORS ON ODD-DIMENSIONAL SPHERES

YIFEI ZHU

(Communicated by Michael A. Mandell)

ABSTRACT. As an application of Behrens and Rezk’s spectral algebra model for unstable  $v_n$ -periodic homotopy theory, we give explicit presentations for the completed  $E$ -homology of the Bousfield-Kuhn functor on odd-dimensional spheres at chromatic level 2, and compare them to the level 1 case. The latter reflects earlier work in the literature on  $K$ -theory localizations.

### 1. INTRODUCTION

The rational homotopy theory of Quillen and Sullivan studies unstable homotopy types of topological spaces modulo torsion, or equivalently, after inverting primes. Such homotopy types are computable by means of their *algebraic models*. In particular, Quillen showed that there are equivalences of homotopy categories

$$\mathrm{Ho}_{\mathbb{Q}}(\mathrm{Top}_*)_2 \simeq \mathrm{Ho}_{\mathbb{Q}}(\mathrm{DGL})_1 \simeq \mathrm{Ho}_{\mathbb{Q}}(\mathrm{DGC})_2$$

between simply-connected pointed topological spaces localized with respect to rational homotopy equivalences, connected differential graded Lie algebras over  $\mathbb{Q}$ , and simply-connected differential graded cocommutative coalgebras over  $\mathbb{Q}$  [21, Theorem I].

Let  $p$  be a prime,  $\mathbb{F}_p$  be the field with  $p$  elements, and  $\overline{\mathbb{F}}_p$  be its algebraic closure. Working prime by prime, one has  $p$ -adic analogues where equivalences detected through  $H_*(-;\mathbb{Q})$  are replaced by those through  $H_*(-;\overline{\mathbb{F}}_p)$ . Various algebraic models for  $p$ -adic homotopy types of spaces were developed (Kříž [17], Goerss [12], Mandell [20]). In the modern language of homotopy theory, these models are often formulated in terms of “spectral” algebra. For example, Mandell’s model is given by the functor that takes a connected  $p$ -complete nilpotent space  $X$  of finite  $p$ -type to the  $\overline{\mathbb{F}}_p$ -cochains  $H\overline{\mathbb{F}}_p^X$ , where  $H\overline{\mathbb{F}}_p^X$  denotes the function spectrum  $F(\Sigma^\infty X, H\overline{\mathbb{F}}_p)$ . This spectrum is a commutative algebra over  $H\overline{\mathbb{F}}_p$ .

Recently, through the prism of chromatic homotopy theory, Behrens and Rezk have established in [6] spectral algebra models for unstable  $v_n$ -periodic homotopy types (cf. Arone and Ching [4], Heuts [14], and see also [7, 1, 3, 13]). Here, instead of inverting primes, they work  $p$ -locally for a fixed prime  $p$  and invert classes of maps called “ $v_n$ -self maps” (the case of  $n = 0$  recovers rational homotopy). Correspondingly, there is the  $n$ ’th unstable monochromatic category  $M_n^f\mathrm{Top}_*$  in the

---

2010 *Mathematics Subject Classification*. Primary 55S25; Secondary 55N20, 55N34, 55Q51.

sense of Bousfield [9]. They study the functor

$$(1) \quad \mathbf{S}_{T(n)}^{(-)}: \mathrm{Ho}(M_n^f \mathrm{Top}_*)^{\mathrm{op}} \rightarrow \mathrm{Ho}(\mathrm{Alg}_{\mathrm{Comm}}(\mathrm{Sp}_{T(n)}))$$

that sends a space  $X$  to the  $\mathbf{S}_{T(n)}$ -valued cochains  $\mathbf{S}_{T(n)}^X$ . This last spectrum is an algebra for the reduced commutative operad  $\mathrm{Comm}$  in modules over  $\mathbf{S}_{T(n)}$ , the localization of the sphere spectrum with respect to the telescope of a  $v_n$ -self map.

Considering a variant of localization with respect to the Morava  $K$ -theory  $K(n)$ , Behrens and Rezk have obtained an equivalence

$$c_X^{K(n)}: \Phi_{K(n)}(X) \xrightarrow{\sim} \mathrm{TAQ}_{\mathbf{S}_{K(n)}}(\mathbf{S}_{K(n)}^X)$$

of  $K(n)$ -local spectra, on a class of spaces  $X$  including spheres [6, Theorem 8.1] (cf. [7, Section 8]). On the left-hand side,  $\Phi_{K(n)} = L_{K(n)}\Phi_n$  is a version of the Bousfield-Kuhn functor (cf. Kuhn [19]). The right-hand side is the topological André-Quillen cohomology of  $\mathbf{S}_{K(n)}^X$  as an algebra over the operad  $\mathrm{Comm}$  in  $\mathbf{S}_{K(n)}$ -modules. The map  $c_X^{K(n)}$  arises from (1) as follows. Computing homotopy groups of spaces in the source category of (1), we have a natural transformation

$$v_n^{-1}\pi_*(-; V) \cong [\Sigma^*V, M_n^f(-)]_{\mathrm{Top}_*} \rightarrow [\mathbf{S}_{T(n)}^{(-)}, \mathbf{S}_{T(n)}^{\Sigma^*V}]_{\mathrm{Alg}_{\mathrm{Comm}}(\mathrm{Sp}_{T(n)})}$$

where  $V$  is any  $p$ -local finite complex of type  $n$  with a  $v_n$ -self map  $v: \Sigma^k V \rightarrow V$  for some  $k$ ,  $v_n^{-1}\pi_*(-; V) := v^{-1}[\Sigma^*V, -]_{\mathrm{Top}_*}$  are the  $v_n$ -periodic homotopy groups with coefficients in  $V$ , and  $M_n^f(-) := \mathrm{hofib}(L_n^f(-) \rightarrow L_{n-1}^f(-))$  is the  $n$ 'th monochromatic layer. Both sides of this natural transformation can be realized by functors valued in spectra. Behrens and Rezk take a homotopy limit over suitable complexes  $V$  of such natural transformations, while replacing  $\mathbf{S}_{T(n)}$  with  $\mathbf{S}_{K(n)}$  everywhere. The resulting limit is  $c_{(-)}^{K(n)}$ , which they show is an equivalence on certain  $X$ . Via a suitable Koszul duality between  $\mathrm{Comm}$  and the Lie operad, we may view the spectrum  $\mathrm{TAQ}_{\mathbf{S}_{K(n)}}(\mathbf{S}_{K(n)}^X)$  as a Lie algebra model for the unstable  $v_n$ -periodic homotopy type of  $X$ .

**1.1. Main results.** The purpose of this paper is to make available calculations that apply Behrens and Rezk's theory to obtain quantitative information about unstable  $v_n$ -periodic homotopy types, in the case of  $n = 2$ . These are based on our computation of power operations for Morava  $E$ -theory in [31].

Let  $E$  be a Morava  $E$ -theory spectrum of height 2 with  $E_* \cong \mathbb{WF}_p[[a]][u^{\pm 1}]$ , where  $|a| = 0$  and  $|u| = -2$ . Recall that the *completed  $E$ -homology* functor is defined as  $E_*^\wedge(-) := \pi_*(E \wedge -)_{K(2)}$ . It is  $E_0$ -linear dual to  $E^*(-)$  with more convenient properties than  $E_*(-)$  (see Rezk [22, Section 3]).

Building on and strengthening Rezk's results in [24, §2.13 and §10], we obtain the following.

**Theorem 1.** *Given any non-negative integer  $m$ , denote by  $E_*^\wedge(\Phi_2 S^{2m+1})$  the completed  $E$ -homology groups of the Bousfield-Kuhn functor applied to the  $(2m+1)$ -dimensional sphere.*

- (i) *The group  $E_1^\wedge(\Phi_2 S^{2m+1}) \cong 0$  if  $m = 0$ . As an  $E_0$ -module, it equals  $(E_0/p)^{\oplus p-1}$  if  $m = 1$ . It is a quotient of  $(E_0/p^m)^{\oplus p-1} \oplus E_0/p^{m-1}$  if  $m > 1$ .*

(ii) More explicitly,

$$E_1^\wedge(\Phi_2 S^{2m+1}) \cong \begin{cases} \frac{\bigoplus_{i=1}^{p-1} (E_0/p^m) \cdot x_i \oplus (E_0/p^{m-1}) \cdot x_p}{(r_1, \dots, r_{m-1})} & \text{if } 2 \leq m \leq p+2 \\ \frac{\bigoplus_{i=1}^{p-1} (E_0/p^m) \cdot x_i \oplus (E_0/p^{m-1}) \cdot x_p}{(r_{m-p-1}, \dots, r_{m-1})} & \text{if } m > p+2 \end{cases}$$

where  $r_j = r_j(x_1, \dots, x_p) = w_0^{m-1-j} \sum_{i=1}^p d_{i,j+1} x_i$ . Here, as in Zhu [31, Theorem 1.6],

$$d_{i,\tau} = \sum_{n=0}^{\tau-1} (-1)^{\tau-n} w_0^n \sum_{\substack{m_1 + \dots + m_{\tau-n} = \tau+i \\ 1 \leq m_s \leq p+1 \\ m_{\tau-n} \geq i+1}} w_{m_1} \cdots w_{m_{\tau-n}}$$

where the coefficients  $w_i \in E_0 \cong \mathbb{WF}_p[[a]]$  are defined by the identity

$$\sum_{i=0}^{p+1} w_i b^i = (b-p)(b+(-1)^p)^p - (a-p^2+(-1)^p)b$$

in the variable  $b$ , so that  $w_{p+1} = 1$ ,  $w_1 = -a$ ,  $w_0 = (-1)^{p+1}p$ , and the remaining coefficients

$$w_i = (-1)^{p(p-i+1)} \left[ \binom{p}{i-1} + (-1)^{p+1} p \binom{p}{i} \right]$$

In particular, each relation  $r_j$  contains a term  $(-1)^{j+1} w_0^{m-1-j} w_1^j x_p$ .

(iii) The group  $E_0^\wedge(\Phi_2 S^{2m+1}) \cong 0$  for any  $m \geq 0$ .

Since  $E$  is 2-periodic, the above determines the completed  $E$ -homology in all degrees.

*Remark 2.* There is a ring structure on  $E_1^\wedge(\Phi_2 S^{2m+1})$ . Indeed, each generator  $x_i$  is a power  $b^i$  of a certain element  $b$ . See Section 3.1 for details.

*Remark 3.* In [29, Sections 5.3–5.4], Wang obtained an equivalent presentation of  $E_*^\wedge(\Phi_2 S^{2m+1})$  for  $m = 1$  and any prime  $p$ . He then used it as the input of a spectral sequence and computed  $\pi_*(\Phi_{K(2)} S^3)$  at  $p \geq 5$ .

**Example 4.** We apply Theorem 1 and compute  $E_1^\wedge(\Phi_2 S^{2m+1})$  at  $p = 2$  for small values of  $m$ . We have  $w(a, b) = b^3 - ab - 2$  so that  $w_3 = 1$ ,  $w_2 = 0$ ,  $w_1 = -a$ , and  $w_0 = -2$ .

- When  $m = 2$ , since  $r_1 = 2x_1 - ax_2$ ,  $E_1^\wedge(\Phi_2 S^5)$  is the quotient of  $(E_0/4) \cdot x_1 \oplus (E_0/2) \cdot x_2$  subject to the relation  $ax_2 = 2x_1$ .
- When  $m = 3$ , we have  $r_1 = -4x_1 + 2ax_2$  and  $r_2 = 2ax_1 - a^2x_2$  so that the relations are

$$\begin{aligned} a^2 x_2 &= 2ax_1 \\ 2ax_2 &= 4x_1 \end{aligned}$$

- When  $m = 4$ , we have  $r_1 = 8x_1 - 4ax_2$ ,  $r_2 = -4ax_1 + 2a^2x_2$ , and  $r_3 = 2a^2x_1 + (-a^3 + 4)x_2$ . Thus the relations are

$$\begin{aligned} a^3x_2 &= 2a^2x_1 + 4x_2 \\ 2a^2x_2 &= 4ax_1 \\ 4ax_2 &= 8x_1 \end{aligned}$$

- When  $m = 5 > p+2$ , we have  $r_2 = 8ax_1 - 4a^2x_2$ ,  $r_3 = -4a^2x_1 + (2a^3 - 8)x_2$ , and  $r_4 = (2a^3 - 8)x_1 + (-a^4 + 8a)x_2$ . Thus the relations are

$$\begin{aligned} a^4x_2 &= (2a^3 - 8)x_1 + 8ax_2 \\ 2a^3x_2 &= 4a^2x_1 + 8x_2 \\ 4a^2x_2 &= 8ax_1 \end{aligned}$$

The relations above show that the exponent bounds for  $p$ -power torsion in Theorem 1 are sharp, e.g.  $2^{m-1}x_1 \neq 0$  and  $2^{m-2}x_2 \neq 0$  in  $E_1^\wedge(\Phi_2 S^{2m+1})$  in these cases (see Bousfield [10, §2.5] and Selick [26]). Also, as in part (ii) of the theorem, each  $r_j$  contains a term  $2^{m-1-j}a^jx_2$ . Unfortunately, it is impossible to simplify the relations for  $E_1^\wedge(\Phi_2 S^{2m+1})$  into  $2^{m-1-j}a^jx_2 = 0$  by an  $E_0$ -linear change of variables with  $x_i$ . See Remark 9 below.

**1.2. A comparison to the case of  $n = 1$ .** As an application of Behrens and Rezk's theory, Theorem 1 is a step toward the program initiated in Arone and Mahowald [2] to compute the unstable  $v_n$ -periodic homotopy groups of spheres using stable  $v_n$ -periodic homotopy groups and Goodwillie calculus. See also Wang [28, 29]. Here we discuss a version of Theorem 1 at height  $n = 1$  according to this program.

**Proposition 5.** *Let  $E$  be a Morava  $E$ -theory spectrum of height 1, with  $E_0 \cong \mathbb{WF}_p$ . Then we have*

$$E_0^\wedge(\Phi_1 S^{2m+1}) \cong E_0/p^m \quad \text{and} \quad E_1^\wedge(\Phi_1 S^{2m+1}) \cong 0 \quad m \geq 0$$

*Proof.* This is stated in Rezk [24, §2.13], based specifically on computations in §§8.3–8.4 there.  $\square$

*Remark 6.* There has been extensive work on the case of  $v_1$ -periodic homotopy theory. See Bousfield [8, §9.11] and [10, §§8.6–8.7] for an alternative approach to the above theorem. In particular,

$$\Phi_1 S^{2m+1} \simeq \mathbf{S}_{K(1)}^{2m}/p^m$$

unless  $p = 2$  and  $m \equiv 1, 2 \pmod{4}$ , where  $\mathbf{S}_{K(1)}^{2m}/p^m$  is the  $K(1)$ -localization of the Moore spectrum with  $i$ 'th space  $S^{i+2m} \cup_{p^m} e^{i+2m+1}$ . See also Davis [11, esp. Theorem 3.1] and the references therein for related information.

Rezk's proof generalizes to height  $n = 2$ . Indeed, we will show Theorem 1 by combining his framework and our explicit formulas from [31]. It is this framework which relies on an interaction with Goodwillie calculus (see also Kuhn [18]). Specifically, the  $E$ -homology of a  $v_n$ -periodic Goodwillie tower of the identity functor evaluated on odd-spheres is isomorphic to the dual of a Koszul resolution of a Dyer-Lashof algebra for the  $E$ -theory (Behrens and Rezk [6, Theorem 9.1]).

*Remark 7.* For any  $n$ , since  $\Phi_n$  is a reduced homotopy functor (i.e.  $\Phi_n(*) \simeq *$  and  $\Phi_n$  preserves weak equivalences), there is a natural “stabilization” map

$$\Sigma^2 \Phi_n(X) \rightarrow \Phi_n(\Sigma^2 X)$$

In view of the 2-periodicity of  $E$ , this induces a map on completed  $E$ -homology in the same degree. Thus the groups  $\{E_*^\wedge(\Phi_n S^{2m+1})\}_{m \geq 0}$  form a directed system. It would be interesting to understand the colimit of this system and its implication for homotopy types. Note that completed  $E$ -homology does not send homotopy colimits to colimits in analytically complete  $E_*$ -modules in the sense of Rezk [24, §2.5] (see Hovey [15]). Nevertheless, based on computational evidence from Theorem 1, Proposition 5, and further, we hope to study the relationship between the  $K(n)$ -local sphere and the Bousfield-Kuhn functor on odd-spheres hinted in Rezk [25, §§3.20–3.21].

Indeed, consider the case of  $n = 1$  and set

$$X_m = (E \wedge \Omega^{2m+1} \Phi_1(S^{2m+1}))_{K(1)}$$

in Hovey [15, Corollary 3.5]. Therefore, in view of Proposition 5, the stabilization  $\text{hocolim}_k \Omega^{2m+1} \Phi_1(S^{2m+1})$  has  $p$ -completed  $K$ -theory isomorphic to  $\mathbb{Z}_p$ , concentrated in even degrees. On the other hand, given any  $n$ ,  $\Phi_n$  preserves filtered homotopy colimits. Thus,  $K(n)$ -locally, we have

$$\begin{aligned} \text{hocolim}_k \Omega^k \Phi_n S^k &\simeq \text{hocolim}_k \Phi_n \Omega^k S^k \\ &\simeq \Phi_n \text{hocolim}_k \Omega^k S^k \\ &\simeq \Phi_n \Omega^\infty \mathbf{S} \\ &\simeq \mathbf{S}_{K(n)} \end{aligned}$$

(cf. Remark 6 and [15, example following Corollary 3.5]).

## 2. KOSZUL COMPLEXES FOR MODULES OVER THE DYER-LASHOF ALGEBRA OF MORAVA $E$ -THEORY

Let  $E$  be a Morava  $E$ -theory spectrum of height  $n$  at the prime  $p$ . Its formal group  $\text{Spf } E^0 \mathbb{C}\mathbb{P}^\infty$  over  $E_0 \cong \mathbb{W}\bar{\mathbb{F}}_p[[u_1, \dots, u_{n-1}]]$  is the Lubin-Tate universal deformation of a formal group  $\mathbb{G}$  over  $\bar{\mathbb{F}}_p$  of height  $n$ .

Generalizing the Lubin-Tate deformation theory, Strickland shows that for each  $k \geq 0$  there is a ring  $A_k \cong E^0 B\Sigma_{p^k}/I_k$  classifying subgroups of degree  $p^k$  in the universal deformation, where  $I_k$  is the ideal generated by the image of all transfer maps from inclusions of the form  $\Sigma_i \times \Sigma_{p^k-i} \subset \Sigma_{p^k}$  with  $0 < i < p^k$  [27, Theorem 1.1]. In particular,  $A_0 \cong E_0$  and there are ring homomorphisms

$$s = s_k, t = t_k: A_0 \rightarrow A_k \quad \text{and} \quad \mu = \mu_{k,m}: A_{k+m} \rightarrow A_k {}^s \otimes_{A_0} {}^t A_m$$

classifying the source and target of an isogeny of degree  $p^k$  on the universal deformation and the composition of two isogenies.

As  $E$  is an  $E_\infty$ -ring spectrum, there are (additive) power operations acting on the homotopy of  $K(n)$ -local commutative  $E$ -algebra spectra. A  $\Gamma$ -module is a left  $A_0$ -module  $M$  equipped with structure maps (the power operations)

$$P_k: M \rightarrow {}^t A_k {}^s \otimes_{A_0} M \quad k \geq 0$$

which are a compatible family of  $A_0$ -module homomorphisms. These power operations form the *Dyer-Lashof algebra*  $\Gamma$  for the  $E$ -theory, with graded pieces

$\Gamma[k] := \text{Hom}_{A_0}({}^s A_k, A_0)$ ,  $k \geq 0$ . There is a tensor product  $\otimes$  for  $\Gamma$ -modules (Rezk [24, §4.1]).

The structure of a  $\Gamma$ -module is determined by  $P_1$ , subject to a condition involving  $A_2$ , i.e. the existence of the dashed arrow in the diagram

$$(2) \quad \begin{array}{ccc} M & \xrightarrow{P_1} & {}^t A_1 {}^s \otimes_{A_0} M \\ \downarrow & & \downarrow \text{id} \otimes P_1 \\ {}^t A_2 {}^s \otimes_{A_0} M & \xrightarrow{\mu \otimes \text{id}} & {}^t A_1 {}^s \otimes_{A_0} {}^t A_1 {}^s \otimes_{A_0} M \end{array}$$

[24, Proposition 7.2]. This manifests the fact that the ring  $\Gamma$  is *Koszul* and, in particular, *quadratic* (Rezk [23]).

Let  $D_0 := A_0$ ,  $D_1 := A_1$ , and

$$D_k := \text{coker} \left( \bigoplus_{i=0}^{k-2} A_1^{\otimes i} \otimes A_2 \otimes A_1^{k-i-2} \xrightarrow{\text{id} \otimes \mu \otimes \text{id}} A_1^{\otimes k} \right) \quad k \geq 2$$

Given  $\Gamma$ -modules  $M$  and  $N$ , Rezk defines the *Koszul complex*  $\mathcal{C}^\bullet(M, N)$  by

$$\mathcal{C}^k(M, N) := \text{Hom}_{A_0}(M, D_k \otimes_{A_0} N)$$

with appropriate coboundary maps [24, §7.3].

**Proposition 8.** *If  $M$  is projective as an  $A_0$ -module, then*

$$\text{Ext}_\Gamma^k(M, N) \cong H^k \mathcal{C}^\bullet(M, N)$$

In particular, if  $k > n$ ,  $D_k \cong 0$  and so  $\text{Ext}_\Gamma^k(M, N) \cong 0$ .

*Proof.* This is Rezk [24, Proposition 7.4].  $\square$

**2.1. The case of  $n = 2$ .** Choose a *preferred  $\mathcal{P}_N$ -model* for  $E$  in the sense of Zhu [30, Definition 3.29] so that the formal group of  $E$  is isomorphic to the formal group of a universal deformation of a supersingular elliptic curve satisfying a list of properties.

Using the theory of dual isogenies of elliptic curves, Rezk identifies that  $D_2 \cong A_1 / s(A_0)$  [24, Proposition 9.3]. He also classifies  $\Gamma$ -modules with underlying  $A_0$ -module free of rank 1 [24, Proposition 9.7]. In particular, each of them takes the form  $L_\beta$  with structure map

$$\begin{aligned} P: L_\beta &\rightarrow {}^t A_1 {}^s \otimes_{A_0} L_\beta \\ x &\mapsto \beta \otimes x \end{aligned}$$

where  $x$  is a generator for the underlying  $A_0$ -module, and  $\beta \in A_1$  is such that  $\iota(\beta) \cdot \beta \in s(A_0)$  with  $\iota(-)$  the *Atkin-Lehner involution*<sup>1</sup> (this condition on  $\beta$  corresponds to the condition in (2)). Moreover,  $L_1$  is the unit object in the symmetric monoidal category of  $\Gamma$ -modules with respect to  $\otimes$  and  $L_{\beta_1} \otimes L_{\beta_2} \cong L_{\beta_1 \beta_2}$ . Thus  $L_\beta$  is  $\otimes$ -invertible as a  $\Gamma$ -module if and only if  $\beta \in A_1^\times$ .

<sup>1</sup>Cf. Katz and Mazur [16, 11.3.1], Atkin and Lehner [5, Lemmas 7–10]. Note that  $\iota(-)$  is *not* an involution in general (see Rezk [24, §9.1 and §9.8]).

Now let  $M = L_\alpha$  and  $N = L_\beta$ . We have identifications

$$\begin{aligned} A_0 &\xrightarrow{\sim} \mathcal{C}^0(M, N) = \text{Hom}_{A_0}(M, N) & f \mapsto (x \mapsto fy) \\ A_1 &\xrightarrow{\sim} \mathcal{C}^1(M, N) = \text{Hom}_{A_0}(M, {}^t A_1 {}^s \otimes_{A_0} N) & g \mapsto (x \mapsto g \otimes y) \\ A_1/s(A_0) &\xrightarrow{\sim} \mathcal{C}^2(M, N) = \text{Hom}_{A_0}\left(M, {}^{\iota^2 s}(A_1/s(A_0)) {}^s \otimes_{A_0} N\right) & h \mapsto (x \mapsto h \otimes y) \end{aligned}$$

Thus the Koszul complex in this case is

$$A_0 \xrightarrow{d_0} A_1 \xrightarrow{d_1} A_1/s(A_0)$$

with  $d_0 f = \iota(f)\beta - f\alpha$  and  $d_1 g = \iota(g)\beta + g\iota(\alpha)$  [24, §9.18].

More explicitly, we have identifications

$$A_0 \cong \mathbb{WF}_p[[a]] \quad \text{and} \quad A_1 \cong \mathbb{WF}_p[[a, b]]/(w(a, b))$$

where

$$w(a, b) = \sum_{i=0}^{p+1} w_i b^i = (b-p)(b+(-1)^p)^p - (a-p^2+(-1)^p)b$$

(Zhu [31, Theorem 1.2]). Note that the parameters  $a$  and  $b$  are chosen as in Rezk [24, §9.15] and they correspond precisely to  $h$  and  $\alpha$  in Zhu [30, 31]. In particular, the  $\Gamma$ -module of invariant 1-forms is  $\omega = L_b$ .

*Remark 9.* As we will see in Section 4, the generators  $x_i$  in Theorem 1 (ii) depend on the choice of the parameter  $b$  for  $A_1$ . We do not know if a different choice would make the presentations simpler.

The ring homomorphism  $s: A_0 \rightarrow A_1$  is simply the inclusion of scalars, as  $A_1$  is a free left module over  $A_0$  of rank  $p+1$ . We will thus abbreviate  $s(A_0)$  as  $A_0$ . Following Rezk [24], we will also abbreviate  $\iota(x)$  as  $x'$ , which is written as  $\tilde{x}$  in Zhu [30, 31]. Note that  $w_{p+1} = 1$ ,  $p|w_i$  for  $2 \leq i \leq p$ ,  $w_1 = -a$ , and

$$(3) \quad w_0 = (-1)^{p+1}p = bb'$$

[30, (3.30)]. Also, we have

$$(4) \quad b' = -b^p - w_p b^{p-1} - \dots - w_2 b + a$$

(cf. [24, §9.15]).

### 3. COMPUTING WITH KOSZUL COMPLEXES

Recall that  $\omega = L_b$  is the  $\Gamma$ -module of invariant 1-forms defined in Section 2.1. Write  $\text{nul} := L_0$ , the  $\Gamma$ -module annihilated by  $\Gamma$ . In this section, we compute  $\text{Ext}_\Gamma^*(\omega^m, \text{nul})$  for  $m \geq 0$ . By Proposition 8,

$$\text{Ext}_\Gamma^*(\omega^m, \text{nul}) \cong H^* \mathcal{C}^\bullet(L_{b^m}, L_0)$$

where

$$\mathcal{C}^\bullet(L_{b^m}, L_0): A_0 \xrightarrow{-b^m} A_1 \xrightarrow{b'^m} A_1/A_0$$

**Proposition 10.** *For all  $m \geq 0$ ,  $H^0 \mathcal{C}^\bullet(L_{b^m}, L_0) \cong H^1 \mathcal{C}^\bullet(L_{b^m}, L_0) \cong 0$ .*

*Proof.* This is proven in Rezk [24, §10].  $\square$

Rezk also gave a description of  $H^2 \mathcal{C}^\bullet(L_{b^m}, L_0) \cong A_1/(A_0 + b'^m A_1)$  in [24, §2.13] (the general case of “ $P_m$ ” as given is not completely correct; cf. Theorem 1). In the following we will compute this second cohomology explicitly.

**3.1. The second cohomology.** Write  $B_m := H^2 \mathcal{C}^\bullet(L_{b^m}, L_0) \cong A_1/(A_0 + b'^m A_1)$ . Clearly,  $B_0 \cong 0$ . Let  $m > 0$  for the rest of this section.

As a free module over  $A_0$ , the ring  $A_1$  has a basis consisting of

$$(5) \quad 1, b, b^2, \dots, b^p$$

**Proposition 11.** *In the  $A_0$ -module  $B_m$ ,  $p^m b^i = 0$  for  $1 \leq i \leq p-1$  and  $p^{m-1} b^p = 0$ .*

*Proof.* Given (3) and (4), in  $B_m$  we have  $w_0^m b^i = b'^m b^m b^i = b'^m b^{m+i} = 0$  and

$$(6) \quad \begin{aligned} w_0^{m-1} b^p &= w_0^{m-1} (-b' - w_p b^{p-1} - \dots - w_2 b) \\ &= -b^{m-1} b'^m - w_0^{m-1} w_p b^{p-1} - \dots - w_0^{m-1} w_2 b \\ &= -w_0^{m-1} w_p b^{p-1} - \dots - w_0^{m-1} w_2 b \end{aligned}$$

Since  $p|w_i$  for  $2 \leq i \leq p$ , the last expression has a factor of  $w_0^m$  and so vanishes in  $B_m$  as we have just shown.  $\square$

Let  $1 \leq m \leq p$ . Under the map of multiplication by  $b'^m$ , the elements in (5) become

$$(7) \quad b'^m, w_0 b'^{m-1}, w_0^2 b'^{m-2}, \dots, w_0^{m-1} b', w_0^m, w_0^m b, \dots, w_0^m b^{p-m}$$

Note that  $w_0^{m-1} b' = 0$  in  $B_m$  is equivalent to (6). Thus, as a quotient of the ring  $(A_0/p^m)^{\oplus p-1} \oplus A_0/p^{m-1}$  from the above proposition,  $B_m$  has relations given precisely by the vanishing of the first  $(m-1)$  terms in (7).

To write down these  $(m-1)$  relations explicitly, we recall the formulas of  $b'^\tau$ ,  $2 \leq \tau \leq m$  from Zhu [31, Section 4.1]. There,  $b'$  is written as  $\tilde{\alpha}$ . We have

$$(8) \quad b'^\tau = d_{p,\tau} b^p + d_{p-1,\tau} b^{p-1} + \dots + d_{0,\tau} \quad 1 \leq \tau \leq p$$

where

$$d_{i,\tau} = \sum_{n=0}^{\tau-1} (-1)^{\tau-n} w_0^n \sum_{\substack{m_1 + \dots + m_{\tau-n} = \tau+i \\ 1 \leq m_s \leq p+1 \\ m_{\tau-n} \geq i+1}} w_{m_1} \cdots w_{m_{\tau-n}}$$

as in [31, Theorem 1.6 (ii)]. In particular, the formula of the coefficient  $d_{p,\tau}$  has a leading term  $(-1)^\tau w_1^{\tau-1} w_{p+1}$ . Thus the term  $w_0^{m-\tau} w_1^{\tau-1} b^p$  must appear in the relation  $w_0^{m-\tau} b'^\tau = 0$  in  $B_m$ .

Next, consider the case of  $m > p$ . Under multiplication by  $b'^m$ , (5) becomes

$$b'^m, w_0 b'^{m-1}, w_0^2 b'^{m-2}, \dots, w_0^p b'^{m-p}$$

Similarly, it remains to determine formulas for  $b'^\tau$ ,  $m-p \leq \tau \leq m$  (ignoring  $w_0^p b'^{m-p}$  if  $m=p+1$ , as above). As noted in [31, Section 4.1], the exact expression of (8) holds more generally for any  $\tau \geq 1$ , if one bears in mind the convention that  $w_\tau = 0$  whenever  $\tau > p+1$ . In fact, the computation of  $d_\tau := d_{0,\tau}$  for  $\tilde{\alpha}^\tau$  in the case  $i=0$  and  $1 \leq \tau \leq p$  from Zhu [30, proof of Proposition 6.4] goes, mutatis mutandis, for the general case.

#### 4. PROOF OF THEOREM 1

Recall that given a Morava  $E$ -theory  $E$  of height  $n$ , the completed  $E$ -homology functor is defined as  $E_*^\wedge(-) := \pi_*(E \wedge -)_{K(n)}$ . In particular,

$$(9) \quad E_*^\wedge(\Phi_n X) \cong E_*^\wedge(\Phi_{K(n)} X)$$

since the map  $\text{id} \wedge L_{K(n)} : E \wedge \Phi_n X \rightarrow E \wedge L_{K(n)} \Phi_n X$  induces a  $K(n)$ -equivalence by the Künneth isomorphism.

In [24], Rezk sets up a composite functor spectral sequence (CFSS) followed by a mapping space spectral sequence (MSSS) to compute the homotopy groups of derived mapping spaces  $\widehat{\mathcal{R}}_E(A, B)$  between  $K(n)$ -local augmented commutative  $E$ -algebras  $A$  and  $B$ . He identifies the  $E_2$ -term in the CFSS as Ext-groups over the Dyer-Lashof algebra  $\Gamma$ . The CFSS converges to the  $E_2$ -term in the MSSS.

In particular, [24, §2.13] shows that this setup specializes to compute the  $E$ -cohomology of the topological André-Quillen homology  $\text{TAQ}^{\mathbf{S}_{K(n)}}(\mathbf{S}_{K(n)}^{S^{2m+1}})$ , and that the two spectral sequences both collapse at the  $E_2$ -term when  $n = 2$ . Here  $A = E_{+}^{S^{2m+1}} := F(\Sigma_{+}^{\infty} S^{2m+1}, E)$  and  $B = E \rtimes E$  is a *square-zero extension* (see [24, §5.10]).

Now, by Behrens and Rezk [6, Theorem 8.1] and (9), we identify the abutment of the MSSS as

$$\pi_{t-s} \widehat{\mathcal{R}}_E(E_{+}^{S^{2m+1}}, E \rtimes E) \cong \pi_{t-s} F(\text{TAQ}^{\mathbf{S}_{K(2)}}(\mathbf{S}_{K(2)}^{S^{2m+1}}), E) \cong E_{t-s}^{\wedge}(\Phi_2 S^{2m+1})$$

For a fixed  $t$ , Rezk identifies the possibly nonzero terms on the  $E_2$ -page of the CFSS as  $\text{Ext}_{\text{Mod}_{\Gamma}^{\star}}^s(\omega^m, \omega^{(t-1)/2} \otimes \text{nul})$ , where  $\text{Mod}_{\Gamma}^{\star}$  is the category of  $\mathbb{Z}/2$ -graded  $\Gamma$ -modules in the sense of [24, §5.6]. Thus for degree and periodicity reasons, we may set  $t = 1$  and the calculations in Section 3 then complete the proof, with  $b^i$  written as  $x_i$  in Theorem 1.  $\square$

#### ACKNOWLEDGEMENTS

I thank Mark Behrens, Paul Goerss, Guchuan Li, Charles Rezk, Guozhen Wang, and Zhouli Xu for helpful discussions. I am especially grateful to Mark for introducing me into the program of computing unstable periodic homotopy groups of spheres and to Charles for sharing his knowledge in power operations.

I thank Lennart Meier for his helpful remarks on an earlier draft of this paper, particularly one that saved me from an error in the main results.

I thank the anonymous referee for a quick, careful reading of the paper and helpful comments and suggestions. In particular, the second paragraph of Remark 7 was included in consultation with the referee.

The paper was written and reported during my visit to the Institute of Mathematics, Chinese Academy of Sciences. I thank the institute for its hospitality.

#### REFERENCES

1. Gregory Arone, *Taylor towers in situations when Tate homology vanishes*, available at <https://www.youtube.com/watch?v=aQd-bDbL4L4>.
2. Greg Arone and Mark Mahowald, *The Goodwillie tower of the identity functor and the unstable periodic homotopy of spheres*, Invent. Math. **135** (1999), no. 3, 743–788. MR 1669268
3. Gregory Arone and Michael Ching, *A classification of Taylor towers of functors of spaces and spectra*, Adv. Math. **272** (2015), 471–552. MR 3303239
4. ———, *Localized Taylor towers*, in preparation.
5. A. O. L. Atkin and J. Lehner, *Hecke operators on  $\Gamma_0(m)$* , Math. Ann. **185** (1970), 134–160. MR 0268123
6. Mark Behrens and Charles Rezk, *The Bousfield-Kuhn functor and topological André-Quillen cohomology*, available at <http://www3.nd.edu/~mbehren1/papers/BKTAQ6.pdf> (version: 2015-06-11).
7. ———, *Spectral algebra models of unstable  $v_n$ -periodic homotopy theory*. arXiv 1703.02186

8. A. K. Bousfield, *The K-theory localizations and  $v_1$ -periodic homotopy groups of H-spaces*, Topology **38** (1999), no. 6, 1239–1264. MR 1690156
9. ———, *On the telescopic homotopy theory of spaces*, Trans. Amer. Math. Soc. **353** (2001), no. 6, 2391–2426 (electronic). MR 1814075
10. ———, *On the 2-primary  $v_1$ -periodic homotopy groups of spaces*, Topology **44** (2005), no. 2, 381–413. MR 2114954
11. Donald M. Davis, *Computing  $v_1$ -periodic homotopy groups of spheres and some compact Lie groups*, Handbook of algebraic topology, North-Holland, Amsterdam, 1995, pp. 993–1048. MR 1361905
12. Paul G. Goerss, *Simplicial chains over a field and p-local homotopy theory*, Math. Z. **220** (1995), no. 4, 523–544. MR 1363853
13. Gijs Heuts, *Goodwillie approximations to higher categories*. arXiv 1510.03304
14. ———, *Periodicity in unstable homotopy theory*, in preparation.
15. Mark Hovey, *Morava E-theory of filtered colimits*, Trans. Amer. Math. Soc. **360** (2008), no. 1, 369–382. MR 2342007
16. Nicholas M. Katz and Barry Mazur, *Arithmetic moduli of elliptic curves*, Annals of Mathematics Studies, vol. 108, Princeton University Press, Princeton, NJ, 1985. MR 772569
17. Igor Kríž, *p-adic homotopy theory*, Topology Appl. **52** (1993), no. 3, 279–308. MR 1243609
18. Nicholas J. Kuhn, *Goodwillie towers and chromatic homotopy: an overview*, Proceedings of the Nishida Fest (Kinosaki 2003), Geom. Topol. Monogr., vol. 10, Geom. Topol. Publ., Coventry, 2007, pp. 245–279. MR 2402789
19. ———, *A guide to telescopic functors*, Homology, Homotopy Appl. **10** (2008), no. 3, 291–319. MR 2475626
20. Michael A. Mandell,  *$E_\infty$  algebras and p-adic homotopy theory*, Topology **40** (2001), no. 1, 43–94. MR 1791268
21. Daniel Quillen, *Rational homotopy theory*, Ann. of Math. (2) **90** (1969), 205–295. MR 0258031
22. Charles Rezk, *The congruence criterion for power operations in Morava E-theory*, Homology Homotopy Appl. **11** (2009), no. 2, 327–379. MR 2591924
23. ———, *Rings of power operations for Morava E-theories are Koszul*. arXiv 1204.4831
24. ———, *Power operations in Morava E-theory: structure and calculations*, available at <http://www.math.uiuc.edu/~rezk/power-ops-ht-2.pdf> (version: 2013-09-13).
25. ———, *Elliptic cohomology and elliptic curves*, Felix Klein Lectures, Bonn 2015, available at <http://www.math.uiuc.edu/~rezk/felix-klein-lectures-notes.pdf> (version: 2016-06-06).
26. Paul Selick, *Moore conjectures*, Algebraic topology—rational homotopy (Louvain-la-Neuve, 1986), Lecture Notes in Math., vol. 1318, Springer, Berlin, 1988, pp. 219–227. MR 952582
27. N. P. Strickland, *Morava E-theory of symmetric groups*, Topology **37** (1998), no. 4, 757–779. MR 1607736
28. Guozhen Wang, *The monochromatic Hopf invariant*. arXiv 1410.7292
29. ———, *Unstable chromatic homotopy theory*, ProQuest LLC, Ann Arbor, MI, 2015, Thesis (Ph.D.)—Massachusetts Institute of Technology. MR 3427198
30. Yifei Zhu, *The Hecke algebra action on Morava E-theory of height 2*. arXiv 1505.06377
31. ———, *Modular equations for Lubin-Tate formal groups at chromatic level 2*. arXiv 1508.03358

DEPARTMENT OF MATHEMATICS, SOUTHERN UNIVERSITY OF SCIENCE AND TECHNOLOGY, SHENZHEN, GUANGDONG 518055, CHINA

*E-mail address:* zhuyf@sustc.edu.cn