

# FOURIER BASES AND A DISTANCE PROBLEM OF ERDŐS

ALEX IOSEVICH, NETS KATZ, AND STEEN PEDERSEN

ABSTRACT. We prove that no ball admits a non-harmonic orthogonal basis of exponentials. We use a combinatorial result, originally studied by Erdős, which says that the number of distances determined by  $n$  points in  $\mathbb{R}^d$  is at least  $C_d n^{\frac{1}{d} + \epsilon_d}$ ,  $\epsilon_d > 0$ .

## INTRODUCTION AND STATEMENT OF RESULTS

**Fourier bases.** Let  $D$  be a domain in  $\mathbb{R}^d$ , i.e.,  $D$  is a Lebesgue measurable subset of  $\mathbb{R}^d$  with finite non-zero Lebesgue measure. We say that  $D$  is a *spectral set* if  $L^2(D)$  has orthogonal basis of the form  $E_\Lambda = \{e^{2\pi i x \cdot \lambda}\}_{\lambda \in \Lambda}$ , where  $\Lambda$  is an infinite subset of  $\mathbb{R}^d$ . We shall refer to  $\Lambda$  as a *spectrum* for  $D$ .

We say that a family  $D + t$ ,  $t \in T$ , of translates of a domain  $D$  *tiles*  $\mathbb{R}^d$  if  $\cup_{t \in T} (D + t)$  is a partition of  $\mathbb{R}^d$  up to sets of Lebesgue measure zero.

**Conjecture.** It has been conjectured (see [Fug]) that a domain  $D$  is a spectral set if and only if it is possible to tile  $\mathbb{R}^d$  by a family of translates of  $D$ .

This conjecture is nowhere near resolution, even in dimension one. It has been the subject of recent research, see for example [JoPe2], [LaWa], and [Ped].

In this paper we address the following special case of the conjecture. Let  $B_d = \{x \in \mathbb{R}^d : |x| \leq 1\}$  denote the unit ball. We prove that

**Theorem 1.** *An affine image of  $D = B_d$ ,  $d \geq 2$ , is not a spectral set.*

If  $A$  is a (possibly unbounded) self-adjoint operator acting on some Hilbert space, then we may define  $\exp(-\sqrt{-1}A)$  using the Spectral Theorem. We say that two (unbounded) self-adjoint operators  $A$  and  $B$  acting on the same Hilbert space *commute* if the bounded unitary operators  $\exp(-\sqrt{-1}sA)$  and  $\exp(-\sqrt{-1}tB)$  commute for all real numbers  $s$  and  $t$ . See, for example, [ReSi] for more details on the needed operator theory. As an immediate consequence of [Fug] and Theorem 1 we have:

---

1991 *Mathematics Subject Classification.* 42B.

Research supported in part by NSF grants DMS97-06825 and DMS-9801410

**Corollary.** *There do not exist commuting self-adjoint operators  $H_j$  acting on  $L^2(B_d)$  such that  $H_j f = -\sqrt{-1} \partial f / \partial x_j$  for  $f$  in the domain of the unbounded operator  $H_j$  and  $1 \leq j \leq d$ . The derivatives  $\partial / \partial x_j$  act on  $L^2(B_d)$  in the distribution sense.*

In other words, there do not exist commuting self-adjoint restrictions of the partial derivative operators  $-\sqrt{-1} \partial / \partial x_j$ ,  $j = 1, \dots, d$ , acting on  $L^2(B_d)$  in the distribution sense.

The two-dimensional case of Theorem 1 was proved by Fuglede in [Fug]. Our proof uses the following combinatorial result. See for example [AgPa], Theorem 12.13.

**Theorem 2.** *Let  $g_d(n)$ ,  $d \geq 2$ , denote the minimum number of distances determined by  $n$  points in  $\mathbb{R}^d$ . Then*

$$(*) \quad g_d(n) \geq C_d n^{\frac{3}{3d-2}}.$$

*Remark.* The study of the problem addressed in Theorem 2 was initiated by Erdős. He proved that  $g_2(n) \geq Cn^{\frac{1}{2}}$ . See [Erd]. Moser proved in [Mos] that  $g_2(n) \geq Cn^{\frac{2}{3}}$ . More recently, Chung, Szeremedi, and Trotter proved that  $g_2(n) \geq C \frac{n^{\frac{4}{5}}}{\log^c(n)}$  for some  $c > 0$ . See [CST]. Theorem 2 above is proved by induction using the  $g_2(n) \geq Cn^{\frac{3}{4}}$  result proved by Clarkson et al. in [C].

As the reader shall see, Theorem 1 does not require the full strength of Theorem 2. We just need the fact  $g_d(n) \geq C_d n^{\frac{1}{d} + \epsilon}$ , for some  $\epsilon > 0$ .

It is interesting to contrast the case of the ball with the case of the cube  $[0, 1]^d$ . It was proved in [IoPe1], (and, independently, in [LRW]; for  $d \leq 3$  this was established in [JoPe2]), that  $\Lambda$  is a spectrum for  $[0, 1]^d$ , in the sense defined above, if and only if  $\Lambda$  is a tiling set for  $[0, 1]^d$ , in the sense that  $[0, 1]^d + \Lambda = \mathbb{R}^d$  without overlaps. It follows that  $[0, 1]^d$  has lots of spectra. The standard integer lattice  $\Lambda = \mathbb{Z}^d$  is an example, though there are many non-trivial examples as well. See [IoPe1] and [LaSh].

Our method of proof is as follows. We shall argue that if  $B_d$  were a spectral set, then any corresponding spectrum  $\Lambda$  would have the property  $\#\{\Lambda \cap B_d(R)\} \approx R^d$ , where  $B_d(R)$  denotes a ball of radius  $R$  and  $f(R) \approx g(R)$  means that there exist constants  $c \leq C$  so that  $c f(R) \leq g(R) \leq C f(R)$  for  $R$  sufficiently large. On the other hand, we will show that the number of distinct distances between the elements of  $\{\Lambda \cap B_d(R)\}$  is  $\approx R$ . Theorem 2 implies that if  $R$  is sufficiently large, this is not possible.

Kolountzakis ([Kol]) recently proved that if  $D$  is any convex non-symmetric domain in  $\mathbb{R}^d$ , then  $D$  is not a spectral set. Theorem 1 is a step in the direction of proving that if  $D$  is a convex domain such that  $\partial D$  has at least one point where the Gaussian curvature does not vanish, then  $D$  is not a spectral set. This, in its turn, would be a step towards proving the conjecture of Fuglede mentioned above.

## ORTHOGONALITY

For a domain  $D$  let

$$Z_D = \{\xi \in \mathbb{R} : \widehat{\chi}_D(\xi) = 0\}.$$

Consider a set of exponentials  $E_\Lambda$ . Observe that

$$\widehat{\chi}_D(\lambda - \lambda') = \int_D e_\lambda(x) \overline{e_{\lambda'}(x)} dx.$$

It follows that the exponentials  $E_\Lambda$  are orthogonal in  $L^2(D)$  iff

$$\Lambda - \Lambda \subseteq Z_D \cup \{0\}.$$

**Proposition 1.** *If  $E_\Lambda$  is an orthogonal subset of  $L^2(D)$  then there exists a constant  $C$  depending only on  $D$  such that*

$$\#(\Lambda \cap B_d(R)) \leq C R^d$$

for any ball  $B_d(R)$  of radius  $R$  in  $\mathbb{R}^d$ .

*Proof.* Since  $\widehat{\chi}_D$  is continuous and  $\widehat{\chi}_D(0) = |D|$  it follows that

$$\inf\{|\xi| : \widehat{\chi}_D(\xi) = 0\} = r > 0.$$

If  $\xi_1, \dots, \xi_n$  are in  $\Lambda \cap B_d(R)$  then the balls  $B(\xi_j, r/2)$  are disjoint and contained in  $B_d(R + r/2)$ . Since  $r$  only depends on  $D$  the desired inequality follows.

To study the exact possibilities for sets  $\Lambda$  so that  $E_\Lambda$  is orthogonal it is of interest to us to compute the set  $Z_D$ . We will without loss of generality assume that  $0 \in \Lambda$ . We again compare the sets  $Z_D$  for the cases where  $D$  is the cube and the ball.

Let  $Q_d = [0, 1]^d$  be the cube in  $\mathbb{R}^d$ . The zero set  $Z_Q$  for  $\widehat{\chi}_Q$  is the union of the hyperplanes  $\{x \in \mathbb{R}^d : x_i = z\}$ , where the union is taken over  $1 \leq i \leq d$ , and over all non-zero integers  $z$ .

Let  $B_d = \{x \in \mathbb{R}^d : \|x\| \leq 1\}$  be the unit ball in  $\mathbb{R}^d$ . The zero set  $Z_{B_d}$  for  $\widehat{\chi}_B$  is the union of the spheres  $\{x \in \mathbb{R}^d : \|x\| = r\}$ , where the union is over all the positive roots  $r$  of an appropriate Bessel function.

For the cube  $Q_d$  it is easy to find a large set  $\Lambda \subseteq Z_{Q_d} \cup \{0\}$  so that  $\Lambda - \Lambda \subseteq Z_{Q_d} \cup \{0\}$ . For example, we may take  $\Lambda = \mathbb{Z}^d$ . In the case of the ball  $B_d$ , we will show that only relatively small sets  $\Lambda \subseteq Z_{B_d} \cup \{0\}$  satisfy  $\Lambda - \Lambda \subseteq Z_{B_d} \cup \{0\}$ .

#### PROOF OF THEOREM 1

We shall need the following result.

**Theorem 3.** *Suppose that  $D$  is a spectral set and that  $\Lambda$  is a spectrum for  $D$  in the sense defined above, where  $D$  is a bounded domain. There exists an  $r > 0$  so that any ball of radius  $r$  contains at least one point from  $\Lambda$ .*

*Proof.* This is a special case of [IoPe2]. See also [Beu], [Lan], and [GrRa].

It is a consequence of Theorem 3 that if  $D$  is a spectral set then there exists a constant  $C > 0$  such that if  $\Lambda$  is a spectrum for  $D$  then  $\#\{\Lambda \cap B_d(R)\} \geq C R^d$  for any ball  $B_d(R)$

of radius  $R$  provided that  $R$  is sufficiently large. Combining this with Proposition 1 we see that  $\#\{\Lambda \cap B_d(R)\} \approx R^d$ .

Suppose  $\Lambda$  is a spectrum for the unit ball  $B_d$  centered at the origin in  $\mathbb{R}^d$ . Let  $B_d(R)$  be a ball of radius  $R$ . Since  $\#\{\Lambda \cap B_d(R)\} \approx R^d$  it follows from Theorem 2 that

$$(**) \quad \#\{|\lambda - \lambda'| : \lambda, \lambda' \in \Lambda \cap B_d(R)\} \geq C R^{\frac{3d}{3d-2}}.$$

Now, since  $\widehat{\chi}_{B_d}$  is an analytic radial function, it follows that if  $f$  is given by  $f(|\xi|) = \widehat{\chi}_{B_d}(\xi)$ , then the number of zeros of  $f$  in the interval  $[-R, R]$  is bounded above by a multiple of  $R$ . In fact an explicit calculation shows that  $\widehat{\chi}_{B_d}(\xi) = |\xi|^{\frac{d}{2}} J_{\frac{d}{2}}(2\pi|\xi|)$ , where  $J_\nu$  denotes the usual Bessel function of order  $\nu$ . See, for example, [BCT, p. 265].

If  $\lambda, \lambda' \in \Lambda$  then

$$f(|\lambda - \lambda'|) = \widehat{\chi}_{B_d}(\lambda - \lambda') = 0.$$

Combining the upper bound on the number of zeros of  $f$  in  $[-R, R]$  with the lower bound (\*\*\*) we derived from Theorem 2 above we have

$$C' R \geq \#\{|\lambda - \lambda'| : \lambda, \lambda' \in \Lambda \cap B_d(R)\} \geq C R^{\frac{3d}{3d-2}}.$$

Since  $1 < \frac{3d}{3d-2}$  this leads to a contradiction by choosing  $R$  sufficiently large. This completes the proof of Theorem 1.

#### REFERENCES

- [AgPa] P. Agarwal and J. Pach, *Combinatorial Geometry*, Wiley-Interscience Series, 1995.
- [Beu] A. Beurling, *Local harmonic analysis with some applications to differential operators*, Some Recent Advances in the Basic Sciences, Academic Press **1** (1966).
- [BCT] L. Brandolini, L. Colzani, and G. Travaglini, *Average decay of Fourier transforms and integer points in polyhedra*, Ark. Mat. **35** (1997), 253-275.
- [CST] F. Chung, E. Szeremedi, and W. Trotter, *The number of distinct distances determined by a set of points in the Euclidean plane*, Discrete and Computational Geometry **7** (1992).
- [C] K. Clarkson, H. Edelsbrunner, L. Guibas, M. Sharir, and E. Welzl, *Combinatorial complexity bounds for arrangements of curves and surfaces*, Discrete and Computational Geometry **5** (1990).
- [Erd] P. Erdős, *On sets of distances of  $n$  points*, American Mathematical Monthly **53** (1946).
- [Fug] B. Fuglede, *Commuting self-adjoint partial differential operators and a group theoretic problem*, J. Funct. Anal. **16** (1974), 101-121.
- [GrRa] K. Gröchenig and H. Razafinjato, *On Landau's necessary density conditions for sampling and interpolation of band-limited functions*, J. London. Math. Soc. **54** (1996), 557-565.

- [IoPe1] A. Iosevich and S. Pedersen, *Spectral and tiling properties of the unit cube*, Internat. Math Research Notices **16** (1998), 819-828.
- [IoPe2] A. Iosevich and S. Pedersen, *How large are the spectral gaps?*, Pacific J. Math. (to appear) (1998).
- [JoPe1] P. E. T. Jorgensen and S. Pedersen, *Spectral pairs in Cartesian coordinates*, J. Fourier Anal. Appl. (to appear) (1998).
- [JoPe2] P. E. T. Jorgensen and S. Pedersen, *Orthogonal harmonic analysis of fractal measures*, ERA Amer. Math. Soc. **4** (1998), 35-42.
- [Kol] M. Kolountzakis, *Non-symmetric convex domains have no basis of exponentials*, (preprint) (1999).
- [LRW] J. Lagarias, J. Reed, and Y. Wang, *Orthonormal bases of exponentials for the  $n$ -cube*, (preprint) (1998).
- [LaSh] J. Lagarias and P. Shor, *Keller's cube tiling conjecture is false in high dimensions*, Bull. Amer. Math. Soc. **27** (1992), 279-283.
- [LaWa] J. Lagarias and Y. Wang, *Spectral sets and factorizations of finite abelian groups*, J. Funct. Anal. **145** (1997), 73-98.
- [Lan] H. Landau, *Necessary density conditions for sampling and interpolation of certain entire functions*, Acta Math. **117** (1967), 37-52.
- [Mon] H. Montgomery, *Ten lectures on the interface between analytic number theory and harmonic analysis*, CBMS Regional Conference Series in Mathematics (1994).
- [Mos] L. Moser, *On different distances determined by  $n$  points*, American Mathematical Monthly **59** (1952).
- [Ped] S. Pedersen, *Spectral sets whose spectrum is a lattice with a base*, J. Funct. Anal. **141** (1992), 496-509.
- [ReSi] M. Reed and B. Simon, *Modern Mathematical Physics, Vol. I: Functional Analysis*, Academic Press, 1972.

DEPARTMENT OF MATHEMATICS, GEORGETOWN UNIVERSITY, WASHINGTON, DC 20057, USA  
*E-mail address:* `iosevich@math.georgetown.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS AT CHICAGO, CHICAGO, ILLINOIS, 60607  
*E-mail address:* `nets@math.uic.edu`

DEPARTMENT OF MATHEMATICS, WRIGHT STATE UNIVERSITY, DAYTON, OH 45435, USA  
*E-mail address:* `steen@math.wright.edu`