

We will use the following (proof done earlier) to calculate rings of integers.

Recall the following two propositions

Proposition 12.1. *Let A be Dedekind. Let \mathfrak{p} be a maximal ideal of A and let α be an integral element of a finite separable extension of the field of fractions of A . Suppose that G is the minimal monic for α over A and that the reduction mod \mathfrak{p} of G , which we call \bar{G} factors as*

$$\bar{G} = \bar{g}_1^{r_1} \cdots \bar{g}_m^{r_m},$$

with the \bar{g}_i distinct, irreducible, and monic. Then choosing monic $g_i \in A[x]$ such that $g_i \equiv \bar{g}_i \pmod{\mathfrak{p}}$, we have

- (1) $\mathfrak{q}_i = A[\alpha](g_i(\alpha), \mathfrak{p})$ is a prime for each i ; and
- (2) r_i is the smallest positive integer such that

$$R_{\mathfrak{q}_i}(\mathfrak{q}_i)^{r_i} \subseteq R_{\mathfrak{q}_i}\mathfrak{p}.$$

Proposition 12.2. *With notation as above, if $r_i = 1$ then the prime $A[\alpha](\mathfrak{p}, g_i(\alpha))$ is invertible. If $r_i > 1$, then \mathfrak{q}_i is not invertible if and only if all the coefficients of the remainder mod g_i of G are in \mathfrak{p}^2 , i.e. if writing*

$$G(x) = q(x)g_i(x) + r(x),$$

we have $r(x) \in \mathfrak{p}^2[x]$.

Now back to discriminants It is easy to see that $\Delta(F) \in K$. To see this, note that if the roots of F are distinct, then $K(\alpha_1, \dots, \alpha_n)$ is Galois over K and $\prod_{i \neq j} (\alpha_i - \alpha_j)$ is certainly invariant under the Galois group of $K(\alpha_1, \dots, \alpha_n)$ over K . It follows that $\Delta(F) \in K$. To see this, note that if the roots of F are distinct, then $K(\alpha_1, \dots, \alpha_n)$ is Galois over K and $\prod_{i \neq j} (\alpha_i - \alpha_j)$ is certainly invariant under the Galois group of $K(\alpha_1, \dots, \alpha_n)$ over K .

Here are some other, often easier ways of writing the discriminant...

Let F be monic over K . Then

$$\Delta(F) = (-1)^{n(n-1)/2} \prod_{i=1}^n F'(\alpha_i).$$

This is quite easy to see, since if $F(X) = \prod_{i=1}^n (X - \alpha_i)$, then by the product rule, $F'(X) = \sum_{i=1}^m \prod_{i \neq j} (\alpha_i - \alpha_j)$, so $F'(\alpha_i) = \prod_{j \neq i} (\alpha_i - \alpha_j)$ and $\prod_{i=1}^n F'(\alpha_i) = \prod_{i \neq j} (\alpha_i - \alpha_j)$.

When F is monic and irreducible with and $L = K(\alpha)$ is separable for a root α of F , this yields

$$\Delta(F) = (-1)^{n(n-1)/2} N_{L/K}(F'(\alpha)).$$

Since F' has coefficients in K , we see that if $\alpha_1, \dots, \alpha_n$ are the conjugates of α , then $N_{L/K}(F'(\alpha)) = \prod_{i=1}^m F'(\alpha_i)$ and we are done.

Recall this key fact from last time:

Corollary 12.3. *Let A be a Dedekind domain with field of fractions K and let \mathfrak{p} be a maximal prime in A and suppose that $A/\mathfrak{p} = k$ is a perfect field. Then the reduction \bar{F} of F modulo \mathfrak{p} has distinct roots in the algebraic closure of A/\mathfrak{p} if and only if $\Delta(F) \notin \mathfrak{p}$.*

Let's do some examples of Dedekind domains today. We'll start with $\mathbb{Q}(\sqrt[3]{5})$, which we will show is Dedekind. First of all, we'll calculate the discriminant of $\mathbb{Z}[\sqrt[3]{5}]$. We see that the minimal polynomial of $\sqrt[3]{5}$ is $F(X) = X^3 - 5$, which has derivative $3X^2$, so

$$\Delta(F) = N_{\mathbb{Q}(\sqrt[3]{5})/\mathbb{Q}}(F'(\sqrt[3]{5})) = N_{\mathbb{Q}(\sqrt[3]{5})/\mathbb{Q}}(3\sqrt[3]{5}^2) = 3^3 5^2,$$

so we know that any non-invertible primes must lie over 3 or 5, since a prime $(\mathcal{Q}, g_i(\sqrt[3]{5}))$ can fail to be invertible if and only if $g^2 \mid F \pmod{p\mathbb{Z}}$ where $\mathcal{Q} \cap \mathbb{Z} = p\mathbb{Z}$.

Let's factor over 5 and see what happens... We get $X^3 - 5 \equiv X^3 \pmod{5}$, so we get the prime $(\sqrt[3]{5}, 5)$ which is certainly generated by $\sqrt[3]{5}$ and hence is principal and thus invertible. Over 3, things are a bit more complicated. We factor as $X^3 - 5 \equiv (X - 5)^3 \pmod{3}$, so we have the ideal $(\sqrt[3]{5} - 5, 3)$, which we denote as \mathcal{Q} . How can we tell whether or not this is locally principal? Let's recall a bit about the norm.

One way to check if an integer n is in the ideal generated by an element β in an integral extension ring is to see if n is the ideal generated by the norm of β . Let's apply this idea to the above we see that

$$N_{\mathbb{Q}(\sqrt[3]{5})/\mathbb{Q}}(\sqrt[3]{5} - 5) = (1 - \sqrt[3]{5})(1 + \sqrt[3]{5} + \sqrt[3]{5}^2) = 5 - 125 = -120 = (-40) \cdot 3.$$

Since -40 is unit in $\mathbb{Z}[\sqrt[3]{5}]_{\mathcal{Q}}$, it follows that

$$\mathbb{Z}[\sqrt[3]{5}]_{\mathcal{Q}}(\sqrt[3]{5} - 5) = \mathbb{Z}[\sqrt[3]{5}]_{\mathcal{Q}}\mathcal{Q},$$

so \mathcal{Q} is locally principal, as desired. Thus, we see that $\mathbb{Z}[\sqrt[3]{5}]$ is a Dedekind domain as desired.

What about $\mathbb{Z}[\sqrt[3]{19}]$? Calculating the discriminant yields $3^3 \cdot 19^2$. Again, it is easy to see that the prime lying over 19 is just $\sqrt[3]{19}$. But the prime lying over 3 is trickier. We see that the only prime $\mathcal{Q} \in \mathbb{Z}[\sqrt[3]{19}]$ such that $\mathcal{Q} \cap \mathbb{Z} = 3\mathbb{Z}$ is the prime $(\sqrt[3]{19} - 19, 3)$. Modulo 3 we have

$$(X - 19)^3 = X - 19 \pmod{3}.$$

From some work from last time, $(\sqrt[3]{19} - 19, 3)$ is invertible if and only if the remainder of $X^3 - 19$ modulo $X - 19$ is divisible by 3^2 . We see that

$$(X^3 - 19) = (X - 19)(X^2 + 19X + 19^2) + 19^3 - 19.$$

Since

$$19^3 - 19 \equiv -18 \pmod{9} \equiv 0 \pmod{19}$$

we see that $(\sqrt[3]{19} - 19, 3)$ is not invertible.

In fact, we can generalize this to show that if a is a square-free integer and p is a prime, then $\mathbb{Z}[\sqrt[p]{a}]$ is Dedekind if and only if $a^p - a \not\equiv 0 \pmod{p^2}$. This will be on your homework.

For an element $\alpha \notin A$ that is integral over A , we define the discriminant $\Delta(\alpha/A)$ to be $\Delta(F)$ where F is the minimal monic for α over A . We also define the discriminant $\Delta(A[\alpha])$ to be $\Delta(A[\alpha])$.

Given a Dedekind domain A with field of fractions K and a finite separable extension L of K of degree n we want to be able to define a discriminant $\Delta(B'/A)$ of *any* subring B' of L . This will involve working with a basis for L over K that consists entirely of elements contained in B'

A bit more on subrings of the integral closure.

Proposition 12.4. *Let A be an integral domain with field of fractions K and let L be a finite extension of K . Suppose that $B' \subset L$ has field of fractions L and is integral over A . Then, for every element $y \in L$ there exists $a \in A$ such that $ay \in B'$.*

Proof. Let $y = \alpha/\beta$ for $\alpha, \beta \in B'$ with $\alpha, \beta \neq 0$. We will show that $\alpha/\beta = b/a$ for $b \in B'$ and $a \in A$. We know that the ideal $B'\beta$ has nonzero intersection with A by taking the constant term of the minimal monic polynomial for β over A . Thus, we can write $\gamma\beta = a$ for some nonzero $a \in A$. Then $1/\beta = \gamma/a$, so $\alpha/\beta = \alpha\gamma/a$ and we are done, since this means that $a(\alpha/\beta) \in B'$. \square

For the rest of class, A is Dedekind with field of fractions K , the field L is a finite separable extension of K of degree n , and B' is a subring of L that is integral over A . We will also assume that for every maximal ideal \mathfrak{p} of A , the residue field A/\mathfrak{p} is perfect.

We'll begin with a definition that works when B' is a free A -module, i.e. when B' is isomorphic as an A -module to A^n , where $n = [L : K]$. In this case, we choose a basis w_1, \dots, w_n for B' over A and we let M be the matrix $[m_{ij}]$ where $m_{ij} = \text{T}_{L/K}(w_i w_j)$. Then we define

$$(1) \quad \Delta(B') = \det M.$$

How do we know that this agrees with our earlier definition in the case $B' = A[\alpha]$? In fact, it more or less follows from some earlier work we did. Recall that in this case, we can choose the basis $1, \alpha, \dots, \alpha^{n-1}$, so that $[m_{ij}] = [\text{T}_{L/K}(\alpha^{i+j-2})]$, which we recall is equal to

$$\sum_{\ell=1}^n \alpha_\ell^{i+j-2}.$$

As we saw earlier, letting N be the van der Monde matrix

$$\begin{pmatrix} 1 & \cdots & 1 \\ \alpha_1 & \cdots & \alpha_n \\ \cdots & \cdots & \cdots \\ \alpha_1^{n-1} & \cdots & \alpha_n^{n-1} \end{pmatrix},$$

we have $NN^t = M$, so

$$\det M = (\det N)^2 = \prod_{i < j} (\alpha_i - \alpha_j)^2,$$

which is the same as $\Delta(\alpha)$, so our definitions agree.

Not all B' will be free A -modules, however, so we have the more general definition below.

Definition 12.5. With notation as above $\Delta(B'/A)$ is defined to be ideal generated by the determinants of all matrices $M = [\text{T}_{L/K}(w_i w_j)]$ as w_1, \dots, w_n range over all bases for L consisting of elements contained in B' .